

***Maypaisomys insularis* (Hutterer, López-Martínez & Michaux 1988) and *Haliaeetus* sp. Comments on the terrestrial fauna of the Pleistocene of the Jandía Isthmus, Fuerteventura**

Betancort JF¹, Lmoschitz A², Hernández-Acosta N³, López O¹, Gallardo A⁴ & López Jurado LF¹

¹ Departamento de Biología, ² IOCAG (Instituto de Oceanografía y Cambio Global, ³ PAMEV, Biblioteca Universitaria.

Universidad de Las Palmas de Gran Canaria, Spain.

⁴ Reserva de la Biosfera de Fuerteventura

Along the Jandia Isthmus, on Southern Fuerteventura Island, Pleistocene fossil dunes and paleosoils are particularly well developed. They are above Miocene lava flows and dikes and a Pliocene marine conglomerate. On those fossil dunes and in the windward coast (Laderas del Veril site) has been found remains of: the extinct “lava mouse” *Maypaisomys insularis* (Hutterer, López-Martínez & Michaux 1988); osprey (*Haliaeetus* sp.); seabird (*Puffinus holei* Walker, Wragg y Harrison 1990); terrestrial gastropods and locust oothecas.



Fossil remains, egg and bones, of seabirds (*Puffinus holei*)




Location of Laderas del Veril



General view of Laderas del Veril

Besides, in the close site of Hueso del Caballo, three dunes are distinguished and the upper one contains fossil remains similar to those of Laderas del Veril. There has been, radiocarbon dated giving an age of > 40 ka, then attributed to the upper Pleistocene.




Fossils remains of *Puffinus holei* (a,b &c)



Claw of *Haliaeetus* sp.



Fossils remains in Ladera del Veril



Jaws of *Maypaisomys insularis*

These fossil dunes looks like a complex fossil ecosystem being the nesting area of those seabirds and where the colonies of mouses where developed. Finally, a great osprey occupied the roll of super predator within this ecosystem.

References

- Hutterer R., Lopez-Martínez N. and Michaux J., (1988) A new rodent from quaternary deposits of the Canary Island and its relationships with Neogene and Recent Murids of Europe and Africa. *Palaeovertebrata* 18(4):241-262, 7 figs., 5 tabl., 2 pl.
- Meco J., Betancort J.F., Ballester J., Fontugne M., Guillou H., Scaillet S., Lmoschitz A., Cilleros A., Carracedo J.C., Petit-Maire N., Ramos A.J.G., Perera M.A., Soler-Onis E., Medina P., Montesinos M., Meco J.M. (2008) Historia Geológica del Clima en Canarias. 296 pp.
- Meco J., Petit-Maire N., Ballester J., Betancort J.F., Ramos A.J.G. (2010) The Acridian plagues, a new Holocene and Pleistocene palaeoclimatic indicator. *Global and Planetary Change*. Vol. 72. nº 4, pp. 318-330.
- Rando J.C. (1995) Presencia de restos de pigargo (*Haliaeetus* sp.) (Aves: Accipitridae) en yacimientos Paleontológicos de Fuerteventura. *Vieraea* 24:65-69.
- Rando J.C., Alcover J.A., Navarro J.F., García-Talavera F., Hutterer R. and Michaux J. (2008) Chronology and causes of the extinction of the Lava Mouse, *Malpaisomys insularis* (Rodentia: Muridae) from the Canary Islands. *Quaternary Research* 70:141-148.