

Pump-underway ship intake: an opportunity for Marine Strategy Framework Directive (MSFD) monitoring needs. First observations of microplastics on oceanic and coastal waters off the Canary Islands (Subtropical NE Atlantic)

Tania Montoto-Martínez^{1*}, José Joaquín Hernández-Brito^{1,2,3} & María Dolores Gelado-Caballero¹

* ✉ tania.montoto@ulpgc.es

⁽¹⁾ Research group on Environmental Technologies, Management and Biogeochemistry. University of Las Palmas de Gran Canaria (ULPGC); ⁽²⁾ Oceanic Platform of the Canary Islands (PLOCAN); ⁽³⁾ Atlantic International Research Centre (AIR)

Keywords: microplastics, surface waters, continuous monitoring, pump-underway, MSFD, Canary Islands, Atlantic Ocean.

Abstract

Broad scale sampling methods for microplastic monitoring in open ocean waters are a current challenge in oceanography. Large amount of samples is required to understand distribution, abundance and fate of these particles in the environment. The underway water system of research vessels (RVs) has been approached for microplastic sampling in some studies up to date ^[1-5]. We present a sampling methodology founded on the design of a microplastic sampling device connected to the pump-underway ship intake system as an opportunity for oceanic monitoring needs concerning microplastics (**Fig. 1**).

This methodology provides four main advantages: (1) the device is fully made with standard materials, cost-effective and affordable, and it can be self-mounted by the researchers without additional skilled personnel or equipment on-board; (2) it can be employed to report data taking advantage of oceanographic campaigns without interfering their regular vessel activities; (3) it is highly versatile and reduces the time needed to recover each sample; (4) it is suitable to sample microplastic particles down to 50 µm.

As preliminary field application results and first reported data from the Canary Islands oceanic and coastal waters, we investigated microplastics (0,05 – 1 mm) sampled from subsurface waters (-4 m) using the pump-underway system of a RV in the Subtropical NE Atlantic. Sampling was performed in three consecutive oceanographic campaigns over a year, repeating the same procedure, retrieving water while on navigation and on coastal and oceanic stations. Microplastic particles (> 50 µm) were found in the total stations and transects sampled (**Fig. 2**). Fibres (64.42%) were predominant over fragments (35.58%), being the concentration values over the data reported in other areas in the Atlantic.

This system was proved to be an efficient sampling method to report data on microplastic abundance and distribution, addressing the threat that microplastics pose to the marine environment and ecosystems.

Figures




Fig. 1 Sampling set up diagram, showing (a) the pump-underway system of the RV; and (b) the filtering device diagram.




Fig. 2 Pictures showing the typical microplastic particles found: (a) a specially long red fibre, and (b) a red microplastic fragment, showing characteristics for microplastic identification such as irregular edges.

References

- [1] Lusher, A. L., Burke, A., O'Connor, I. & Officer, R.: Microplastic pollution in the Northeast Atlantic Ocean: Validated and opportunistic sampling. *Marine Pollution Bulletin* 88, 325–333 (2014).
- [2] Desforges, J.-P. W., Galbraith, M., Dangerfield, N. & Ross, P. S.: Widespread distribution of microplastics in subsurface seawater in the NE Pacific Ocean. *Marine Pollution Bulletin* 79, 94–99 (2014).
- [3] Enders, K., Lenz, R., Stedmon, C. A. & Nielsen, T. G.: Abundance, size and polymer composition of marine microplastics $\geq 10\mu\text{m}$ in the Atlantic Ocean and their modelled vertical distribution. *Marine Pollution Bulletin* 100, 70–81 (2015).
- [4] Setälä, O., Magnusson, K., Lehtiniemi, M. & Norén, F.: Distribution and abundance of surface water microlitter in the Baltic Sea: A comparison of two sampling methods. *Marine Pollution Bulletin* 110, 177–183 (2016).
- [5] Kanhai, L. D. K., Officer, R., Lyashevskaya, O., Thompson, R. C. & O'Connor, I.: Microplastic abundance, distribution and composition along a latitudinal gradient in the Atlantic Ocean. *Marine Pollution Bulletin* 115, 307–314 (2017).