

de Jong Menno Jeroen (Orcid ID: 0000-0003-2131-9048)  
Tardy Céline (Orcid ID: 0000-0001-5671-5804)  
Hoelzel Rus (Orcid ID: 0000-0002-7265-4180)

## Genomics reveals the role of admixture in the evolution of structure among sperm whale populations within the Mediterranean Sea

Biagio Violi<sup>1,2,3</sup>, Menno J de Jong<sup>2</sup>, Alexandros Frantzis<sup>4</sup>, Paraskevi Alexiadou<sup>4</sup>, Céline Tardy<sup>5</sup>, Denis Ody<sup>5</sup>, Renaud de Stephanis<sup>6</sup>, Joan Giménez<sup>7</sup>, Giuseppe Lucifora<sup>8</sup>, Mónica A. e Silva<sup>9</sup>, Cláudia Oliveira<sup>9</sup>, Filipe Alves<sup>10</sup>, Ana Dinis<sup>10</sup>, Marisa Tejedor<sup>11</sup>, Antonio Fernández<sup>12</sup>, Marina Arregui<sup>12</sup>, Manuel Arbelo<sup>12</sup>, Alfredo Lopez<sup>13,14</sup>, Pablo Coveló<sup>14</sup>, A. Rus Hoelzel<sup>2</sup>.

### Correspondence:

A. Rus Hoelzel and Biagio Violi, Department of Biosciences, Durham University, South Road, Durham, DH1 3LE, UK

Email: [a.r.hoelzel@dur.ac.uk](mailto:a.r.hoelzel@dur.ac.uk) and [violi.biagio@gmail.com](mailto:violi.biagio@gmail.com)

### Affiliations

<sup>1</sup>Department of Earth, Environment and Life Sciences, Genova University, Italy

<sup>2</sup>Department of Biosciences, Durham University, UK

<sup>3</sup>MENKAB: il respiro del mare, Savona, Italy

<sup>4</sup>Pelagos Cetacean Research Institute, Vouliagmeni, Greece

<sup>5</sup>WWF France, Marseille, France

<sup>6</sup>CIRCE, Conservation Information and Research on Cetaceans, Cadiz, Spain

<sup>7</sup>ICM-CSIC, Institut de Ciències del Mar Barcelona, Spain

<sup>8</sup>Istituto Zooprofilattico Sperimentale del Mezzogiorno, Italy

<sup>9</sup>Institute of Marine Sciences – Okeanos and Institute of Marine Research - IMAR, University of the Azores, Rua Professor Doutor Frederico Machado 4, 9901-862 Horta, Portugal

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](https://doi.org/10.1111/mec.16898). Please cite this article as doi: [10.1111/mec.16898](https://doi.org/10.1111/mec.16898)

This article is protected by copyright. All rights reserved.

<sup>10</sup>MARE - Marine and Environmental Sciences Centre / ARNET - Aquatic Research Network, ARDITI, Madeira, Portugal

<sup>11</sup>Oriental Canary Islands Strandings Cetaceans Network, Lanzarote, Spain

<sup>12</sup>Veterinary Histology and Pathology, Atlantic Center for Cetacean Research, University Institute of Animal Health and Food Safety (IUSA), Veterinary School, University of Las Palmas de Gran Canaria, Canary Islands, Spain

<sup>13</sup>Departamento de Biología & CESAM, Universidade de Aveiro, Portugal

<sup>14</sup>CEMMA, Coordinadora para o Estudo dos Mamíferos Mariños, Galicia, Spain.

## KEYWORDS

Genomics, Admixture, Evolution, Population structure, Demography, Cetaceans

## ORCID

*Biagio Violi* <https://orcid.org/0000-0001-9170-1395>

*Menno J. de Jong* <https://orcid.org/0000-0003-2131-9048>

*A. Rus Hoelzel* <https://orcid.org/0000-0002-7265-4180>

## ABSTRACT

In oceanic ecosystems, the nature of barriers to gene flow, and the processes by which populations may become isolated are different from the terrestrial environment, and less well understood. In this study we investigate a highly mobile species (the sperm whale, *Physeter macrocephalus*) that is genetically differentiated between an open North Atlantic population and the populations in the Mediterranean Sea. We apply high resolution single nucleotide polymorphism (SNP) analysis to study the nature of barriers to gene flow in this system, assessing the putative boundary into the Mediterranean (Strait of Gibraltar and Alboran Sea region), and including novel analyses on structuring among sperm whale populations within the Mediterranean basin. Our data support a recent founding of the Mediterranean, around the time of the last glacial maximum, and show concerted historical demographic profiles in both the Atlantic and in the Mediterranean. In each region there is evidence for a population decline around the time of the founder event. The largest decline was seen within the Mediterranean Sea where effective population size is substantially lower (especially in the eastern basin). While differentiation is strongest at the Atlantic/Mediterranean boundary, there is also weaker but significant differentiation between the eastern and western basins of the Mediterranean Sea. We propose, however, that the mechanisms are different. While post-founding gene flow was reduced between the Mediterranean and Atlantic populations, within the Mediterranean an important factor differentiating the basins is likely a greater degree of admixture between the western basin and the North Atlantic and some level of isolation between the western and eastern Mediterranean basins. Subdivision within the Mediterranean Sea exacerbates conservation concerns and will require consideration of what distinct impacts may affect populations in the two basins.

## INTRODUCTION

In oceanic ecosystems, geographic barriers are often less relevant than oceanographic and other environmental factors in shaping the genetic structure of populations (e.g., Hoelzel, 1994; Palumbi, 2004; Van Cise et al., 2019; Westbury et al., 2022). Varying selection pressures, patterns of connectivity and genetic drift shape the diversity of species, potentially leading to greater geographic diversity in species with a wide latitudinal distribution range (Ralph & Coop, 2010). In general, cetaceans are highly mobile with high dispersal potential (see Stevick et al., 2011; Mrusczok et al., 2021; Violi et al., 2021; Peres dos Santos et al., 2022). Even so, they often show population genetic structure over relatively small geographic ranges (e.g., Hoelzel, 2009). In some cases, migratory whales from different stocks may mix on feeding grounds (Palumbi & Baker, 1994; Larsen et al., 1996). In conservation biology, the identification and protection of genetically distinct local populations, including those in mixed assemblages, is essential for conserving evolutionary potential and reducing extinction risks (e.g., Avise, 2009). Given the potential for population structure to be cryptic in the marine environment, both empirical data defining populations and an understanding of the evolutionary mechanisms generating structure are required to facilitate effective conservation and management (e.g., Luck et al., 2003; Holderegger et al., 2020; Hohenlohe et al., 2021).

In this study, we investigate fine-scale population structure in the sperm whale (*Physeter macrocephalus*), a species with a global range across ocean basins (see Whitehead, 2018), considered to play a fundamental role in the balance of the oceanic ecosystems (Würtz & Simard, 2007). Worldwide and for up to three centuries, this species was the focus of two intensive hunting periods (Whitehead, 2002). From a pre-whaling abundance of over 1 million, today there are an estimated 844,761 (Whitehead & Shin, 2022) and they are now globally classified as Vulnerable on the IUCN red list (Taylor et al., 2019).

Range-wide population genetic studies of sperm whales have identified several key features. Genetic variation is low especially at mtDNA (Alexander et al., 2013, 2016; Morin et al., 2018), and low enough to suggest a historical population bottleneck (see Lyrholm & Gyllenstein, 1998; Morin et al., 2018) or cultural hitchhiking (genetic variation co-segregating according to cultural structure; see Whitehead, 2017). In both the Atlantic and Indian Oceans, mtDNA structure was found despite the absence of clear geographic boundaries (Engelhaupt et al., 2009; Alexander et al., 2016). Within the Pacific Ocean the lack of genetic structure at mtDNA was proposed to be consistent with the wider dispersal scale of females compared to Atlantic populations (Rendell et al., 2012; Mesnick et al., 2011; Whitehead et al., 2012). Discrepancy between mtDNA and nuclear-DNA based population structure was suggested to imply male-mediated gene flow (Engelhaupt et al., 2009; Mesnick et al., 2011).

Here, we focus on the populations of sperm whales within the Mediterranean Sea and in nearby Atlantic waters to investigate the mechanisms for maintaining fine-scale population structure in this region. Engelhaupt et al. (2009) reported differentiation between the Mediterranean Sea and North Atlantic at microsatellite DNA and mtDNA loci. Based on a mitochondrial genome phylogeny, Morin et al. (2018) found evidence for differentiation, but proposed that the Mediterranean lineage was founded from the Atlantic population only ~20k years ago, at the end of the last glacial maximum. Consistent with genetic differentiation, sperm whales show differential usage of vocal ‘codas’ (see Frantzis & Alexiadou, 2008) within the Mediterranean compared to elsewhere (Pavan et al., 2000; Drouot et al., 2004b; Teloni, 2005).

Within the northeast Atlantic, sperm whales are regularly present (see Fais et al. 2014; Silva et al. 2014; Alves et al., 2018; Boys et al., 2019; Ferreira et al., 2022). Within the Mediterranean, sperm whales are widely distributed in both the western and eastern basins (see Notarbartolo di Sciara et al., 2008; Pirota et al., 2011, 2020a,b; Aïssi et al., 2012, 2014; Fiori et al., 2014; Mussi et al., 2014; Bellomo et al., 2019; Frantzis et al., 2003, 2014; Rendell & Frantzis, 2016; Tepsich et

al., 2014). In the Strait of Gibraltar, they are regularly seen both east and west of the Strait (de Stephanis et al., 2008), suggesting the potential for gene flow despite evidence for differentiation (e.g. Engelhaupt et al., 2009). Movement by males within the Mediterranean has been broadly documented (e.g., Carpinelli et al., 2014; Rendell et al., 2014; Mussi et al., 2014; Alessi et al., 2014) including movement from the western to the eastern basin (Frantzis et al., 2011), though such direct observations do not necessarily imply local mating success and effective gene flow.

To date, no studies have assessed sperm whale population genetics within the Mediterranean Sea, though this information is essential in support of effective conservation strategies (see Notarbartolo di Sciara, 2014). Conservation planning could furthermore benefit from information on current and past demographic trajectory. Plastic debris ingestion, collisions with ships and entanglements put a strain on the Mediterranean population (Notarbartolo di Sciara, 2014), which numbers fewer than 2,500 mature individuals and is classified as Endangered in the IUCN Red List (Pirotta et al., 2021). Although this assessment is justified by the observed mortality levels at present, no robust data regarding population trends exist (Rendell & Frantzis, 2016).

Here we assess the pattern and mechanisms of demographic trends and population structure of a highly mobile species showing population structure within an oceanic basin. We use high resolution nuclear genetic analyses (*double digest restriction site-associated DNA sequencing: ddRADseq*) to assess the pattern and level of gene flow within the Mediterranean Sea and between the Atlantic Ocean and the Mediterranean Sea. We also consider ongoing and historical population dynamics since the putative founding of the population and implications for the effective conservation management of this endangered population. These data provide transferable inference supporting the conservation of other mobile marine species in similar, recently founded habitats.

## MATERIALS AND METHODS

Sperm whale tissue samples were obtained during various research projects between 1999 and 2018. Skin biopsy samples were collected in the Azores, Madeira, Strait of Gibraltar, and the Ligurian-Provençal area, while skin and muscle samples were obtained from stranded individuals in the Canary Islands, Galicia, Tyrrhenian Sea, Adriatic Sea and the Greek Seas. Altogether the Mediterranean samples were sourced from six different areas in four different countries (Greece, Italy, France and Spain) and the Atlantic samples were from four different areas (Galicia, Azores, Canary Islands and Madeira) (Figure 1 and Table S1).

### DNA Extraction and Library prep

DNA was extracted both by kit (OMEGA BIOTEK and MN MACHEREY-NAGEL) following the manufacturers protocol, and by the phenol chloroform method (after Hoelzel, 1998). Genomic DNA concentration was quantified using the Qubit High Sensitivity kit (Thermo Fisher Scientific). We applied the ddRADseq methodology (Peterson et al., 2012) to sperm whales for the first time in this study. Two Sequencing libraries, each of 80 samples, were constructed following the protocol in Peterson et al. (2012). DNA (500ng) was digested using enzymes *MspI* and *HindIII* (New England Biolabs). Adapters were ligated using T4 ligase in a reaction solution with 1 X Buffer, 400 units of T4 Ligase and 1.5  $\mu$ M of adapter that contained a unique in-line barcode for each individual. Uniquely barcoded samples were pooled, and cleaned using calibrated streptavidin coated SpeedBeads (Sera-Mag). Fragments between 360 and 560 bp were size-selected using a Pippin Prep (Sage Science). Sixteen uniquely indexed libraries were prepared, each containing ten individually barcoded samples, with randomized allocation of individuals across libraries. Fragment size distributions of resulting libraries were evaluated using a 2200 TapeStation (Agilent Technologies), and library DNA concentrations were estimated using qPCR. Two final libraries

were then sequenced on an Illumina HiSeq 2500 using 125 bp paired-end sequencing at the DBS Genomics facility at Durham University (<https://www.dur.ac.uk/biosciences/services/dna/dnasequencing/>).

## SNP calling

Reads were trimmed to 110 bp, demultiplexed and filtered using `process_radtags` in STACKS v. 2.2 (Rochette et al., 2019). Unpaired reads were discarded. Paired reads were aligned using the very-sensitive mode of Bowtie v. 2.2.5 (Reinert et al., 2015) against a sperm whale reference genome downloaded from the NCBI website (GCA\_002837175.2). Samtools v. 1.9 (Li et al., 2009) was used to filter out reads which aligned discordantly, as well as reads with a mapping quality below 20. SNPs were called using the STACKS refmap pipeline with default settings. Sites were not filtered based on depth (an option which has purposely been removed from STACKS version 2.0 onwards), but instead based on a likelihood ratio test which weighs the evidence for heterozygous versus homozygous calls (which depends partially on coverage), using the default p-value threshold of 0.05. The Populations command of STACKS was run with  $r=0.7$  and  $p=3$ . We accepted multiple SNPs per read and thinned the dataset downstream when appropriate. PGD-Spider (Lischer & Excoffier, 2012) or vcftools (Danecek et al., 2011) was used to convert the SNP data into PED and MAP format. Binary files (BED, RAW and BIM) were generated from PED and MAP files using Plink (Purcell et al., 2007). The vcftools flags `--depth` and `--site-depth` were used to calculate read depth per individual and per SNP. SNP data management and analyses were performed in R v. 3.6.2 (R Core Team, 2019) using wrapper functions of the R package SambaR (De Jong et al., 2021). The data was imported into R and stored in a `genlight` object using the function `'read.PLINK'` of the R package Adegenet v. 2.1.1 (Jombart & Ahmed, 2011).



## SNP and sample filtering

Two datasets were obtained using two different threshold filters. Dataset A excluded samples with more than 50% missing data and all loci with greater than 5% missing data. Dataset B excluded all samples with more than 25% missing data and all loci with greater than 5% missing data (see Table S2). Both datasets were thinned selecting one SNP per 500 bp region, in order to limit SNPs to one per paired-end read, and used for further structure and diversity analyses.

## Population structure

The two datasets were used in most of the analyses to test for divergence between Atlantic and Mediterranean populations and within the Mediterranean putative western and eastern populations. Replicates were detected using KING-robust score calculations (Manichaikul et al., 2010) and were excluded from the analyses. Kinship analyses were performed only on dataset B due to the higher number of SNPs, using the  $\pi_{\text{hat}}$  score of the software Plink (Purcell et al., 2007). The background variation was based on the Atlantic and Med samples separately.

Population structure was assessed in R, using SambaR wrapper functions. Principal coordinate analysis (PCoA) was calculated using the Ape v. 5.3 (Paradis & Schliep, 2019) for both datasets. PCoA is more suitable for the analyses of categorical data (like genotype data: 0, 1 and 2) than PCA. Structure analyses were performed using two methods: the Landscape and Ecological Studies (LEA 3.10.0) package (Frichot & Francois, 2015) setting the putative number of population (K) to 2-4, and Admixture 1.3 (Alexander et al., 2009), giving prior assignment to Atlantic and Mediterranean populations. We did 20 repeat runs using PONG (<https://github.com/ramachandran-lab/pong>; see Behr et al 2016), using a greedy approach to identify modes and default threshold (0.95) for determining modes. F-statistics were assessed in Arlequin v. 3.5 (Excoffier & Lischer, 2010) with 10,000 permutations and the level of missing data set to 0.05.  $F_{\text{ST}}$  was calculated for

both datasets at three levels: 1) all sampled areas; 2) Atlantic vs. Mediterranean populations; and 3) Atlantic vs. western vs eastern putative populations.

#### Migration rates and population modelling

Contemporary migration rates were calculated using BayesAss3-SNPs (Mussmann et al., 2019) using both datasets comparing Atlantic vs Mediterranean populations and Atlantic vs. western vs. eastern Mediterranean populations. BayesAss3 uses allele frequency data and MCMC simulations to find individuals that are migrants or have migrant ancestry “over the last several generations”. The analysis was run with 1,000,000 iterations and a burn-in of 100,000 iterations, seed of 10 and delta value of 0.1. Output matrices were converted into gene flow plots with the use of the R package Circlize v. 0.4.6 (Gu et al., 2014) through SambaR.

We considered 4 scenarios in DIYABC (see Collin et al., 2021; <https://github.com/diyabc>) to test inference about 1 or 2 populations in the Mediterranean. These had the Atlantic as an ancestral population (at t2) and a split within the Mediterranean later at t1, the Western Med as ancestral, the Eastern Med as ancestral, and only two populations, Atlantic and Mediterranean. We excluded admixed individuals, and the number of simulations in the training set was to 120 while the number of trees was 500. The priors were default flat priors.

#### Admixture and population dynamics

Patterns of admixture were interrogated using the Unix software TreeMix v. 1.13 (Pickrell & Pritchard, 2012) using the Atlantic population as an outgroup and testing for 1 migration event, while the parameter f3 was calculated using the R program Admixtools v. 2 using default settings (Patterson et al., 2012). Both analyses were performed on sample set B comparing the putative Atlantic, western Mediterranean and eastern Mediterranean populations. TreeMix uses allele

frequencies to construct a maximum likelihood tree and builds a model which allows for both population splits and gene flow to infer the pattern of admixture among lineages. The analysis was run with Bootstrap replicate over blocks of 500 contiguous SNPs (as suggested in the manual). The  $f_3$  statistic assesses the covariance of allele frequencies between a population pair with a shared outgroup. Measuring the shared drift between the population pairs in the context of drift that is specific to single populations allows inference about their relationship and the chance of admixture.

The historical demography of the Atlantic and the Mediterranean populations was inferred using the Stairway plot analysis (Liu & Fu, 2020). Due to potential issues with admixture affecting demographic inference (e.g., Heller et al., 2013), we analysed pooled population samples that showed no evidence of differentiation. One sample from Galicia from the Atlantic sample was omitted due to missing data, and those individuals from Liguria with a clear signal for admixture with the Atlantic were also excluded. The folded site frequency spectra (SFS), was calculated using the software package ANGSD v. 0.938 (Korneliussen et al., 2014). The proportion of sites used for training was 0.67, and the number of input files created for estimations was 200. As in Morin et al. (2018), the mutation rate was set at  $2.9 \times 10^{-8}$  substitutions per nucleotide per generation based on Dornburg et al., (2012) and the generation time at 31.9 years based on Taylor et al. (2007). A comparative assessment that can estimate relatively recent  $N_e$  with greater accuracy was applied (using the same sample sets as for the stairway plots), drawing inference from linkage disequilibrium (LD). This was assessed using SNeP software v1.11 (Barbato et al., 2015, after Corbin et al., 2012).  $N_e$  estimation was calculated with a minimum and maximum distance between SNPs of 50,000 and 4 Mb, respectively. Data was arranged in 30 bins of 50 kb distance each. Maximum number of SNPs per chromosome was set to 100,000. The formula used to estimate  $N_e$  from LD was as proposed by Sved & Feldman (1973), as recommended from a comparative analysis by García-Cortés et al. (2018).

## RESULTS

### Reads and SNPs.

A total of 160 sperm whale individuals were sequenced. Both sequencing lanes combined produced 628.9 million reads. Up to 16.2 million reads had to be discarded due to either low quality, an ambiguous radtag, or a missing read mate, resulting in an average of 1.9 million read pairs per sample (stdev: 1.3 million, min: 0.003 million, max: 6.5 million). Individuals with less than 10, 1, 0.5 and 0.25 percent missing data had a minimum cover of respectively 3, 8.6, 8.6, and 23.2 reads. The mean sequence depth per individual ranged from 3.2 to 184.4 reads per SNP, with a mean 12.4 reads. From a total of 46,717 SNPs, after filtering and thinning, 132 samples and 4,422 SNPs were retained for dataset *A* and 117 samples, and 12,616 SNPs were retained for dataset *B*. Ten samples found as replicates were excluded (Table S1 and S2).

### Population structure and gene flow

Comparisons assessing genetic differentiation between the Atlantic and Mediterranean populations based on  $F_{ST}$  were significant for both datasets, as was the  $F_{ST}$  value comparing the western and eastern basins (Table 1). The  $F_{ST}$  value within the Mediterranean was smaller than for comparisons with the Atlantic, but significantly greater than zero. A subdivision between the basins is sensible based on physical and oceanographic differences, and based on evidence of differentiation across this boundary for other highly mobile species (such as the bottlenose dolphin, *Tursiops truncatus*; see Natoli et al., 2005 and discussion therein). This established a pattern that we then further investigated using ordination (PCoA) and assignment (LEA and Admixture) methods.

Principal coordinate analyses (PCoA) revealed that Atlantic samples form a tight cluster (blue circle) separated from Mediterranean samples (green circle; both datasets are shown in Figure 1B, C). The Mediterranean samples are distributed over a broader range of Euclidian space in the

plot, and there is only a weak pattern of differentiation among samples from the western and eastern basins (Figure S1). The pattern is somewhat clearer when only samples from within the Mediterranean are included, but there is still overlap (Figure 1D, E). Kinship estimations ( $\pi$ -hat scores) are consistent with lower overall pairwise kinship among samples collected in the Atlantic than among samples collected in the Mediterranean Sea (Figure S2). While the values are consistently higher in the Mediterranean, they also extend over a broader range (Figure S2). Support values are low per axis, but this is typical when based on many loci.

LEA analyses with values of  $K=2-4$  revealed a clear differentiation between the Mediterranean and Atlantic samples and for  $K=3-4$ , inconsistent differentiation between the western and eastern Mediterranean basins (Figure 2). A cross-entropy analysis shows that both  $K=2$  and  $K=3$  are equally well supported (Figure S3). As expected, the pattern was somewhat clearer when only Mediterranean samples were included (Figure 2C). This is confirmed using Pong to repeat the Admixture Mediterranean only analyses 20 times (Figure S4). For  $K=2-4$  for the Med only samples (Figure S5) the lowest validation error is again  $K=2$  ( $K2\ cv = 0.22375$ ;  $K3\ cv = 0.24256$ ;  $K4\ cv = 0.25900$ ). For Bayesian modelling in DIYABC, three populations (Atlantic as ancestral followed by an east vs west split within the Mediterranean) was as well supported as a model that considered only two population (Atlantic and Mediterranean; Figure S6).

Mixed Atlantic/Mediterranean ancestry was found within Mediterranean areas consistently at all  $K$  values. The proportions of the admixture coefficient for  $K=2$  was estimated in Admixture, based on separate Atlantic and Mediterranean putative populations ( $K=2$ ), as  $K=2$  gave the lowest cross validation error when all samples were included ( $K=2$ :  $cv\ 0.2234$ ;  $K=3$ :  $cv\ 0.2318$ ;  $K=4$ :  $cv\ 0.2486$ ). This analysis revealed a subset of Mediterranean samples with mixed Atlantic/Mediterranean ancestry, especially from the western basin (Figure 2B). That was the case for 29 samples ( $29/85=34\%$ ) overall, including 3 ( $3/85=3.5\%$ ) samples from the Strait of Gibraltar, 24 ( $24/85=28.2\%$ ) samples from Ligurian-Provençal area, 1 ( $1/85=1.1\%$ ) sample from Tyrrhenian

Sea and 1 (1/85=1.1%) sample from the Adriatic Sea. These samples showed assignment probabilities to the Atlantic population ranging from 14 to 74 % (Table S3). Averaged over all samples within each basin, there was 12.6% admixture from the Atlantic in the western basin, and 0.7% in the eastern basin. Individuals showing some level of admixture in the Admixture analysis were marked as red dots in the PCoA shown in Figure S1B and all group within the Mediterranean cluster. An  $f_3$  analysis shows marginal values, but a stronger indication of admixture for the western region with the Atlantic than for other possible combinations (Figure 3C and Table 4). The analysis in Treemix showed the same pattern, with admixture suggested from the Atlantic into the western region (Figure 3D), though the drift parameter is low for the tree overall. Neither of these methods are specifically identifying only recent migrants (as in BayesAss3). Also, while BayesAss3 can consider bidirectional gene flow, in TreeMix we could only test one migration edge due to the simplicity of the tree.

Investigations of contemporary migration rate were conducted with BayesAss3-SNPs. A first analysis was run for both datasets between the Atlantic and Mediterranean populations and revealed an estimated proportion of migrants of ~1% through the Strait of Gibraltar in both directions (Figure 3A and Table 2), consistent with either low gene flow or the relatively recent cessation of gene flow. A second analysis, between the Atlantic, western and eastern Mediterranean putative populations indicated recent or ongoing directional movement between the two basins from west to east (about 18-26%), and low (~1%) gene flow between each basin and the Atlantic (Figure 3B, Table 3).

## Demography

All three stairway plot profiles (Figure 4) showed an apparent increase in population size during the early part of the Pleistocene, though recent time periods (within ~10 Ka) are less precise by this method and therefore should be interpreted with caution. All three regions showed a similar

effective population size ( $N_e$ ) (in the 10k to 20k range) and profile prior to the last glacial maximum (LGM ~20ka). After that, all populations showed a decline, but this was most severe within the Mediterranean (down to ~1k while in the Atlantic the decline was only to ~8k). Our comparative assessment using inference from LD (in the program SNeP) showed a similar estimate for  $N_e$  at ~7Ka, but then a steeper decline to the present (Figure S7). The contemporary estimate of  $N_e$  was low by this method (~100; within the confidence limits of estimates from the stairway plots).  $N_e$  for the Atlantic was largest followed by the western and then eastern Mediterranean, but all had a similarly low level for the most recent estimates.

## Discussion

### Atlantic vs. Mediterranean divergence

Our data support the differentiation between the North Atlantic and Mediterranean Sea described in previous genetic and acoustic studies (e.g., Engelhaupt et al., 2009; Drouot et al., 2004a; Rendell & Frantzis, 2016; Mesnick et al., 2011; Morin et al., 2018). Pairwise comparison between these two areas in our study based on 12,616 SNP markers reveals a low but significant value ( $F_{ST}=0.03454$ ,  $P\text{-value}=0.000$ ), consistent with that reported in Engelhaupt et al., (2009; where  $F_{ST}=0.034$ ,  $P\text{-value}=0.022$  based on 16 microsatellite DNA markers). Significant genetic division between the Atlantic and Mediterranean areas has also been documented for other cetaceans such as fin whale (*Balaenoptera physalus*; Berubé et al., 1998; Palsboll et al., 2004), bottlenose dolphin (*Tursiops truncatus*; Natoli et al., 2005), Cuvier's beaked whale (*Ziphius cavirostris*; Onoufriou et al., 2022) and for other top predators such as swordfish (*Xiphias gladius*; Smith et al., 2015) and the blue fin tuna (*Thunnus thynnus*; Puncher et al., 2018). Our findings from LEA and Admixture were consistent with Atlantic ancestry of the Mediterranean populations. Estimates of contemporary migration patterns indicate low gene flow through the Strait of Gibraltar,

approximately 1% migration per generation in both directions. Sperm whales have been routinely observed in the Strait of Gibraltar, both in summer (de Stephanis et al., 2008) and winter (Gauffier et al., 2012, 2018). Some movements from the Strait of Gibraltar to the Balearic Islands and the Ligurian-Provençal area and vice versa have also been documented through photoidentification (Carpinelli et al., 2014). Historical reports from Bolognari (1949, 1950, 1951) suggested seasonal movements through the Strait of Gibraltar – entering in the winter and going out in spring. Movements from the Atlantic into the Mediterranean Sea have also been recorded for other cetacean species (e.g., fin whales, Castellote et al., 2010; Giménez et al., 2013; Gauffier et al., 2018, 2020; humpback whales *Megaptera novaeangliae*, Violi et al., 2021; killer whales *Orcinus orca*, Mrusczok et al., 2021).

A complex network of submarine canyons in the western Mediterranean act as corridors between continental shelf areas and deep-sea regions, enhancing oceanographic processes that enrich the deep-sea food web (De Leo et al., 2010) and create a favourable environment for the cephalopod species (e.g., Quetglas et al., 2000; O'Dor & Coelho, 1993) that sperm whales prey on. Therefore, this region can support the energetic demands of sperm whales and other deep divers (Torreblanca et al., 2022). Distinct resource availability and resource specialisations have been proposed as drivers of differentiation in various cetacean species (see Hoelzel 2009; Louis et al., 2014). However, the umbrella squid *Histioteuthis bonnellii* is evidently the most common prey of sperm whales both in the Mediterranean Sea (Garibaldi & Podestà, 2014; Foskolos et al., 2020; Tonay et al., 2021) and in the Atlantic Ocean (Clarke et al., 1993), and so resource specialisation may not be an important driver in this case. We note that in this study we have not tested for the various other environmental drivers that may be relevant, such as surface temperature and salinity.



## Structure within the Mediterranean

Within the Mediterranean, we report weak differentiation between the basins either side of Italy and the Strait of Sicily, as observed in other species (e.g., bottlenose dolphin in Natoli et al., 2005; Gaspari et al., 2015; striped dolphin in Gaspari et al., 2007; Cuvier's beaked whale in Onoufriou et al., 2022). The relatively shallow sill at the Strait of Messina and the Sicilian Channel, and the differential current patterns in the two basins likely reinforce this boundary. Our results from PcoA and LEA were largely consistent with  $F_{ST}$ , which showed significant differentiation between the western and eastern basins, though weaker than the differentiation between the North Atlantic and Mediterranean Sea. Assignment analysis (using LEA) restricted to the Mediterranean samples showed a weak pattern when  $K=3$  or 4 (Figure 2), but this type of assignment method has low power to resolve a pattern when  $F_{ST}$  is lower than  $\sim 0.02$  (Latch et al., 2006). Ordination (PCoA) showed some separation between western and eastern basin samples, but some overlap as well. We found evidence of admixture between the North Atlantic and Mediterranean Sea, based on analyses using the program Admixture (Figure 2),  $f_3$  statistics and TreeMix (Figure 3). We propose that the signal for differentiation between the two basins is driven largely by the more frequent historical admixture between the Atlantic and the western basin than with the eastern basin. Signals for ongoing migration based on BayesAss also show relatively low, bi-directional gene flow between the Atlantic and the Mediterranean, but relatively high, directional gene flow from the western into the eastern basin (see Figure 3). One possible interpretation would be that the signal for differential admixture from the Atlantic into the western basin predates a more recent increase in gene flow from the western into the eastern basin (since BayesAss should register only recent movement).

## Diversity & Demography

Overall, the Mediterranean samples show higher pairwise kinship than the samples from the North Atlantic (Figure S2), reflecting reduced diversity overall (assuming random sampling). This is consistent with the results from demographic profiling, where the estimated contemporary  $N_e$  is approximately six times lower in the Mediterranean than in the Atlantic based on the stairway plots (Figure 4), though the difference was smaller based on SNeP (Figure S7). After a period of relative stasis through the latter part of the Pleistocene, all putative populations show a decline starting around the time of the LGM (Figure 4), and the overall profile is similar for all. However, the Mediterranean trajectories diverge from the Atlantic trajectory at that time. This would fit with a timeframe of within the last 20 ka for the founding of the Mediterranean by sperm whales, as suggested earlier (Morin et al., 2018). The decline in the Mediterranean might be explained by a relatively small founder population, a low rate of migration between the Atlantic and Mediterranean after founding, and a consequent reduction in  $N_e$  within the Mediterranean. At the same time, the parallel pattern of demography is interesting, suggesting that both populations were impacted by a similar process leading to population decline. Based on mtDNA mismatch distributions, Alexander et al. (2016) suggested a post-bottleneck expansion signal world-wide as recently as 10Ka. The stairway plots and SNeP gave similar  $N_e$  estimates between ~7Ka and 1Ka (especially within the Mediterranean), but the stairway plot estimates then level off, and are known to be less accurate for very recent timeframes (reflected in the confidence limit estimates; Figure 4). SNeP estimates continue to decline, suggesting low  $N_e$  in modern populations both in the Mediterranean and in the Atlantic.

Episodes of ocean cooling/warming are proposed to have influenced the distribution and abundance of various cetacean species and such events could have led to restricted home ranges (Gaskin, 1982). Morin et al. (2018) proposed this scenario for sperm whales, whose suitable habitat for females was reduced up to 50% since the last glacial maximum. Cold regions at high latitude

were covered by ice, and sperm whale males, which usually feed in these regions, may have instead occupied the same home range as females and their social units, which normally do not range into cold waters (Best, 1979). This change of age/sex classes distribution and overlap could have led to competition for resources and a reduction of food availability.

Some species known to be preyed on by sperm whales, such as the giant squid (*Architeuthis* spp. Winkelmann et al., 2013), the Humboldt squid (*Dosidicus gigas* Ibáñez et al., 2011) and the Patagonian longfin squid (*Doryteuthis gahi* Ibanez & Poulin, 2014) show low mitogenomic diversity and signatures of demographic/range expansions associated with the LGM. Some cetaceans species such as long-finned pilot whale (*Globicephala melas* Kraft et al., 2020), northern bottlenose whale (*Hyperoodon ampullatus* Feyrer et al., 2019), white-sided dolphins (*Lagenorhynchus obliquidens* Banguera-Hinestroza et al., 2010) and killer whales (*Orcinus orca* Moura et al., 2014) also show a similar expansion pattern.

The Mediterranean Sea played a key role as a refuge for several marine species during the last glacial period (see Patarnello et al., 2007). During Pleistocene glaciations, sea level was reduced up to 150 m (Lambeck & Purcell, 2005), changing oceanographic features that created isolated refugia and geographical barriers to gene flow, followed by post-glacial dispersal and expansion (Xue et al., 2014). Previous studies have suggested that Mediterranean biodiversity more generally is the result of endemism from glacial refugia (see Patarnello et al., 2007). During the LGM, changes in the pattern of deep-water availability were more pronounced in the eastern basin (Mikolajewicz et al., 2011; Thunell & Williams 1989). In the western basin, sperm whales could have found suitable deep water habitat to colonize where there was likely to be cephalopod prey.

## Conservation

Our data are consistent with earlier studies showing significant genetic differentiation between the North Atlantic and Mediterranean sperm whale populations, and within the

Mediterranean greater identity by descent and consequently lower  $N_e$ . Demographic profiles show a shared history until around the time of the LGM, after which the decline in  $N_e$  in the Mediterranean was likely greater than in the Atlantic. Our estimates of effective population size are low, consistent with the IUCN-Red list assessment, which classified the Mediterranean sperm whale population as Endangered (see Pirotta et al., 2021). Our data supports the need to manage the Atlantic and Mediterranean populations separately. Estimates of gene flow suggest low contemporary levels between the Atlantic and the Mediterranean, but more historical admixture. Greater historical admixture in the west and some level of isolation between the basins appears to explain the pattern of differentiation between basins, though gene flow may have increased recently (from west to east based on the analysis in BayesAss). Any persistent subdivision within the Mediterranean Sea would exacerbate conservation concerns and require consideration of what distinct impacts may affect populations in the two basins. Here we find a weak, but significant subdivision between the eastern and western basins, probably driven by differential patterns of admixture with the Atlantic population (more in the west). We also find evidence for a lower  $N_e$  in the eastern basin (see Figure S7). We therefore would propose that the eastern and western populations be considered separate management units.

It has been proposed that sperm whales in the Mediterranean are not directly threatened by fisheries since their main prey is *H. bonnellii* (Garibaldi et al., 2014; Foskolos et al., 2019), which is not a target of fishery activity. However, Mazzariol et al., (2011) found several hooks within stomach content analyses of stranded sperm whales suggesting possible feeding activity along fishing lines as described in some oceanic regions (e.g., Jacobsen et al., 2010). At the same time, fishery activity can also impact sperm whales through entanglement in fishing nets (e.g., Pace et al., 2008; Blasi et al., 2021). Ghost nets, ship collision and the ingestion of plastic debris are the three main impacts that are seriously threatening these endangered sperm whales in the Mediterranean Sea (de Stephanis et al., 2013; Alexiadou et al., 2019; Frantzis et al., 2019).

## Conclusions

Marine systems can be a challenge for the identification of patterns of population structure and the units of conservation for effective management (e.g., Alves et al., 2019). Apart from obvious continental barriers, physical barriers are few, and even continental barriers can be overcome (by going around; see Hoelzel et al., 2021). However, oceanic structures, such as current systems (e.g., Knutsen et al., 2007) that move or concentrate larvae, prey or predators can determine distribution patterns that remain cryptic unless the relevant drivers are understood (see Gonçalves da Silva et al., 2019). This reflects a need to combine genetics with ecological data to integrate evidence for both genetic and ecological management units (e.g., Esteban et al., 2016; Giménez et al., 2018). Historical changes in species distributions, such as those driven by the glacial cycles of the Pleistocene, can also shape modern day population structure (e.g., Bracken et al., 2015). In our study, we find that a highly mobile species likely founded an isolated population around the end of the last glacial period, and that this founder population was further structured by some period of admixture between the source population and the region within the Mediterranean most proximate to the source. This may be a useful model for investigating other patterns of cryptic structure in marine systems.

## ACKNOWLEDGMENTS

This work was supported by OceanCare (Switzerland), project *Catodon* of the *Menkab, il respiro del mare* association, and MPA Bergeggi Island. Sampling was supported by Loro Parque Foundation, the Fundación Biodiversidad, FCT (IF/00943/2013/CP1199/CT0001, UIDB/05634/2020 & UIDP/05634/2020), FRCT, the EC through research projects TRACE-PTDC/MAR/74071/2006, MAPCET-M2.1.2/F/012/2011, WATCH IT (Acores-01-0145-FEDER-

000057) [FEDER, COMPETE, QREN, POPH, ESF, Portuguese Ministry for Science and Education, OP Azores 2020, the regional government Xunta de Galicia – Dirección Xeral de Patrimonio Natural, research program “Sperm whales of the Greek Seas” of the Pelagos Cetacean Research Institute partially supported by OceanCare since 2008, Oceanic Observatory of Madeira throughout the project M1420-01-0145-FEDER-000001-OOM and of the Oceano Azul Foundation throughout the Whale Tales Project (ODL/2019/003). Biagio Violi was funded by Department of Earth and Life Sciences of Genoa University. Joan Giménez was supported by the Spanish National Program Juan de la Cierva-Formación (FJC2019-040016-I) and the Severo Ochoa Centre of Excellence accreditation (CEX2019-000928-S) given to the Institute of Marine Science (ICM-CSIC). Ana Dinis and Filipe Alves were funded by ARDITI throughout the project M1420-09-5369-FSE-000002 and by FCT throughout the projects UIDB/04292/2020 and UIDP/04292/2020 granted to MARE and project LA/P/0069/2020 granted to the Associate Laboratory ARNET. Mónica Almeida e Silva was co-funded by OP Azores 2020 (Fund 01-0145-FEDER-000140-MarAZ Researchers) of the EU. Cláudia Oliveira was funded by WATCH IT – Acores-01-0145-FEDER-000057, and service contract with SRMCT/DRAM under project INTERTAGUA (MAC2/1.1a/385). Okeanos is supported by FCT (UIDB/05634/2020 & UIDP/05634/2020) and the Regional Government of the Azores (M1.1.A/REEQ.CIENTÍFICOUI&D/2021/010). Alfredo Lopez was supported by the CESAM by FCT/MCTES (UIDP/50017/ 2020+UIDB/ 50017/2020 + LA/P/0094/2020), through national funds and by national funds (OE), through FCT-I.P., in the scope of the framework contract foreseen in the numbers 4, 5 and 6 of the article 23, of the Decree-Law 57/2016, of August 29, changed by Law 57/2017, of July 19. We thank Prof. Maurizio Würtz, Prof. Alberta Mandich, Daniel Moore, Mario Erandi Bonillas, Vania Fonseca, Sofia da Silva, Thomas Keggins, Luke Hecht, Ali Reza Foroozani, Michelle Gaither, Howard Gray, Karis Baker, Andreanna Welch, Fulvio Garibaldi, Phillip Morin, Alana Alexander, Andre Moura, Xiaoming Liu and Mark Dixon for helpful advice. We thank the Prof. Bruno Cozzi, Prof. Sandro Mazzariol,

Mattia Panin and Cinzia Centelleghes, Rita Ferreira, Annalisa Sambolino, Philippe Verborgh, Juan Manuel Salzar Sierra, Pauline Gauffier, Ruth Esteban, Rui Prieto, Marta Tobeña, Irma Cascão, and Sergi Pérez-Jorge for samples. Samples in Madeira were collected under permit number 508/IFCN IP-RAM/2018. Sampling in the Canary Islands was done under permission from the Ministry of Agriculture, Fisheries, Food and Environment of Spain. Sampling in Galicia was done under permission from Xunta de Galicia – Dirección Xeral de Patrimonio Natural.

## REFERENCES

- Aïssi, M., Fiori, C., & Alessi, J. (2012). Mediterranean submarine canyons as stepping stones for pelagic top predators: the case of sperm whale. *Mediterranean Submarine Canyons: Ecology and Governance*. IUCN, 99-103.
- Aïssi, M., Ouammi, A., Fiori, C., & Alessi, J. (2014). Modelling predicted sperm whale habitat in the central Mediterranean Sea: requirement for protection beyond the Pelagos Sanctuary boundaries. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(S1), 50-58. <https://doi.org/10.1002/aqc.2411>
- Alessi, J., Aïssi, M., & Fiori, C. (2014). Photo-identification of sperm whales in the north-Western Mediterranean Sea: an assessment of natural markings. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(S1), 11-22. <https://doi.org/10.1002/aqc.2427>
- Alexander, D.H., Novembre, J., Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Res* 19 (9):1655–1664. <https://doi.org/10.1101/gr.094052.109di>
- Alexander, A., Steel, D., Slikas, B., Hoekzema, K., Carraher, C., Parks, M., ... & Baker, C. S. (2013). Low diversity in the mitogenome of sperm whales revealed by next-generation sequencing. *Genome biology and evolution*, 5(1), 113-129. <https://doi.org/10.1093/gbe/evs126>
- Alexander, A., Steel, D., Hoekzema, K., Mesnick, S. L., Engelhaupt, D., Kerr, I., ... & Baker, C. S. (2016). What influences the worldwide genetic structure of sperm whales (*Physeter macrocephalus*)? *Molecular ecology*, 25(12), 2754-2772. <https://doi.org/10.1111/mec.13638>
- Alexiadou, P., Foskolos, I., & Frantzis, A. (2019). Ingestion of macroplastics by odontocetes of the Greek Seas, Eastern Mediterranean: Often deadly!. *Marine pollution bulletin*, 146, 67-75. <https://doi.org/10.1016/j.marpolbul.2019.05.055>
- Alves, F., Alessandrini, A., Servidio, A., Mendonça, A. S., Hartman, K. L., Prieto, R., ... & Aguilar de Soto, N. (2019). Complex biogeographical patterns support an ecological connectivity network of a large marine predator in the north-east Atlantic. *Diversity and Distributions*, 25(2), 269-284. <https://doi.org/10.1111/ddi.12848>
- Alves F., Ferreira R., Fernandes M., Halicka Z., Dias L., Dinis A. (2018). Analysis of occurrence patterns and biological factors of cetaceans based on long-term and fine-scale data from platforms of opportunity: Madeira island as a case study. *Mar. Ecol.* 39 (2), e12499. doi: 10.1111/maec.12499



- Avise, J. C. (2009). Phylogeography: retrospect and prospect. *Journal of biogeography*, 36(1), 3-15. <https://doi.org/10.1111/j.1365-2699.2008.02032.x>
- Banguera-Hinestroza, E., Bjørge, A., Reid, R. J., Jepson, P., & Hoelzel, A. R. (2010). The influence of glacial epochs and habitat dependence on the diversity and phylogeography of a coastal dolphin species: *Lagenorhynchus albirostris*. *Conservation Genetics*, 11(5), 1823-1836. <https://doi.org/10.1007/s10592-010-0075-y>
- Barbato, M., Orozco-terWengel, P., Tapio, M., & Bruford, M. W. (2015). SNeP: a tool to estimate trends in recent effective population size trajectories using genome-wide SNP data. *Frontiers in genetics*, 6, 109. <https://doi.org/10.3389/fgene.2015.00109>
- Behr, A. A., Liu, K. Z., Liu-Fang, G., Nakka, P., & Ramachandran, S. (2016). Pong: fast analysis and visualization of latent clusters in population genetic data. *Bioinformatics*, 32(18), 2817-2823. <https://doi.org/10.1093/bioinformatics/btw327>
- Bellomo, S., Santacesaria, F. C., Fanizza, C., Cipriano, G., Renò, V., Carlucci, R., & Maglietta, R. (2019). Photo-identification of *Physeter macrocephalus* in the Gulf of Taranto (Northern Ionian Sea, Central-Eastern Mediterranean Sea). In *IEEE metrology for the Sea 2019 IMEKO TC-19 International workshop on metrology for the Sea* (pp. 33-37).
- Best, P.B. (1979). Social organization in sperm whales, *Physeter macrocephalus*. In *Behavior of marine animals*. pp. 227–289. Springer US.
- Blasi, M. F., Caserta, V., Bruno, C., Salzeri, P., Di Paola, A. I., & Lucchetti, A. (2021). Behaviour and vocalizations of two sperm whales (*Physeter macrocephalus*) entangled in illegal driftnets in the Mediterranean Sea. *PloS one*, 16(4). <https://doi.org/10.1371/journal.pone.0250888>
- Bolognari, A. (1949). A proposito della recente cattura di alcuni esemplari di Capodoglio (*Physeter macrocephalus* L.) nel Mediterraneo. *Bull. Inst. Océan.*, Monaco; n. 99.
- Bolognari, A. (1950). Ancora sulla comparsa del Capodoglio: (*Physeter macrocephalus* L.) nel Mediterraneo. *Italian Journal of Zoology*, 17(1-3), 29-37.
- Bolognari, A., (1951). La migrazione del Capodoglio nel Mediterraneo. *Bolletino di Zoologia*, 18, 253-256. Centro Studi Cetacei, 1986-2000. Cetacei spiaggiati lungo le coste italiane. Rendiconto. Atti della Società Italiana di Scienze Naturali di Milano.
- Boys, R. M., Oliveira, C., Pérez-Jorge, S., Prieto, R., Steiner, L., & Silva, M. A. (2019). Multi-state open robust design applied to opportunistic data reveals dynamics of wide-ranging taxa: the sperm whale case. *Ecosphere*, 10(3), e02610. <https://doi.org/10.1002/ecs2.2610>
- Bracken, F.S.A, Hoelzel A.R., Hulme, J.B. & Lucas, M.C. (2015). Contrasting population genetic structure among freshwater-resident and anadromous lampreys: the role of demographic history, differential dispersal, and anthropogenic barriers to movement. *Mol. Ecol.* 24, 1188–1204. <https://doi.org/10.1111/mec.13112>
- Carpinelli, E., Gauffier, P., Verborgh, P., Airoidi, S., David, L., Di-Méglio, N., ... & de Stephanis, R. (2014). Assessing sperm whale (*Physeter macrocephalus*) movements within the West Mediterranean Sea through photo-identification. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(S1), 23-30. <https://doi.org/10.1002/aqc.2446>
- Castellote, M., Clark, C. W., & Lammers, M. O. (2010). Population identity and migration movements of fin whales (*Balaenoptera physalus*) in the Mediterranean Sea and Strait of Gibraltar. *IWC Scientific Committee, Agadir, Morocco*.
- Clarke, M. R., Martins, H. R., & Pascoe, P. (1993). The diet of sperm whales (*Physeter macrocephalus* Linnaeus 1758) off the Azores. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 339(1287), 67-82. <https://doi.org/10.1098/rstb.1993.0005>
- Collin, F. D., Durif, G., Raynal, L., Lombaert, E., Gautier, M., Vitalis, R., ... & Estoup, A. (2021). Extending approximate Bayesian computation with supervised machine learning to infer



- demographic history from genetic polymorphisms using DIYABC Random Forest. *Molecular Ecology Resources*, 21(8), 2598-2613. <https://doi.org/10.1111/1755-0998.13413>
- Corbin, L. J., Liu, A. Y. H., Bishop, S. C., & Woolliams, J. A. (2012). Estimation of historical effective population size using linkage disequilibria with marker data. *Journal of Animal Breeding and Genetics = Zeitschrift Für Tierzüchtung Und Züchtungsbiologie*, 129(4), 257–70. doi:10.1111/j.1439-0388.2012.01003.x
- Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., ... & 1000 Genomes Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*, 27(15), 2156-2158. <https://doi.org/10.1093/bioinformatics/btr330>
- De Leo, F. C., Smith, C. R., Rowden, A. A., Bowden, D. A., & Clark, M. R. (2010). Submarine canyons: hotspots of benthic biomass and productivity in the deep sea. *Proceedings of the Royal Society B: Biological Sciences*, 277(1695), 2783-2792. <https://doi.org/10.1098/rspb.2010.0462>
- de Jong, M. J., de Jong, J. F., Hoelzel, A. R., & Janke, A. (2021). SambaR: An R package for fast, easy and reproducible population-genetic analyses of biallelic SNP data sets. *Molecular Ecology Resources*, 21(4), 1369-1379. <https://doi.org/10.1111/1755-0998.13339>
- de Stephanis, R., Cornulier, T., Verborgh, P., Sierra, J. S., Gimeno, N. P., & Guinet, C. (2008). Summer spatial distribution of cetaceans in the Strait of Gibraltar in relation to the oceanographic context. *Marine Ecology Progress Series*, 353, 275-288. <https://doi.org/10.3354/meps07164>
- de Stephanis, R., Giménez, J., Carpinelli, E., Gutierrez-Exposito, C., & Cañadas, A. (2013). As main meal for sperm whales: plastics debris. *Marine pollution bulletin*, 69(1-2), 206-214. <https://doi.org/10.1016/j.marpolbul.2013.01.033>
- Dornburg, A., Brandley, M. C., McGowen, M. R., & Near, T. J. (2012). Relaxed clocks and inferences of heterogeneous patterns of nucleotide substitution and divergence time estimates across whales and dolphins (Mammalia: Cetacea). *Molecular Biology and Evolution*, 29(2), 721-736. <https://doi.org/10.1093/molbev/msr228>
- Drouot, V., Berube, M., Gannier, A., Goold, J. C., Reid, R. J., & Palsboll, P. J. (2004a). A note on genetic isolation of Mediterranean sperm whales (*Physeter macrocephalus*) suggested by mitochondrial DNA. *Journal of Cetacean Research and Management*, 6(1), 29-32.
- Drouot, V., Goold, J. C., & Gannier, A. (2004b). Regional diversity in the social vocalizations of sperm whale in the Mediterranean Sea. *Revue d'écologie*. <http://hdl.handle.net/2042/55620>
- Engelhaupt, D., Rus Hoelzel, A., Nicholson, C., Frantzis, A., Mesnick, S., Gero, S., ... & Mignucci-Giannoni, A. A. (2009). Female philopatry in coastal basins and male dispersion across the North Atlantic in a highly mobile marine species, the sperm whale (*Physeter macrocephalus*). *Molecular Ecology*, 18(20), 4193-4205. <https://doi.org/10.1111/j.1365-294X.2009.04355.x>
- Esteban, R., Verborgh, P., Gauffier, P., Giménez, J., Martín, V., Pérez-Gil, M., ... & de Stephanis, R. (2016). Using a multi-disciplinary approach to identify a critically endangered killer whale management unit. *Ecological Indicators*, 66, 291-300. <https://doi.org/10.1016/j.ecolind.2016.01.043>
- Excoffier, L., & Lischer, H. E. (2010). Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular ecology resources*, 10(3), 564-567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Fan, G., Zhang, Y., Liu, X., Wang, J., Sun, Z., Sun, S., ... & Liu, X. (2019). The first chromosome-level genome for a marine mammal as a resource to study ecology and evolution. *Molecular ecology resources*, 19(4), 944-956. <https://doi.org/10.1111/1755-0998.13003>
- Fais, A., Lewis, T. P., Zitterbart, D. P., Álvarez, O., Tejedor, A., & Aguilar Soto, N. (2016). Abundance and distribution of sperm whales in the Canary Islands: Can sperm whales in the

- archipelago sustain the current level of ship-strike mortalities?. *PLoS One*, 11(3), e0150660. <https://doi.org/10.1371/journal.pone.0150660>
- Ferreira, R., Steiner, L., Martín, V., Fusar Poli, F., Dinis, A., Kaufmann, M., ... & Alves, F. (2022). Unraveling site fidelity and residency patterns of sperm whales in the insular oceanic waters of Macaronesia. *Frontiers in Marine Science*, 9, 2546. doi: 10.3389/fmars.2022.1021635
- Feyrer, L. J., Bentzen, P., Whitehead, H., Paterson, I. G., & Einfeldt, A. (2019). Evolutionary impacts differ between two exploited populations of northern bottlenose whale (*Hyperoodon ampullatus*). *Ecology and evolution*, 9(23), 13567-13584. <https://doi.org/10.1002/ece3.5813>
- Fiori, C., Giancardo, L., Aïssi, M., Alessi, J., & Vassallo, P. (2014). Geostatistical modelling of spatial distribution of sperm whales in the Pelagos Sanctuary based on sparse count data and heterogeneous observations. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(S1), 41-49. <https://doi.org/10.1002/aqc.2428>
- Foskolos, I., Koutouzi, N., Polychronidis, L., Alexiadou, P., & Frantzis, A. (2020). A taste for squid: the diet of sperm whales stranded in Greece, East Mediterranean. *Deep Sea Research Part I: Oceanographic Research Papers*, 155, 103-164. <https://doi.org/10.1016/j.dsr.2019.103164>
- Frantzis, A., Alexiadou, P., Paximadis, G., Politi, E., Gannier, A., Corsini-Foka, M. (2003) Current knowledge on the cetacean fauna of the Greek seas. *Journal of Cetacean Research and Management* 5: 219–232.
- Frantzis, A., & Alexiadou, P. (2008). Male sperm whale (*Physeter macrocephalus*) coda production and coda-type usage depend on the presence of conspecifics and the behavioural context. *Canadian Journal of Zoology*, 86(1), 62-75. <https://doi.org/10.1139/Z07-114>
- Frantzis, A., Airoidi, S., Notarbartolo-di-Sciara, G., Johnson, C., & Mazzariol, S. (2011). Inter-basin movements of Mediterranean sperm whales provide insight into their population structure and conservation. *Deep Sea Research Part I: Oceanographic Research Papers*, 58(4), 454-459. <https://doi.org/10.1016/j.dsr.2011.02.005>
- Frantzis, A., Alexiadou, P., & Gkikopoulou, K. C. (2014). Sperm whale occurrence, site fidelity and population structure along the Hellenic Trench (Greece, Mediterranean Sea). *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(S1), 83-102. <https://doi.org/10.1002/aqc.2435>
- Frantzis, A., Leaper, R., Alexiadou, P., Prospathopoulos, A., & Lekkas, D. (2019). Shipping routes through core habitat of endangered sperm whales along the Hellenic Trench, Greece: Can we reduce collision risks?. *PloS one*, 14(2). <https://doi.org/10.1371/journal.pone.0212016>
- Frichot, E., & François, O. (2015). LEA: An R package for landscape and ecological association studies. *Methods in Ecology and Evolution*, 6(8), 925-929. <https://doi.org/10.1111/2041-210X.12382>
- García-Cortés L.A., Austerlitz F., de Cara MAR. (2019). An evaluation of the methods to estimate effective population size from measures of linkage disequilibrium. *J Evol Biol.* 32:267–277. <https://doi.org/10.1111/jeb.13411>
- Garibaldi, F., & Podestà, M., (2014). Stomach contents of a sperm whale (*Physeter macrocephalus*) stranded in Italy (Ligurian Sea, north-West Mediterranean). *J. Mar. Biol. Assoc. UK* 94, 1087–1091. <http://dx.doi.org/10.1017/S0025315413000428>
- Gaskin, D. E. (1982). *Ecology of whales and dolphins*. Heinemann
- Gaspari, S., Azzellino, A., Airoidi, S., & Hoelzel, A. R. (2007). Social kin associations and genetic structuring of striped dolphin populations (*Stenella coeruleoalba*) in the Mediterranean Sea. *Molecular Ecology*, 16(14), 2922-2933. <https://doi.org/10.1111/j.1365-294X.2007.03295.x>

- Gaspari, S., Scheinin, A., Holcer, D., Fortuna, C., Natali, C., Genov, T., ... & Moura, A. E. (2015). Drivers of population structure of the bottlenose dolphin (*Tursiops truncatus*) in the East Mediterranean Sea. *Evolutionary Biology*, 42(2), 177-190. <https://doi.org/10.1007/s11692-015-9309-8>
- Gauffier P, Moral Cendón M, Blasi A, Carpinelli E, Jiménez Torres C, Giménez Verdugo J, Esteban R, de Stephanis R, & Verborgh P. (2012). Winter presence of large cetaceans in the Strait of Gibraltar. 26th Annual Meeting European Cetacean Society, Galway, Ireland, 26–28 March 2012. <https://www.europeancetaceansociety.eu/sites/default/files/ecs2012galwayabstractbook.pdf>
- Gauffier, P., Verborgh, P., Giménez, J., Esteban, R., Sierra, J. M. S., & de Stephanis, R. (2018). Contemporary migration of fin whales through the Strait of Gibraltar. *Marine Ecology Progress Series*, 588, 215-228. <https://doi.org/10.3354/meps12449>
- Gauffier, P., Borrell, A., Silva, M. A., Víkingsson, G. A., López, A., Giménez, J., ... & Aguilar, A. (2020). Wait your turn, North Atlantic fin whales share a common feeding ground sequentially. *Marine Environmental Research*, 155, 104884. <https://doi.org/10.1016/j.marenvres.2020.104884>
- Giménez, J., Gómez-Campos, E., Borrell, A., Cardona, L., & Aguilar, A. (2013). Isotopic evidence of limited exchange between Mediterranean and East North Atlantic fin whales. *Rapid Communications in Mass Spectrometry*, 27(15), 1801-1806. <https://doi.org/10.1002/rcm.6633>
- Giménez, J., Louis, M., Barón, E., Ramírez, F., Verborgh, P., Gauffier, P., ... & de Stephanis, R. (2018). Towards the identification of ecological management units: a multidisciplinary approach for the effective management of bottlenose dolphins in the southern Iberian Peninsula. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 28(1), 205-215. <https://doi.org/10.1002/aqc.2814>
- Goncalves da Silva, A., Barendse, B., Kijas, J., England, P.R., & Hoelzel, A.R. (2019) Genomic data suggest environmental drivers of fish population structure in the deep sea; a case study for the orange roughy (*Hoplostethus atlanticus*) *J. Applied Ecol.* 57, 296-306. <https://doi.org/10.1111/1365-2664.13534>
- Gu, Z., Gu, L., Eils, R., Schlesner, M., & Brors, B. (2014). circlize implements and enhances circular visualization in R. *Bioinformatics*, 30(19), 2811-2812. <https://doi.org/10.1093/bioinformatics/btu393>
- Heller, R., Chikhi, L., & Siegmund, H. R. (2013). The confounding effect of population structure on Bayesian skyline plot inferences of demographic history. *PloS one*, 8(5), e62992. <https://doi.org/10.1371/journal.pone.0062992>
- Hoelzel, A.R. (1994). Ecology and genetics of whales and dolphins. *Annual Review of Ecology and Systematics*, 25:377-399 <https://www.jstor.org/stable/2097317>
- Hoelzel, A. R. (1998) Molecular analysis of populations; a practical approach. *Oxford, UK: Oxford University Press*.
- Hoelzel, A. R. (2009). Evolution of population genetic structure in marine mammal species. *Population genetics for animal conservation*. Cambridge University Press, Cambridge, 294-318.
- Hoelzel, A.R., Sarigol, F., Gridley, T., & Elwen, S.H. (2021) Natal origin of Namibian grey whale implies new distance record for in water migration. *Biol. Lett.* 17, 20210136 <https://doi.org/10.1098/rsbl.2021.0136>
- Holderegger, R., Schmidt, B. R., Grünig, C., Meier, R., Csencsics, D., Gassner, M., ... & Stapfer, A. (2020). Ready-to-use workflows for the implementation of genetic tools in conservation

- management. *Conservation Genetics Resources*, 12(4), 691-700.  
<https://doi.org/10.1007/s12686-020-01165-5>
- Hohenlohe, P. A., Funk, W. C., & Rajora, O. P. (2021). Population genomics for wildlife conservation and management. *Molecular Ecology*, 30(1), 62-82.  
<https://doi.org/10.1111/mec.15720>
- Ibáñez, C. M., Cubillos, L. A., Tafur, R., Argüelles, J., Yamashiro, C., & Poulin, E. (2011). Genetic diversity and demographic history of *Dosidicus gigas* (Cephalopoda: Ommastrephidae) in the Humboldt Current System. *Marine Ecology Progress Series*, 431, 163-171.  
<https://doi.org/10.3354/meps09133>
- Ibáñez, C. M., & Poulin, E. (2014). Genetic structure and diversity of squids with contrasting life histories in the Humboldt Current System. *Hidrobiologica*, 24(1), 1-10.
- Jacobsen, J. K., Massey, L., & Gulland, F. (2010). Fatal ingestion of floating net debris by two sperm whales (*Physeter macrocephalus*). *Marine Pollution Bulletin*, 60(5), 765-767.  
<https://doi.org/10.1016/j.marpolbul.2010.03.008>
- Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics*, 27(21), 3070-3071. <https://doi.org/10.1093/bioinformatics/btr521>
- Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: analysis of next generation sequencing data. *BMC bioinformatics*, 15(1), 1-13. <https://doi.org/10.1186/s12859-014-0356-4>
- Knutsen, H., Jorde, P.E., Albert, O.T., Hoelzel, A.R., & Stenseth, N.C. (2007). Population genetic structure in the North Atlantic Greenland halibut: influenced by oceanic current systems? *Can. J. Fish. Aquat. Sci.* 64, 857-866. <https://doi.org/10.1139/f07-070>
- Kraft, S., Pérez-Álvarez, M., Olavarria, C., & Poulin, E. (2020). Global phylogeography and genetic diversity of the long-finned pilot whale *Globicephala melas*, with new data from the southeastern Pacific. *Scientific reports*, 10(1), 1-13. <https://doi.org/10.1038/s41598-020-58532-3>
- Lambeck, K., & Purcell, A. (2005). Sea-level change in the Mediterranean Sea since the LGM: model predictions for tectonically stable areas. *Quaternary Science Reviews*, 24(18-19), 1969-1988. <https://doi.org/10.1016/j.quascirev.2004.06.025>
- Larsen, A. H., Sigurjonsson, J., Oien, N., Vikingsson, G., & Palsboll, P. J. (1996). Population genetic analysis of mitochondrial and nuclear genetic loci in skin biopsies collected from central and north Eastern North Atlantic humpback whales (*Megaptera novaeangliae*): Population identity and migratory destinations. *Proc. R. Soc. Lond B.* 263:1611-1618
- Latch, E.K., Dharmarajan, G., Glaubitz, J.C. & Rhodes, O.E. (2006) Relative performance of Bayesian clustering software for inferring population substructure and individual assignment at low levels of population differentiation. *Cons. Genet.* 7: 295-302.  
<https://doi.org/10.1007/s10592-005-9098-1>
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... & Durbin, R. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics*, 25(16), 2078-2079.  
<https://doi.org/10.1093/bioinformatics/btp352>
- Lischer, H. E., & Excoffier, L. (2012). PGDSpider: an automated data conversion tool for connecting population genetics and genomics programs. *Bioinformatics*, 28(2), 298-299.  
<https://doi.org/10.1093/bioinformatics/btr642>
- Liu, X., & Fu, Y. X. (2020). Stairway Plot 2: demographic history inference with folded SNP frequency spectra. *Genome Biology*, 21(1), 1-9. <https://doi.org/10.1186/s13059-020-02196-9>
- Louis, M., Viricel, A., Lucas, T., Peltier, H., Alfonsi, E., Berrow, S., ... & Simon-Bouhet, B. (2014). Habitat-driven population structure of bottlenose dolphins, *Tursiops truncatus*, in the North-East Atlantic. *Molecular Ecology*, 23(4), 857-874. <https://doi.org/10.1111/mec.12653>



- Luck, G. W., Daily, G. C., & Ehrlich, P. R. (2003). Population diversity and ecosystem services. *Trends in Ecology & Evolution*, 18(7), 331-336. [https://doi.org/10.1016/S0169-5347\(03\)00100-9](https://doi.org/10.1016/S0169-5347(03)00100-9)
- Lyrholm, T., Leimar, O., & Gyllenstein, U. (1996). Low diversity and biased substitution patterns in the mitochondrial DNA control region of sperm whales: implications for estimates of time since common ancestry. *Molecular Biology and Evolution*, 13(10), 1318-1326. <https://doi.org/10.1093/oxfordjournals.molbev.a025578>
- Lyrholm, T., & Gyllenstein, U. (1998). Global matrilineal population structure in sperm whales as indicated by mitochondrial DNA sequences. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 265(1406), 1679-1684. <https://doi.org/10.1098/rspb.1998.0488>
- Manichaikul, A., Mychaleckyj, J. C., Rich, S. S., Daly, K., Sale, M., & Chen, W. M. (2010). Robust relationship inference in genome-wide association studies. *Bioinformatics*, 26(22), 2867-2873. <https://doi.org/10.1093/bioinformatics/btq559>
- Mazzariol, S., Di Guardo, G., Petrella, A., Marsili, L., Fossi, C. M., Leonzio, C., ... & Fernández, A. (2011). Sometimes sperm whales (*Physeter macrocephalus*) cannot find their way back to the high seas: a multidisciplinary study on a mass stranding. *PLoS One*, 6(5), e19417. <https://doi.org/10.1371/journal.pone.0019417>
- Mesnick, S. L., Taylor, B. L., Archer, F. I., Martien, K. K., Treviño, S. E., Hancock-Hanser, B. L., ... & Morin, P. A. (2011). Sperm whale population structure in the Eastern and central North Pacific inferred by the use of single-nucleotide polymorphisms, microsatellites and mitochondrial DNA. *Molecular Ecology Resources*, 11, 278-298. <https://doi.org/10.1111/j.1755-0998.2010.02973.x>
- Mikolajewicz, U. (2011). Modeling mediterranean ocean climate of the last glacial maximum. *Climate of the Past*, 7(1), 161-180. <https://doi.org/10.5194/cp-7-161-2011>
- Morin, P. A., Foote, A. D., Baker, C. S., Hancock-Hanser, B. L., Kaschner, K., Mate, B. R., ... & Alexander, A. (2018). Demography or selection on linked cultural traits or genes? Investigating the driver of low mtDNA diversity in the sperm whale using complementary mitochondrial and nuclear genome analyses. *Molecular ecology*, 27(11), 2604-2619. <https://doi.org/10.1111/mec.14698>
- Moura, A. E., Janse van Rensburg, C., Pilot, M., Tehrani, A., Best, P. B., Thornton, M., ... & Hoelzel, A. R. (2014). Killer whale nuclear genome and mtDNA reveal widespread population bottleneck during the last glacial maximum. *Molecular biology and evolution*, 31(5), 1121-1131. <https://doi.org/10.1093/molbev/msu058>
- Mrusczok, M.T., Violi, B., Fakhri, M., Calogero, G., Biasissi, E., Jaouhar, A., Scullion, A., & Würtz, M. (2021). Long-Distance Movements of North Atlantic Killer Whales (*Orcinus orca*) from Iceland via Spain and Italy to Lebanon. *Marine Mammal Science*, 1–10. <https://doi.org/10.1111/mms.12866>
- Mussi, B., Miragliuolo, A., Zucchini, A., & Pace, D. S. (2014). Occurrence and spatio-temporal distribution of sperm whale (*Physeter macrocephalus*) in the submarine canyon of Cuma (Tyrrhenian Sea, Italy). *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(S1), 59-70. <https://doi.org/10.1002/aqc.2460>
- Mussmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2019). BA3-SNPs: Contemporary migration reconfigured in BayesAss for next-generation sequence data. *Methods in Ecology and Evolution*, 10(10), 1808-1813. <https://doi.org/10.1111/2041-210X.13252>
- Natoli, A., Birkun, A., Aguilar, A., Lopez, A., & Hoelzel, A. R. (2005). Habitat structure and the dispersal of male and female bottlenose dolphins (*Tursiops truncatus*). *Proceedings of the*

- Royal Society B: Biological Sciences*, 272(1569), 1217-1226.  
<https://doi.org/10.1098/rspb.2005.3076>
- Notarbartolo di Sciara, G., Agardy, T., Hyrenbach, D., Scovazzi, T., & Van Klaveren, P. (2008). The Pelagos Sanctuary for Mediterranean marine mammals. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 18(4), pp.367–391. <https://doi.org/10.1002/aqc.855>
- Notarbartolo di Sciara, G. (2014). Sperm whales, *Physeter macrocephalus*, in the Mediterranean Sea: a summary of status, threats, and conservation recommendations. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(S1), 4-10.  
<https://doi.org/10.1002/aqc.2409>
- Onoufriou, A. B., Gaggiotti, O. E., de Soto, N. A., McCarthy, M. L., Morin, P. A., Rosso, M., ... & Carroll, E. L. (2022). Biogeography in the deep: Hierarchical population genomic structure of two beaked whale species. *Global Ecology and Conservation*, e02308.  
<https://doi.org/10.1016/j.gecco.2022.e02308>
- Pace, D.S., Miragliuolo, A., & Mussi, B. (2008) Behaviour of a social unit of sperm whales (*Physeter macrocephalus*) entangled in a driftnet off Capo Palinuro (Southern Tyrrhenian Sea, Italy). *Journal of Cetacean Research and Management* 10: 131–135.
- Palumbi, S. R., & Baker, C. S. (1994). Contrasting population structure from nuclear intron sequences and mtDNA of humpback whales. *Mol. Biol. Evol.* 11:426-435.  
<https://doi.org/10.1093/oxfordjournals.molbev.a040115>
- Palumbi, S. R. (2004). Marine reserves and ocean neighbourhoods: The spatial scale of marine populations and their management. *Annual Rev. of Env. And Resources*, 29:31–68.  
<https://doi.org/10.1146/annurev.energy.29.062403.102254>
- Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, 35(3), 526-528.  
<https://doi.org/10.1093/bioinformatics/bty633>
- Patarnello, T., Volckaert, F. A., & Castilho, R. (2007). Pillars of Hercules: is the Atlantic–Mediterranean transition a phylogeographical break?. *Molecular ecology*, 16(21), 4426-4444. <https://doi.org/10.1111/j.1365-294X.2007.03477.x>
- Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., ... & Reich, D. (2012). Ancient admixture in human history. *Genetics*, 192(3), 1065-1093.  
<https://doi.org/10.1534/genetics.112.145037>
- Pavan, G., Hayward, T.J., Borsani, J.F., Priano, M., Manghi, M., Fossati, C., & Gordon, J., (2000). Time patterns of sperm whale codas recorded in the Mediterranean Sea 1985–1996. J. Acoust. Soc. Am. 107, 3487–3495. <http://dx.doi.org/10.1121/1.429419>.
- Peres dos Santos, R., Martins, R., Chaiko, A., Cheeseman, T., Jones, L. S., & Wenzel, F. W. (2022). First humpback whale (*Megaptera novaeangliae*) re-sighting between Azorean waters and the Barents Sea (Murman coast, northWestern Russia). *Polar Biology*, 1-5.  
<https://doi.org/10.1007/s00300-021-02998-y>
- Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PloS one*, 7(5), e37135. <https://doi.org/10.1371/journal.pone.0037135>
- Pickrell, J., & Pritchard, J. (2012). Inference of population splits and mixtures from genome-wide allele frequency data. *Nature Precedings*, 1-1. <https://doi.org/10.1038/npre.2012.6956.1>
- Pirotta, E., Matthiopoulos, J., MacKenzie, M., Scott-Hayward, L., & Rendell, L. (2011). Modelling sperm whale habitat preference: a novel approach combining transect and follow data. *Marine Ecology Progress Series*, 436, 257-272. <https://doi.org/10.3354/meps09236>
- Pirotta, E., Brotons, J. M., Cerdà, M., Bakkens, S., & Rendell, L. E. (2020a). Multi-scale analysis reveals changing distribution patterns and the influence of social structure on the habitat use of an endangered marine predator, the sperm whale *Physeter macrocephalus* in the Western

- Mediterranean Sea. *Deep Sea Research Part I: Oceanographic Research Papers*, 155, 103169. <https://doi.org/10.1016/j.dsr.2019.103169>
- Pirotta, E., Vighi, M., Brotons, J. M., Dillane, E., Cerdà, M., & Rendell, L. (2020b). Stable isotopes suggest fine-scale sexual segregation in an isolated, endangered sperm whale population. *Marine Ecology Progress Series*, 654, 209-218. <https://doi.org/10.3354/meps13502>
- Pirotta, E., Carpinelli, E., Frantzis, A., Gauffier, P., Lanfredi, C., Pace, D.S. & Rendell, L.E. (2021). *Physeter macrocephalus* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 2021: e.T16370739A50285671. <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T16370739A50285671.en>. Accessed on 20 January 2023.
- Puncher, G. N., Cariani, A., Maes, G. E., Van Houdt, J., Herten, K., Cannas, R., ... & Tinti, F. (2018). Spatial dynamics and mixing of bluefin tuna in the Atlantic Ocean and Mediterranean Sea revealed using next-generation sequencing. *Molecular Ecology Resources*, 18(3), 620-638. <https://doi.org/10.1111/1755-0998.12764>
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., ... & Sham, P. C. (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *The American journal of human genetics*, 81(3), 559-575. <https://doi.org/10.1086/519795>
- Quetglas, A., Carbonell, A., & Sanchez, P. (2000). Demersal continental shelf and upper slope cephalopod assemblages from the Balearic Sea (north-Western Mediterranean). Biological aspects of some deep-sea species. *Estuarine, Coastal and Shelf Science*, 50(6), 739-749. <https://doi.org/10.1006/ecss.1999.0603>
- Ralph, P., & Coop, G. (2010). Parallel adaptation: one or many waves of advance of an advantageous allele?. *Genetics*, 186(2), 647-668. <https://doi.org/10.1534/genetics.110.119594>
- Reinert, K., Langmead, B., Weese, D., & Evers, D. J. (2015). Alignment of next-generation sequencing reads. *Annual review of genomics and human genetics*, 16, 133-151. <https://doi.org/10.1146/annurev-genom-090413-025358>
- Rendell, L., Mesnick, S., Dalebout, M., Burtenshaw, J., & Whitehead, H. (2012) Can genetic differences explain vocal dialect variation in sperm whales, *Physeter macrocephalus*? *Behavior Genetics*, 42, 332–343. <https://doi.org/10.1007/s10519-011-9513-y>
- Rendell, L., Simiao, S., Brotons, J.M., Airoidi, S., Fasano, D., & Gannier, A. (2014). Abundance and movements of sperm whales in the Western Mediterranean basin. *Aquat. Conserv. Mar. Freshwat. Ecosyst.* 24, 31–40. <https://doi.org/10.1002/aqc.2426>
- Rendell, L., & Frantzis, A. (2016). Mediterranean sperm whales, *Physeter macrocephalus*: the precarious state of a lost tribe. *Advances in marine biology*, 75, 37-74. <https://doi.org/10.1016/bs.amb.2016.08.001>
- Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2: Analytical methods for paired-end sequencing improve RADseq-based population genomics. *Molecular Ecology*, 28(21), 4737-4754. <https://doi.org/10.1111/mec.15253>
- Silva M., Prieto R., Cascão I., Seabra M., Machete M., Baumgartner M., et al. (2014). Spatial and temporal distribution of cetaceans in the mid-Atlantic waters around the Azores. *Mar. Biol. Res.* 10 (2), 123–137. doi: 10.1080/17451000.2013.793814
- Smith, B. L., Lu, C. P., García-Cortés, B., Viñas, J., Yeh, S. Y., & Alvarado Bremer, J. R. (2015). Multilocus Bayesian estimates of intra-oceanic genetic differentiation, connectivity, and admixture in Atlantic swordfish (*Xiphias gladius* L.). *PLoS One*, 10(6), e0127979. <https://doi.org/10.1371/journal.pone.0127979>

- Stevick, P. T., Neves, M. C., Johansen, F., Engel, M. H., Allen, J., Marcondes, M. C., & Carlson, C. (2011). A quarter of a world away: female humpback whale moves 10 000 km between breeding areas. *Biology letters*, 7(2), 299-302. <https://doi.org/10.1098/rsbl.2010.0717>
- Sved, J. a, & Feldman, M. W. (1973). Correlation and probability methods for one and two loci. *Theoretical Population Biology*, 4(1), 129–132. doi:10.1016/0040-5809(73)90008-7
- Taylor, B. L., Chivers, S. J., Larese, J., & Perrin, W. F. (2007). Generation length and percent mature estimates for IUCN assessments of cetaceans. *NOAA, NMFS, Southwest Fisheries Science Center Administrative Report LJ-07–01*, 21. <http://hdl.handle.net/1834/41281>
- Taylor, B.L., Baird, R., Barlow, J., Dawson, S.M., Ford, J., Mead, J.G., Notarbartolo di Sciara, G., Wade, P. & Pitman, R.L. (2019). *Physeter macrocephalus* (amended version of 2008 assessment). *The IUCN Red List of Threatened Species* 2019: e.T41755A160983555.
- Teloni, V., (2005). Patterns of sound production in diving sperm whales in the north Western Mediterranean. *Mar. Mammal Sci.* 21, 446–457. <http://dx.doi.org/10.1111/j.1748-7692.2005.tb01243.x>
- Tepsich, P., Rosso, M., Halpin, P. N., & Moulins, A. (2014). Habitat preferences of two deep-diving cetacean species in the northern Ligurian Sea. *Marine Ecology Progress Series*, 508, 247-260. <https://doi.org/10.3354/meps10851>
- Thompson, K. F., Patel, S. E. L. I. N. A., Baker, C. S., Constantine, R. O. C. H. E. L. L. E., & Millar, C. D. (2016). Bucking the trend: genetic analysis reveals high diversity, large population size and low differentiation in a deep ocean cetacean. *Heredity*, 116(3), 277-285. <https://doi.org/10.1038/hdy.2015.99>
- Thunell, R. C., & Williams, D. F. (1989). Glacial-Holocene salinity changes in the Mediterranean Sea: hydrographic and depositional effects. *Nature*, 338(6215), 493-496. <https://doi.org/10.1038/338493a0>
- Tonay, A. M., Öztürk, A. A., Salman, A., Dede, A., Danyer, I. A., Danyer, E., & Öztürk, B. Stranding records of sperm whale (*Physeter macrocephalus*) on the Turkish coast in 2019-2020 with a note on the opportunistic sampling of stomach content. *J. Black Sea/Mediterranean Environment*, 27, 3: 281-293. [https://blackmeditjournal.org/wp-content/uploads/1-2021-3\\_281-293.pdf](https://blackmeditjournal.org/wp-content/uploads/1-2021-3_281-293.pdf)
- Torreblanca, E., Báez, J. C., Real, R., Macías, D., García-Barcelona, S., Ferri-Yañez, F., & Camiñas, J. A. (2022). Factors associated with the differential distribution of cetaceans linked with deep habitats in the Western Mediterranean Sea. *Scientific Reports*, 12(1), 1-16. <https://doi.org/10.1038/s41598-022-14369-6>
- Van Cise, A. M., Baird, R. W., Baker, C. S., Cerchio, S., Claridge, D., Fielding, R., ... & Morin, P. A. (2019). Oceanographic barriers, divergence, and admixture: phylogeography and taxonomy of two putative subspecies of short-finned pilot whale. *Molecular Ecology*, 28(11), 2886-2902. <https://doi.org/10.1111/mec.15107>
- Violi, B., Verga, A., Jones, L. S., Calogero, G., Soldano, G., Cheeseman, T., & Wenzel, F. W. (2021). A wanderer in the Mediterranean Sea: the case of a humpback whale (*Megaptera novaeangliae*) from the West Indies. *Aquatic Mammals*, 47(6), 599-611. <https://doi.org/10.1578/AM.47.6.2021.599>
- Westbury, M. V., Cabrera, A. A., Rey-Iglesia, A., De Cahsan, B., Duchêne, D. A., Hartmann, S., & Lorenzen, E. D. (2022). A genomic assessment of the marine-speciation paradox within the toothed whale superfamily Delphinoidea. *bioRxiv*, 2020-10. <https://doi.org/10.1101/2020.10.23.352286>
- Whitehead, H. (2002). Estimates of the current global population size and historical trajectory for sperm whales. *Marine Ecology Progress Series*, 242, 295-304. <http://doi:10.3354/meps242295>



- Whitehead, H., Antunes, R., Gero, S., Wong, S. N., Engelhaupt, D., & Rendell, L. (2012). Multilevel societies of female sperm whales (*Physeter macrocephalus*) in the Atlantic and Pacific: why are they so different?. *International Journal of Primatology*, 33(5), 1142-1164. <https://doi.org/10.1007/s10764-012-9598-z>
- Whitehead, H. (2017). Gene–culture coevolution in whales and dolphins. *Proceedings of the National Academy of Sciences*, 114(30), 7814-7821. <https://doi.org/10.1073/pnas.1620736114>
- Whitehead, H. (2018). Sperm whale: *Physeter macrocephalus*. In *Encyclopedia of marine mammals* (pp. 919-925). Academic Press.
- Whitehead, H., & Shin, M. (2022). Current global population size, post-whaling trend and historical trajectory of sperm whales. *Scientific Reports*, 12(1), 1-12. <https://doi.org/10.1038/s41598-022-24107-7>
- Winkelmann, I., Campos, P. F., Strugnelli, J., Cherel, Y., Smith, P. J., Kubodera, T., ... & Gilbert, M. T. P. (2013). Mitochondrial genome diversity and population structure of the giant squid *Architeuthis*: genetics sheds new light on one of the most enigmatic marine species. *Proceedings of the Royal Society B: Biological Sciences*, 280(1759), 20130273. <https://doi.org/10.1098/rspb.2013.0273>
- Würtz, M., & Simard, F. (2007). Following the food chain: an ecosystem approach to pelagic protected areas in the Mediterranean by mean of cetacean presence. *US Rapport de la Commission internationale pour la Mer Méditerranée*.
- Xue, D. X., Wang, H. Y., Zhang, T., & Liu, J. X. (2014). Population genetic structure and demographic history of *Atrina pectinata* based on mitochondrial DNA and microsatellite markers. *PLoS ONE*, 9, e95436. <https://doi.org/10.1371/journal.pone.0095436>

### Data Accessibility

Sequences associated with ddRADseq analyses are deposited at Genbank under BioProject accession PRJNA936197. ddRADseq genotype files and relevant code are provided on Dryad at: <https://doi.org/10.5061/dryad.dv41ns233>. There are no restrictions on data availability.

## TABLES

Table 1. Pairwise  $F_{st}$  values calculated using Arlequin v3.5 and based on dataset A (below the diagonal) and dataset B (above the diagonal). \*values are significant at  $p=0.000$  (to the resolution provided by the program). Cell colour corresponds to comparisons between the Atlantic and Mediterranean (dark blue) or within the Mediterranean (light blue).

	Atlantic	West Mediterranean	East Mediterranean
Atlantic	-	0.03009*	0.04085*
West Mediterranean	0.03099*	-	0.01276*
East Mediterranean	0.03583*	0.01203*	-

Table 2. Migration rates between Atlantic and Mediterranean populations using dataset A (above) and dataset B (below). Rates with standard deviations among parentheses defined as the proportion of individuals in a ‘current’ population (in rows) that are migrants derived from a ‘source’ population (in column), per generation. Values inferred in BayesAss3-SNPs.

Current \ Source	Atlantic	Mediterranean
	Atlantic	Mediterranean
Atlantic	0.9916 (0.0081)	0.0084 (0.0081)
	0.9903 (0.0094)	0.0097 (0.0094)
Mediterranean	0.0063 (0.0044)	0.9937 (0.0044)
	0.0088 (0.0074)	0.9912 (0.0074)

Table 3. Migration rates between Atlantic West Mediterranean and East Mediterranean putative populations using dataset A (values above) and dataset B (values below). Rates with standard deviations among parenthesis defined as the proportion of individuals in a ‘current’ population (in rows) that are migrants derived from a ‘source’ population in column, per generation. Values inferred in BayesAss3-SNPs.

Current \ Source	Atlantic	W Mediterranean	E Mediterranean
	Atlantic	W Mediterranean	E Mediterranean
Atlantic	0.9757 (0.0136)	0.0081 (0.0079)	0.0162 (0.0113)
	0.9810 (0.0129)	0.0191 (0.0130)	0.0094 (0.0091)
W Mediterranean	0.0042 (0.0041)	0.9639 (0.0114)	0.0320 (0.0108)
	0.0129 (0.0123)	0.9826 (0.0103)	0.0101 (0.0079)
E Mediterranean	0.0117 (0.0114)	0.2467 (0.0293)	0.7416 (0.0280)
	0.0099 (0.0071)	0.2355 (0.0759)	0.7516 (0.0755)

Table 4:  $F_3$  statistic among Atlantic West Mediterranean and East Mediterranean putative populations.

<b>pop1</b>		<b>pop2</b>	<b>pop3</b>	<b>est</b>	<b>se</b>	<b>z</b>	<b>P-value</b>
Atlantic	W Mediterranean	E Mediterranean		0.0018	6.25 E-05	29.7818	p<0.00001
W Mediterranean	E Mediterranean		Atlantic	-7.9 E-05	3.16 E-05	-2.4851	p=0.00459
E Mediterranean	Atlantic	W Mediterranean		0.0009	4.58 E-05	19.9024	p<0.00001

## Figure Captions:

Figure 1: Population Structure analyses. a) Sampling locations, b) Principal coordinate analyses for dataset A (4,422 SNPs) and c) Principal coordinate analyses for dataset B (12,616 SNPs); d) and e) principal coordinate analyses on samples using dataset A or B (respectively) restricted to the Mediterranean. Eastern basin samples are represented by triangles.

Figure 2: a) LEA plots of population structure including all samples for  $K=2-4$ . b) Estimated proportions of admixture for  $K=2$ . Atlantic assignment is in blue and Mediterranean in green. Estimates developed in Admixture and visualized in R. c) LEA plots of population structure including only Mediterranean samples for  $K=2-4$ .

Figure 3: Gene flow analyses. Circosplots showing migration rates between populations as calculated by BayesAss3-SNPs. a) Atlantic (blue) and Mediterranean (green) putative populations. b) Atlantic (blue) and Western (orange) and Eastern (green) Mediterranean putative populations. c)  $F_3$  statistic values using Atlantic, west Mediterranean and east Mediterranean putative populations. All values are significant at the  $p<0.05$  level based on the Z-scores shown. d) ML tree of Atlantic generated with TreeMix, Western and Eastern Mediterranean populations with inferred migration edges.

Figure 4: Demography analyses using stairway plots. a) Atlantic subset; b) western Mediterranean subset; c) eastern Mediterranean subset. Last glacial period (LGP) and Last glacial maximum (LGM) indicated by shading. Red line indicates median values. Black lines indicate 12.5% and 87.5% percentile values, and grey margins indicate 2.5% and 97.5% percentile values.









