

BOCAGIANA

Museu Municipal do Funchal (História Natural)

Madeira

31.XII.2009

No. 230

**FIRST RECORD OF THE DEEP-SEA RED CRAB
Chaceon inglei (DECAPODA: GERYONIDAE)
FROM MADEIRA AND THE CANARY ISLANDS
(NORTHEASTERN ATLANTIC OCEAN) ***

R. ARAÚJO ¹, M. BISCOITO ¹, J. I. SANTANA ² & J. A. GONZÁLEZ ²

With 2 figures and 1 table

ABSTRACT. The deep-sea red crab *Chaceon inglei* Manning & Holthuis, 1989 is recorded for the first time from the waters of Madeira and the Canary Islands. This is the second species of the genus *Chaceon* to occur in Madeira Island and the third in the Canaries. These collections set a new bathymetric record for this species and represent the southern limit of its distribution in the eastern Atlantic Ocean.

KEY WORDS: Crustacea, Decapoda, Geryonidae, *Chaceon inglei*, new record, Madeira, Canary Islands, NE Atlantic Ocean.

¹ Museu Municipal do Funchal (História Natural), Rua da Mouraria, 31, 9000-047 Funchal, Madeira, Portugal. E-mail: ricardo.araujo@cm-funchal.pt

² Grupo de Biología Pesquera, Instituto Canario de Ciencias Marinas (ICCM – ACIISI), P. O. Box 56, Telde, 35200 Las Palmas, Canary Islands, Spain.

* Contribution no. 150 of the Marine Biology Station of Funchal.

RESUMO. Neste artigo é assinalada pela primeira vez para as ilhas da Madeira e das Canárias uma nova espécie de caranguejo da fundura, *Chaceon inglei* Manning & Holthuis, 1989. Esta é a segunda espécie do género *Chaceon* assinalada para a ilha da Madeira e a terceira para as ilhas Canárias. Estas colheitas estabelecem um novo limite Sul de distribuição no Oceano Atlântico nordeste e constituem as mais profundas desta espécie registadas até ao presente.

INTRODUCTION

The genus *Chaceon* Manning & Holthuis, 1989 (Decapoda: Geryonidae) is represented in the NE Atlantic Ocean and the Mediterranean Sea by five species: *Chaceon affinis* (A. Milne-Edwards & Bouvier, 1894), *C. gordonaee* (Ingle, 1985), *C. inglei* Manning & Holthuis, 1989, *C. maritae* (Manning & Holthuis, 1981) and *C. mediterraneus* Manning & Holthuis, 1989 (DAWSON & WEBBER, 1991; d'UDEKEM d'ACOZ, 1999). In the Canary Islands waters only two species have been recorded: *Chaceon affinis* and *C. maritae* (GONZÁLEZ, 1995; GONZÁLEZ *et al.*, 1996) and in Madeira only *C. affinis* was known to occur previously to the present report (TÜRKAY, 1976). Recent collections in both archipelagos have shown that another species, *C. inglei*, is also present and therefore it is herein recorded for the first time.

MATERIAL AND METHODS

All specimens were collected with bottom traps either of the model described by BISCOITO (1993) or a cubic experimental model with 1m side (GONZÁLEZ *et al.*, 1988). Traps were laid down for 24 hours at depths between 1500 and 2500 m and baited with salted mackerel. The specimens were preserved in 70% ethanol and deposited in the collections of MMF (Museu Municipal do Funchal (História Natural)) and ICCM (Instituto Canario de Ciencias Marinas).

All measurements were taken to the nearest 0.1 mm and follow ATTRILL *et al.* (1991).

Abbreviations used: CW – carapace width; CL – carapace length; RCHW – right chela width; RCHL – right chela length; LCHW – left chela width; LCHL – left chela length; W – weight.

RESULTS

***Chaceon inglei* Manning & Holthuis, 1989 (Figs. 1 and 2, Table 1)**

Material examined: MMF25576, 53.2 mm CW, ♂, off Funchal, Madeira, 32° 34' N 16° 52' W, ca. 1700 m, R/V Auriga, 09.07.1992, bottom trap; MMF35981, 62.7 mm CW, ♂, off Funchal, Madeira, 32° 30' N 16° 50' W, ca. 2500 m, RECPROFMAD cruise, R/V Arquipélago,

20.10.2004, bottom trap; ICCM-0063, 46.2 mm CW, non-ovigerous ♀, Arguineguín, Gran Canaria, Canary Islands, 27° 38' N 15° 57' W, ca. 2156 m, RECPROFCAN-4 cruise, F/V Mary Nere, stn. 2, 20.10.2004, bottom trap.



Fig. 1 - Male specimen of *Chaceon inglei* from Madeira (MMF35981), dorsal view.



Fig. 2 - Female specimen of *Chaceon inglei* from the Canaries (ICCM0063), ventral view.

TABLE 1 - Measurements (in mm) and weights (in g) of the specimens of *C. inglei* studied.

	MMF25576	MMF35981	ICCM0063
CW	53.2	62.7	46.2
CL	48.2	51.9	39.4
RCHW	13.6	15.2	8.9
RCHL	39.6	35.1	25.8
LCHW	11.8	13.7	9.3
LCHL	39.0	42.9	26.9
W	49.8	71.7	27.4

DISCUSSION

All the specimens of *C. inglei* collected are in agreement with the descriptions and illustrations given by MANNING & HOLTHUIS (1989). The specimen in Fig. 1 shows the well developed antero-lateral spines on the carapace, as pointed out by MANNING & HOLTHUIS (1989).

C. inglei was recorded in the northeastern Atlantic Ocean from 61° 33' N, south of Iceland, to 39° N, off the Azores including off the west of Scotland, off the southwest of England, the Bay of Biscay and off Vigo, Spain, at a depth range of 1640-2245 m (HANSEN, 1908 as *Geryon affinis*; INGLE, 1985 as *Geryon gordonaee*; MANNING & HOLTHUIS, 1989; DAWSON & WEBBER, 1991; d'UDEKEM d'ACOZ, 1999).

This species is here recorded for the first time from the archipelagos of Madeira and Canaries, the latter setting a new southern limit (27° 38' N) of its distribution. Specimens studied were collected between 1700 and 2500 m. The specimen (MMF35981) captured at 2500 m in Funchal Bay sets a new bathymetric record for this species.

ACKNOWLEDGEMENTS

The authors are indebted to Commander Nuno Bustorff and crew of the N. R. P. Auriga and captains and crews of the R/V Arquipélago and F/V Mary Nere. This research was co-funded by the EU FEDER, Interreg III-B, project PESCPROF-1 (MAC/4.2/M12).

REFERENCES

- ATTRILL, M. J., R. G. HARTNOLL & A. RICE:
1991. Aspects of the biology of the deep-sea crab *Geryon trispinosus* from the Porcupine Seabight. *Journal of the Marine Biological Association of the U. K.*, **71**: 311-328.
- BISCOITO, M.:
1993. An account on the shrimps of the family Pandalidae (Crustacea, Decapoda, Caridea) in Madeiran waters. In: Proceedings of the 5th Symposium "Fauna and Flora of the Cape Verde Islands", Leiden, 4-7 October, 1989. *Courier Forschungs-Institut Senckenberg*, **159**: 321-325.
- DAWSON, E. W. & W. R. WEBBER:
1991. The deep-sea red crab *Chaceon* ("Geryon"): a guide to information and a reference list of the family Geryonidae. *Miscellaneous series of the National Museum of New Zealand*, **24**: pp. 83. National Museum of New Zealand: Wellington.
- GONZÁLEZ, J. A.:
1995. *Catálogo de los Crustáceos Decápodos de las islas Canarias*. pp. 282. Publicaciones Turquesa: Santa Cruz de Tenerife.
- GONZÁLEZ, J. A., I. J. LOZANO, M. A. CALDENTEY, J. I. SANTANA, J. A. GÓMEZ & R. CASTILLO:
1988. Resultados de la campaña de prospección pesquera CANARIAS 85. *Informes Técnicos del Instituto Español de Oceanografía*, **57**: pp. 93.
- GONZÁLEZ, J. A., J. I. SANTANA & V. FERNÁNDEZ-VERGAZ:
1996. The family Geryonidae (Decapoda, Brachyura) in the Canary Islands. *Crustaceana*, **69** (5): 627-635.
- HANSEN, H. J.:
1908. Crustacea Malacostraca. I. – *The Danish Ingolf-Expedition*, **3** (2): 1-120, pls. 1-5.
- INGLE, R. W.:
1985. *Geryon gordonaee* sp. nov. (Decapoda, Brachyura, Geryonidae) from the Northeastern Atlantic Ocean. *Crustaceana*, **48** (1): 88-98.

MANNING, R. B. & L. B. HOLTHUIS:

1981. West African Brachyuran Crabs (Crustacea: Decapoda). *Smithsonian Contribution to Zoology*, **306**: i-xii + 1-379.
1989. Two new genera and nine new species of geryonid crabs (Crustacea, Decapoda, Geryonidae). *Proceedings of the Biological Society of Washington*, **102** (1): 50-77.

TÜRKAY, M.:

1976. Die Madeirischen Brachyuren des Museu Municipal do Funchal und des Forschungs-Instituts Senckenberg I. Familien: Dromiidae, Homolidae, Calappidae, Leucosiidae, Cancridae, Portunidae, Xanthidae, Geryonidae, Gonoplacidae und Palicidae (Crustacea: Decapoda). *Boletim do Museu Municipal do Funchal*, **30**: 57-74.

UDEKEM d'ACOZ, C. (d'):

1999. Inventaire et distribution des crustacés décapodes de l'Atlantique nord-oriental, de la Méditerranée et des eaux continentales adjacentes au nord de 25° N. *Patrimoines naturels* (M. N. H. N. / S. P. N.), **40**: pp. 383.