

CULTIVATED SEA BASS INGEST PLASTIC PARTICLES

Stefanie Reinold*¹, Alicia Herrera¹, Francesco Saliu², Carlos Hernández-González³,
Ico Martínez¹, Marina Lasagni² and May Gómez¹

¹ Marine Ecophysiology Group (EOMAR), Iu-ECOQUA, Universidad de Las Palmas de Gran Canaria, Campus Universitario de Tafira, 35017, Canary Islands, Spain
stefanie.reinold101@alu.ulpgc.es, alicia.herrera@ulpgc.es, ico.martinez@ulpgc.es, may.gomez@ulpgc.es

² Earth and Environmental Science Department, University of Milano Bicocca, Piazza della Scienza 1, 20126 Milano, Italy
francesco.saliu@unimib.it, marina.lasagni@unimib.it

³ Centro Oceanográfico de Canarias, Instituto Español de Oceanografía, Santa Cruz de Tenerife, Canary Islands, Spain
carlos.hernandez@ieo.es

Abstract: The presence of microplastics (MPs) in the marine environment is a concerning topic due to the ecotoxicological effects and possible seafood contamination. Data is needed to evaluate human exposure and assess risks, in the context of a healthy and beneficial seafood consumption. While microplastic ingestion by wild fish has been reported since the early 70's, farmed fish are rarely investigated. Here, for the first time the presence of microplastics in fish cultivated in the coastal water of Tenerife (Canary Island, Spain) was evaluated. From 83 examined individuals, 65% displayed microplastics in their gastrointestinal tracts, with averages between 0.6 ± 0.8 (SD) and 2.7 ± 1.85 (SD) particles per fish. The total number of microplastics detected was 119. Fibres (81%) and fragments (12%) were the predominant shapes. FTIR analysis showed that fibres were mostly composed by Cellulose (55%) and Nylon (27%), whereas fragments by PE (25%) and PP (25%).

Key words: Pollution, Plastic, Microparticles, Aquaculture, Seafood, Canary Islands

Acknowledgments: We are grateful to the IEO for the help to process the samples of this study. This work was funded by the project RICOMAR (GOB-ESP2019-03) which supported the Marine Ecophysiology Group (EOMAR).