

International Society for Pediatric and Adolescent Diabetes Clinical Practice Consensus Guidelines 2024: Glycemic Targets

Martin de Bock^{a,b} Juliana Chizo Agwu^c Matt Deabreu^d Klemen Dovc^e
David M. Maahs^f M. Loredana Marcovecchio^g Farid H. Mahmud^h
Yeray Nóvoa-Medinaⁱ Leena Priyambada^j Carmel E. Smart^{k,l}
Linda A. DiMeglio^m

^aDepartment of Paediatrics, University of Otago, Christchurch, New Zealand; ^bDepartment of Paediatrics, Te Whatu Ora, Waitaha, New Zealand; ^cDepartment of Paediatrics, Wye Valley NHS Trust, Hereford, UK; ^dParent and Advocate of Child with Type One Diabetes, Toronto, ON, Canada; ^eUniversity Medical Centre Ljubljana, University Children's Hospital, Department of Endocrinology, Diabetes and Metabolic Diseases and University of Ljubljana Faculty of Medicine, Ljubljana, Slovenia; ^fDivision of Endocrinology, Department of Pediatrics, Stanford University, Stanford, CA, USA; ^gDepartment of Paediatrics, University of Cambridge and Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK; ^hDivision of Endocrinology, Department of Pediatrics, Hospital for Sick Children, University of Toronto, Toronto, ON, Canada; ⁱUniversity Institute of Biomedical and Healthcare Research (IUBS), University of Las Palmas de Gran Canaria, Las Palmas de Gran Canaria, Spain; ^jDepartment of Pediatric Endocrinology, Rainbow Children's Hospital, Hyderabad, India; ^kPaediatric Endocrinology and Diabetes, John Hunter Children's Hospital, Newcastle, NSW, Australia; ^lSchool of Health Sciences, University of Newcastle, Newcastle, NSW, Australia; ^mDivision of Pediatric Endocrinology and Diabetology, Department of Pediatrics, Riley Hospital for Children, Indiana University School of Medicine, Indianapolis, IN USA

Summary of What Is New or Different

1. This chapter recommends a target hemoglobin A1c (HbA1c) of $\leq 6.5\%$ (48 mmol/mol) for those who have access to advanced diabetes technologies like continuous glucose monitoring (CGM) and automated insulin delivery (AID). This target should be encouraged for all children and adolescents living with diabetes when safely achievable. In other settings, the HbA1c target is $\leq 7.0\%$ (53 mmol/mol). The emphasis on the lower HbA1c target is deliberate, based on the growing evidence that achieving HbA1c levels below previous targets can significantly reduce the risk of developing diabetes complications and that, when adequate technology and healthcare professional support are available, these lower glycemic targets can be safely achieved without increasing the risk of hypoglycemia or adding to the care burdens.
2. These glycemic goals may be individualized based on an assessment of the potential challenges for the person with diabetes and their caregivers.
3. The relationship between glycemic targets and rates of excess body weight is discussed.
4. The glycemic target chapter is also more succinct compared to previous iterations. This chapter is meant to provide updated guidance on overall glycemic management alongside other International Society for Pediatric and Adolescent Diabetes (ISPAD) consensus guidelines that provide further context and information, including specific populations (e.g., pre-school children, limited care settings) or situations (e.g., exercise).

Keywords

Glycemic targets · Technology · Burden · Diabetes-related complications

Abstract

The International Society for Pediatric and Adolescent Diabetes (ISPAD) guidelines represent a rich repository that serves as the only comprehensive set of clinical recommendations for children, adolescents, and young adults living with diabetes worldwide. This chapter builds on the 2022 ISPAD guidelines, and updates recommendations on the glycemic targets for children and adolescents living with diabetes. A new target for hemoglobin A1c (HbA1c) of $\leq 6.5\%$ (48 mmol/mol) is recommended for those who have access to advanced diabetes technologies like continuous glucose monitoring and automated insulin delivery. This target should be encouraged for all children and adolescents living with diabetes when safely achievable. In other settings, the HbA1c target is $\leq 7.0\%$ (53 mmol/mol).

© 2024 The Author(s).

Published by S. Karger AG, Basel

List of Abbreviations

AID:	automated insulin delivery
BMI:	body mass index
CGM:	continuous glucose monitoring
HbA1c:	hemoglobin A1c
ISPAD:	International Society for Pediatric and Adolescent Diabetes
SDS:	standard deviation score
SMBG:	self-monitored blood glucose
TIR:	time in range
TITR:	time in tight range

The Purpose of Setting Glycemic Targets

Setting glycemic targets for young persons living with diabetes is vital as optimizing glycemia can greatly minimize both immediate and long-term complications [1–6]. This is especially important for children and adolescents with diabetes as they face many years of managing the condition and lifetime consequences of dysglycemia [7]. Achieving healthier glycemic outcomes has significant benefits for healthcare systems and economic costs, underscoring the importance of targeting glycemic ranges to avert future complications [8]. Yet, it is also important to recognize the day-to-day

challenges glycemic targets can impose on persons living with diabetes, their caregivers, and even their healthcare team. These targets should be seen as part of a balanced approach to long-term management; at times flexibility will be necessary.

Recent data from diabetes registries show that while a minority of young people are meeting current glycemic targets, median HbA1c levels have gradually decreased over the past decades [9–13]. This progress can be attributed to several factors, including better, consistent communication of glycemic goals by healthcare teams, advancements in treatments like insulin analogs and CGM, a well-trained medical workforce, and the development and implementation of AID systems. Additionally, clinical benchmarking activities that use published glycemic targets in their reporting have been shown to be associated with improvement in whole population glycemic metrics [14–17]. However, challenges such as social determinants of health, limited resources for pediatric diabetes care, and restricted access to advanced treatments continue to impede broader achievement of target glycemia and contribute to health inequities [18–20].

For over 20 years, organizations like ISPAD, the American Diabetes Association (ADA), and the National Institute for Health and Care Excellence (NICE) in the UK have established and regularly updated glycemic targets based on evolving evidence. While lower glycemic targets are associated with improved glycemia [21], the path to achieving these targets can be challenging. In 2022, ISPAD set the HbA1c target at 7.0% (53 mmol/mol), with a provision for lower targets where feasible, particularly when there is access to gold standard treatments like AID [22]. Other societies have adopted lower targets. For instance, Sweden [23] and NICE [11] have lowered their targets to 6.5% (48 mmol/mol). Additionally, in 2022, ISPAD communicated that an HbA1c of 6.5% could be safely achieved for pre-school children [24]. The ADA recommends an HbA1c of $< 6.5\%$ as a reasonable goal if “it can be achieved without significant risk of hypoglycemia, or negatively impacts on well-being or undue burden of care,” but also allows flexibility, stating “individualized goals” with $< 7.0\%$ as the “appropriate” target [25].

Healthcare professionals and persons with diabetes have various tools for monitoring glycemia, including self-monitored blood glucose (SMBG) using a glucose meter, CGM and HbA1c. While HbA1c has traditionally been the standard measure, it also has limitations that CGM can address. Despite this, CGM is not yet accessible to all young people, meaning that SMBG and HbA1c




Fig. 1. Summary of glycemic metrics. *For those who do not have access to advanced technologies or unable to safely attain the lower target. †And/or where the pursuit of the lower target does not add burden such that quality of life is impacted. DCCT, diabetes control and complications trial; TITR, time in tight range.

measurements remain critical in many settings, including for benchmarking. Combining data from these sources can also offer the most comprehensive view of glycemia, and in turn, the best opportunity to tailor care to both the immediate and long-term needs of everyone.

Measures of Glycemia and Targets

For educational purposes, three methods of measuring glycemia and their targets are provided in Figure 1. This has been updated from 2022 to specifically emphasize the lower HbA1c target of 6.5% (48 mmol/mol) for those with access to advanced technologies or those who can safely reach that target without additional undue care demands.

Glycated Hemoglobin

- ISPAD recommends an HbA1c target of $\leq 6.5\%$ (48 mmol/mol) for those who can safely reach that target with the support of advanced technologies (CGM and AID) and/or where the pursuit of the lower target does not add burden such that quality of life is impacted [C]
- ISPAD recommends an HbA1c target of $\leq 7\%$ (53 mmol/mol) in all other scenarios [A]

Landmark trials have shown that the relationship between the developments of long-term complications

and glycemia is continuous and non-linear [2, 26]. These robust data justify an HbA1c target of $\leq 7\%$ (53 mmol/mol) (Category A evidence). However, evidence for a lower target of $\leq 6.5\%$ (48 mmol/mol) is still emerging as described below (Category C evidence). Overall, further research is needed to evaluate the outcomes of children achieving HbA1c levels of $\leq 6.5\%$. This should include more comprehensive studies and detailed mathematical modeling of existing data sets. Such analysis would help quantify the incremental improvements in complication rates and assess any potential increased risks when comparing the HbA1c thresholds of $\leq 7\%$ (53 mmol/mol) and $\leq 6.5\%$ (48 mmol/mol). Yet, at this time, communicating at a single threshold is a pragmatic approach to convey clear messaging that can be implemented in clinical practice. This threshold also facilitates the ability to benchmark outcomes, adding to the body of evidence demonstrating the safety and validity of lower target setting to improvements in glycemia [14–17]. Goal setting is important for reducing therapeutic inertia, and therefore, regular measuring of HbA1c (and other metrics) is an important foundation of clinical care – with an accepted convention of 3 monthly assessments, given that HbA1c reflects ~90 days of glycemia. Health care professionals and care teams should consistently and clearly communicate the importance of glycemic targets to optimize long-term health outcomes for youth with diabetes [21].

Refer to the ISPAD CPCG 2024: Diabetes Technologies: Insulin Delivery [27] for the evidence regarding the

efficacy and safety of advanced technologies, including cost-effectiveness [28, 29]. Current glycemic target recommendations are also based on the following observations:

Diabetic Retinopathy

Independent risk factors for developing diabetic retinopathy include not only glycemic management, but also age of onset and duration of diabetes [30, 31], which emphasizes the need to pursue optimal glycemic targets in young people with diabetes from diagnosis. Precursors to retinopathy are now measurable with optical coherence tomography angiography, and microvascular changes are seen in youth with suboptimal glycemic levels [32]. Critically, the lowest rates of diabetic retinopathy at the time of transition to adult services are seen where HbA1c is $\leq 6.5\%$ (48 mmol/mol) [33].

Diabetic Kidney Disease

Multiple studies confirm that lower HbA1c protects against the development of diabetic kidney disease [1–6]. In contrast to the development of diabetic retinopathy, Swedish registry data did not find a similar clear relationship for the development of increased albuminuria at the 6.5% (48 mmol/mol) HbA1c threshold [33]. There is evidence that some individuals from differing racial groups have different predispositions to developing complications early independently of glycemia and socioeconomic status [34, 35]. Indeed, observations from the general population showed that Native Americans at risk of developing type 2 diabetes and HbA1c in a pre-diabetes range (5.7–6.4% [39–47 mmol/mol]) had higher incidence of albuminuria and retinopathy than children with optimal HbA1c levels [36]. However, there is limited evidence supporting reductions in diabetic kidney disease onset and progression at HbA1c levels below 7% in type 1 diabetes.

Diabetic Neuropathy

While systematic reviews confirm the relationship between HbA1c and the development of diabetes neuropathy [36], there is, again, a paucity of data to derive a specific threshold that confers significant additional risk. An HbA1c $>7\%$ (53 mmol/mol) was associated with a higher risk of developing diabetic neuropathy in adolescents with type 1 diabetes for >5 years, but the risk associated with a lower threshold of 6.5% (48 mmol/mol) was not assessed [37]. Further data, or modeling, are needed to confirm the additional benefit of using a target of 6.5% (48 mmol/mol) for protecting against the development of diabetic neuropathy and other vascular complications.

Cognition

Optimal glycemia is associated with improved cognitive function, including memory, learning and attention [13, 38–41]. Yet, in settings where hypoglycemia-protective technologies such as CGM and AID are not available or adopted, there may be unacceptable risks of severe hypoglycemia and associated negative impact on cognitive function. Overall, with respect to cognition, a lower target of 6.5% should be carefully guided on an individual basis, particularly in situations with limited access to glucose monitoring and technologies that suspend insulin when glucose is low or projected to be low.

Burden

Recognizing and addressing the psychosocial and behavioral needs of youth with diabetes and their families is a key element of their care. The relationship between the demands of care and HbA1c is complex; for example, while some reports describe the challenges of diabetes technology, improved quality of life and glycemia have been consistently demonstrated [27]. Burden of care can be considered in the following domains; psychological, behavioral, social, quality of life, and economic. Avoidance of complications with lower HbA1c will lower the health cost of diabetes and can be safely achieved with advanced diabetes technologies, with proven health economic benefits. Adoption of these technologies is associated with potential barriers, which includes perceived and actual workload, physical discomforts, frustrations with technical glitches and alarms, as well as concerns about device size/visibility and stigma [42–44]. Potential benefits of diabetes technologies must also be recognized, with improvements in individual and parental health of life [45], as well as autonomy, greater flexibility in social activities and eating, improved sleep, and higher treatment satisfaction being reported [27].

Body Mass Index

Due to overweight and obesity being established as independent risk factors for developing cardiovascular disorders, any relationship between glycemic targets and body mass index (BMI) is an important consideration. This is especially relevant given that while international registries are demonstrating an overall improvement in HbA1c, there is also an increasing prevalence of higher BMI-standard deviation scores (SDS) [46, 47]. Earlier landmark studies have shown increased excess weight gain and obesity with intensive treatment [26]. However, there are apparent external influences on BMI independent of HbA1c, as is apparent with the U-shaped curve with the highest HbA1c observed in groups with unhealthy weight [48]. Some studies

have reported inverse associations between HbA1c and BMI-SDS [49–51]. However, other studies have found high BMI-SDS significantly related to higher HbA1c levels [47, 52], or reported children with type 1 diabetes can achieve glycemic targets on intensive insulin therapy without excessive weight gain [53, 54]. Use of advanced technologies such as AID is associated with improved glycemia without change in BMI [55, 56]. The emergence of increased overweight and obesity in the pediatric population living with diabetes underpins the importance of dietetic education, to ensure the benefits of improved glycemia are not offset by increasing BMI. These data highlight the need for additional research and innovative care approaches to address rising rates of overweight and obesity and the risk for premature cardiovascular disease in this vulnerable population.

Continuous Glucose Monitoring

- Children and adolescents with diabetes should strive for the following % time spent in the following glycemic ranges [B]:
>70% between 3.9 and 10 mmol/L (70–180 mg/dL),
<4% <3.9 mmol/L (70 mg/dL),
<1% <3.0 mmol/L (54 mg/dL),
<25% >10 mmol/L (180 mg/dL), and
<5% >13.9 mmol/L (250 mg/dL).
These times in range are chosen to align with previously published recommendations [45].
- It is recognized that time spent in range (TIR) may need to be >80% to achieve an optimal HbA1c target of $\leq 6.5\%$ (48 mmol/mol) [E].
- It is also recognized that an emerging metric requiring additional validation of >50% time in tight range (TTR) (3.9–8 mmol/L or 70–144 mg/dL) may allow for greater sensitivity for assessing changes in mean glucose and glucose variability (CV) [C].
- CGM-derived Glucose Management Index metrics can be considered a standalone method for evaluating glycemic outcomes [C].

Average sensor glucose, due to its strong correlation with HbA1c and its association with the risk of microvascular complications [57] and glycemic variability (which can predict hypoglycemia), is a key metric included in standardized CGM reports, known as ambulatory glucose profiles. When available, CGM targets can be considered an alternative or adjunct to HbA1c targets. As CGM technology is still relatively new, further research linking CGM data to the development of diabetes complications is needed to refine these glycemic targets with greater certainty.

Given the strong correlation between CGM metrics and HbA1c [58], it is reasonable in a clinical setting to use CGM along with the Glucose Management Index metric as a standalone method for setting glycemic targets. This

could be especially relevant in situations where CGM data are easier to obtain than an HbA1c (telehealth, limited access to HbA1c measurement).

TTR (3.9–8 mmol/L or 70–144 mg/dL) has recently been suggested as a new metric to characterize optimal glycemia [59, 60]. ISPAD has previously endorsed TTR as a metric in pre-schoolers with T1D, especially during the remission period [61]. TTR may be preferable to TIR when targeting lower HbA1c values since it may be more sensitive to glycemic changes at lower average sensor glucose levels, and it may also better characterize glucose variability [62].

Generally, TTR targets have been set at 50%; however, to reach the lower HbA1c target of $\leq 6.5\%$ (48 mmol/mol), an even greater TTR of > 55% may be needed [63]. While we await additional, high-quality evidence for this metric in youth with diabetes [64], as well as guidance on how best to message and use this metric for clinical care, it is logical that increasing time spent in a tight range may reflect lower glucose variability and reduce the long-term risk of complications. Importantly, when selecting any CGM metric to use in care settings, it is essential to adopt an educational framework that places the child and caregiver at its core and is sensitive to holistic health needs that extend beyond glucose levels.

Self-Monitored Blood Glucose

- In the absence of CGM, pre-prandial SMBG targets should be 4.0–8.0 mmol/L (70–144 mg/dL), and 4.0–10 mmol (70–180 mg/dL) post-prandial [B].
- In the absence of CGM, SMBG should be undertaken at least 6 times a day in persons with diabetes taking insulin [B].

SMBG meter values should be targeted to correspond to an HbA1c $\leq 7\%$ (≤ 53 mmol/mol), to align with the CGM TIR target of >70% between 4 and 10 mmol (70–180 mg/dL), and the strong correlation of CGM metrics with HbA1c reviewed earlier. For educational purposes, 3.9 mmol/L (70 mg/dL) has been intentionally rounded to 4.0 mmol/L; however, some centers may use 3.9 mmol/L as usual practice. Fasting target ranges glucoses of 4.0–8 mmol/L (70–144 mg/dL) are recommended to achieve this HbA1c target. SMBG glucose levels prior to bed of 4.0 – 8 mmol (70–144 mg/dL) are appropriate; however, caregivers may have more confidence in higher levels within the 4.0–10 mmol/L (70–180 mg/dL) range in certain scenarios; for example, if there has been preceding hypoglycemia, exercise, hypoglycemic unawareness.

Practical Considerations and Limitations

Each mode of measuring glycemia comes with its own limitations and practical considerations. HbA1c has long been the accepted benchmark for glycemia. However, it has several limitations, such as (i) for a given HbA1c, there is a wide range of mean glucose concentration values, and for any given mean glucose value there is a wide range of HbA1c values [65], (ii) racial differences [66], and (iii) unreliable measurement in certain conditions including anemia, hemoglobinopathies, and liver disease [58]. CGM may be impacted by various pharmaceutical agents [67] and can be intermittently inaccurate; however, overall accuracy in modern CGM technology is very high [68–70]. SMBG retains a place in measuring glycemia where this is the only mode available, but maintaining the >6 self-tests per day to guide management has long been recognized as a significant barrier to improving overall glycemia [71]. Overall, the strengths and limitations of each mode of measuring glycemia should be considered and interpreted in accordance to the environment and/or clinical characteristics of the person with diabetes.

In addition, while advanced technologies are enabling people with diabetes and healthcare professionals to aim for more optimal glucose levels, the demands of care and the limitations of current treatment options prevent this from being a reality for most young people with diabetes and their caregivers.

Conclusion

Future research is needed to evaluate differences in risk of complications and perceived care burdens between the currently proposed HbA1c target of 6.5% (48 mmol/mol) and the former 7.0% (53 mmol/mol) target. The former might be achieved using mathematical modeling to assess the relative risk of diabetes complications based on historical and contemporary data. These data would help inform the benefits of a lower HbA1c weighed against any increased burden of care and assist an individualized approach to setting glycemic targets.

As we move forward, there is also a need to utilize standardized CGM metrics alongside emerging diagnostic techniques that can predict diabetes complications. This approach will help bridge the gap until long-term registry data can confirm the relationship between CGM metrics and complication development.

Herein, we advocate for a shift in HbA1c targets that aligns with advancements in diabetes technologies and care. This approach is pragmatic to the immediate needs of individuals and their caregivers while embracing anticipatory and preventive care strategies for the long term. To achieve uniform glycemic targets across the ISPAD community, ongoing efforts are needed to address global disparities in access to these technologies.

Methodology

A literature search was conducted to gather updated evidence, using a combination of relevant medical subject headings (MeSH, Emtree) and free text terms specific to each chapter's focus. Studies published from 2021 to 2022 onward, related to children and young adults, were retrieved from MEDLINE. The Project Officer, in collaboration with chapter leads and co-authors, performed the literature searches. The resulting articles were then uploaded to COVIDENCE for screening and review. Two authors/experts involved in drafting this guideline version independently screened the articles. Any disagreements were resolved by a third reviewer. Where relevant, further literature was included.

Recommendations were graded as per the ADA evidence grading system for Standards of Medical Care in Diabetes [72]. This hierarchical A-E grading system sets A as having the highest level of evidence, and E having the lowest.

The draft chapter was posted on the ISPAD forum to allow feedback from the greater ISPAD membership. Modifications were made with authorship consensus, with the chapter receiving endorsement from the ISPAD editorial team. Literature search terms are summarized in online supplementary material (for all online suppl. material, see <https://doi.org/10.1159/000543266>).

Conflict of Interest Statement

M.B. received research funding from Novo Nordisk, Medtronic, Ypsomed, Dexcom, and Insulet; honoraria, travel expenses, or speaking fees from Novo Nordisk, Sanofi, Pfizer, Medtronic, Boehringer Ingelheim, Ypsomed, Dexcom, and Insulet; and advisory boards for Tandem and Dexcom, Nascence technology, and Tautoko Biomedical. K.D. received honoraria for participation in the speaker's bureau of Abbott, Eli Lilly, Medtronic, Novo Nordisk A/S, and Pfizer and advisory board for Medtronic and Novo Nordisk. D.M.M. received research support

from the NIH, JDRF, NSF, and the Helmsley Charitable Trust, and his institution has had research support from Medtronic, Dexcom, Insulet, Bigfoot Biomedical, Tandem, and Roche. D.M.M. has consulted for Abbott, Aditxt, the Helmsley Charitable Trust, Lifescan, Mannkind, Sanofi, Novo Nordisk, Eli Lilly, Medtronic, Insulet, Dompe, BioSpex, Provention Bio, Kriya, Enable Biosciences, and Bayer. F.H.M. received research support from the Canadian Institute for Health Research (CIHR), Physicians Services Incorporated, Heart and Stroke Foundation, Diabetes UK, and Breakthrough Diabetes (formerly JDRF). C.E.S. received speaker honoraria from Medtronic and Eli Lilly and advisory boards for Abbott. M.L.M. received research support from the NIH, JDRF, and the Helmsley Charitable Trust. Her institution has received research support from Dompe, Lilly, Mannkind, Medtronic, Provention Bio, Sanofi, and Zealand. L.A.D. has consulted for Abata, Tandem, Biomea Fusion, and Vertex. She also received payment from Sanofi for a CME talk (content independently developed by her). J.C.A., M.D., Y.N.-M., and L.P. have no conflicts of interest to declare.

Funding Sources

The 2024 Consensus guidelines were supported by unrestricted grants from Abbott Diabetes Care, Dexcom, Medtronic and Sanofi. These companies did not take part in any aspect of the development of these guidelines.

Author Contributions

M.B. and L.A.D. co-directed the guideline development process. M.B., J.C.A., M.D., K.D., D.M.M., M.L.M., F.M.M., Y.N.-M., L.P., C.E.S., and L.A.D. contributed equally to the content of individual chapters and participated in revising both the original manuscript and subsequent versions. M.B. synthesized these contributions to develop the original full draft of the guideline. L.A.D., as the editor of the guideline, supervised the overall development and revision process.

References

- White NH, Cleary PA, Dahms W, Goldstein D, Malone J, Tamborlane WV, et al. Beneficial effects of intensive therapy of diabetes during adolescence: outcomes after the conclusion of the Diabetes Control and Complications Trial (DCCT). *J Pediatr*. 2001; 139(6):804–12. <https://doi.org/10.1067/mpd.2001.118887>
- Diabetes Control and Complications Trial Research Group. Effect of intensive diabetes treatment on the development and progression of long-term complications in adolescents with insulin-dependent diabetes mellitus: diabetes control and complications trial. *J Pediatr*. 1994;125(2):177–88.
- Arnqvist HJ, Ludvigsson J, Nordwall M. Early increase in HbA1c trajectory predicts development of severe microangiopathy in patients with type 1 diabetes: the VISS study. *BMJ Open Diabetes Res Care*. 2024;12(3): e003917. <https://doi.org/10.1136/bmjdc-2023-003917>
- Arnqvist HJ, Westerlund MC, Fredrikson M, Ludvigsson J, Nordwall M. Impact of HbA1c followed 32 years from diagnosis of type 1 diabetes on development of severe retinopathy and nephropathy: the VISS study. *Diabetes Care*. 2022;45(11): 2675–82. <https://doi.org/10.2337/dc22-0239>
- Tienhaara E, Falck AAK, Pokka TML, Tossavainen PH. The natural history of emerging diabetic retinopathy and microalbuminuria from prepuberty to early adulthood in type 1 diabetes: a 19-year prospective clinical follow-up study. *Diabet Med*. 2022;39(1): e14732. <https://doi.org/10.1111/dme.14732>
- Miller RG, Orchard TJ, Costacou T. 30-year cardiovascular disease in type 1 diabetes: risk and risk factors differ by long-term patterns of glycemic control. *Diabetes Care*. 2022; 45(1):142–50. <https://doi.org/10.2337/dc21-1381>
- Miller RG, Orchard TJ. Understanding metabolic memory: a tale of two studies. *Diabetes*. 2020;69(3):291–9. <https://doi.org/10.2337/db19-0514>
- McEwan P, Bennett H, Bolin K, Evans M, Bergenheim K. Assessing the economic value of maintained improvements in type 1 diabetes management, in terms of HbA1c, weight and hypoglycaemic event incidence. *Diabet Med*. 2018;35(5):557–66. <https://doi.org/10.1111/dme.13590>
- Bak JC, Serné EH, Kramer MHH, Nieuwdorp M, Verheugt CL. National diabetes registries: do they make a difference? *Acta Diabetol*. 2021;58(3):267–78. <https://doi.org/10.1007/s00592-020-01576-8>
- Foster NC, Beck RW, Miller KM, Clements MA, Rickels MR, DiMeglio LA, et al. State of type 1 diabetes management and outcomes from the T1D exchange in 2016–2018. *Diabetes Technol Ther*. 2019;21(2):66–72. <https://doi.org/10.1089/dia.2018.0384>
- National Institute for Health and Care Excellence (NICE) (published 2015, updated 2023). Diabetes (type 1 and type 2) in children and young people: diagnosis and management. [cited August 21, 2024]; Available from: <https://www.nice.org.uk/guidance/ng18>
- Holmes-Walker DJ, Abraham MB, Chee M, Jones TW; Australasian Diabetes Data Network ADDN Study Group. Glycaemic outcomes in Australasian children and adults with type 1 diabetes: failure to meet targets across the age spectrum. *Intern Med J*. 2023; 53(1):61–7. <https://doi.org/10.1111/imj.15426>
- Al-Shehaili SM, Al-Johani SS, Al-Sarhan NT, Al-Anazi AA, Al-Mijmaj FF, Al-Qhatani WN, et al. The effect of poor glycemic control on cognitive function in children and adolescents with type 1 diabetes mellitus: a single-center cross-sectional study (2019–2020). *Saudi Med J*. 2023;44(10):1006–12. <https://doi.org/10.15537/smj.2023.44.20230327>
- Prahalad P, Riolo N, Noor N, Rapaport R, Weinstock RS, Ebekozien O, et al. T1D exchange quality improvement collaborative: accelerating change through benchmarking and improvement science for people with type 1 diabetes. *J Diabetes*. 2022;14(1):83–7. <https://doi.org/10.1111/1753-0407.13234>
- Gerhardsson P, Schwandt A, Witsch M, Kordonouri O, Svensson J, Forsander G, et al. The SWEET project 10-year benchmarking in 19 countries worldwide is associated with improved HbA1c and increased use of diabetes technology in youth with type 1 diabetes. *Diabetes Technol Ther*. 2021;23(7):491–9. <https://doi.org/10.1089/dia.2020.0618>
- Alonso GT, Corathers S, Shah A, Clements M, Kamboj M, Sonabend R, et al. Establishment of the T1D exchange quality improvement collaborative (T1DX-QI). *Clin Diabetes*. 2020;38(2):141–51. <https://doi.org/10.2337/cd19-0032>
- Samuelsson U, Åkesson K, Peterson A, Hanas R, Hanberger L. Continued improvement of metabolic control in Swedish pediatric diabetes care. *Pediatr Diabetes*. 2018;19(1):150–7. <https://doi.org/10.1111/pedi.12467>
- Addala A, Auzanneau M, Miller K, Maier W, Foster N, Kapellen T, et al. A decade of disparities in diabetes technology use and HbA1c in pediatric type 1 diabetes: a transatlantic comparison. *Diabetes Care*. 2021; 44(1):133–40. <https://doi.org/10.2337/dc20-0257>

- 19 Burnside MJ, Williman JA, Davies HM, Jefferies CA, Paul RG, Wheeler BJ, et al. Inequity in access to continuous glucose monitoring and health outcomes in paediatric diabetes, a case for national continuous glucose monitoring funding: a cross-sectional population study of children with type 1 diabetes in New Zealand. *Lancet Reg Health West Pac*. 2023;31:100644. <https://doi.org/10.1016/j.lanwpc.2022.100644>
- 20 Virmani A, Brink SJ, Middlehurst A, Mohsin F, Girauo F, Sarda A, et al. ISPAD Clinical Practice Consensus Guidelines 2022: management of the child, adolescent, and young adult with diabetes in limited resource settings. *Pediatr Diabetes*. 2022;23(8):1529–51. <https://doi.org/10.1111/pedi.13456>
- 21 Van Looke M, Battelino T, Tittel SR, Prahalad P, Goksen D, Davis E, et al. Lower HbA1c targets are associated with better metabolic control. *Eur J Pediatr*. 2021;180(5):1513–20. <https://doi.org/10.1007/s00431-020-03891-2>
- 22 de Bock M, Codner E, Craig ME, Huynh T, Maahs DM, Mahmud FH, et al. ISPAD Clinical Practice Consensus Guidelines 2022: glycemic targets and glucose monitoring for children, adolescents, and young people with diabetes. *Pediatr Diabetes*. 2022;23(8):1270–6. <https://doi.org/10.1111/pedi.13455>
- 23 Swedish National Diabetes Register (NDR). Diabetes in children and adolescents annual report 2020. https://www.ndr.nu/pdfs/Annual_Report_Swediabkids_2020.pdf
- 24 Sundberg F, deBeaufort C, Krogvold L, Patton S, Piloya T, Smart C, et al. ISPAD clinical practice consensus guidelines 2022: managing diabetes in preschoolers. *Pediatr Diabetes*. 2022;23(8):1496–511. <https://doi.org/10.1111/pedi.13427>
- 25 American Diabetes Association Professional Practice Committee. 14. Children and adolescents: standards of care in diabetes-2024. *Diabetes Care*. 2024;47(Suppl 1):S258–81. <https://doi.org/10.2337/dc24-S014>
- 26 Diabetes Control, and Complications Trial Research Group, Nathan DM, Genuth S, Lachin J, Cleary P, Crofford O, et al. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med*. 1993;329(14):977–86. <https://doi.org/10.1056/NEJM199309303291401>
- 27 Biester T, Berget C, Boughton C, Cudizio L, Ekhlaspour L, Hilliard ME, et al. ISPAD clinical practice consensus guidelines 2024: diabetes technologies: insulin delivery. *Horm Res Paediatr*. 2024;1–34. In preparation. <https://doi.org/10.1159/000543034>
- 28 Mathieu C, Ahmed W, Gillard P, Cohen O, Vigersky R, de Portu S, et al. The health economics of automated insulin delivery systems and the potential use of time in range in diabetes modeling: a narrative review. *Diabetes Technol Ther*. 2024;26(S3):66–75. <https://doi.org/10.1089/dia.2023.0438>
- 29 Biskupiak JE, Ramos M, Levy CJ, Forlenza G, Hopley C, Boyd J, et al. Cost-effectiveness of the tubeless automated insulin delivery system vs standard of care in the management of type 1 diabetes in the United States. *J Manag Care Spec Pharm*. 2023;29(7):807–17. <https://doi.org/10.18553/jmcp.2023.22331>
- 30 Hautala N, Hannula V, Palosaari T, Ebeling T, Falck A. Prevalence of diabetic retinopathy in young adults with type 1 diabetes since childhood: the Oulu cohort study of diabetic retinopathy. *Acta Ophthalmol*. 2014;92(8):749–52. <https://doi.org/10.1111/aos.12426>
- 31 Schreur V, van Asten F, Ng H, Weeda J, Groenewoud JMM, Tack CJ, et al. Risk factors for development and progression of diabetic retinopathy in Dutch patients with type 1 diabetes mellitus. *Acta Ophthalmol*. 2018;96(5):459–64. <https://doi.org/10.1111/aos.13815>
- 32 Sherif EM, Matter RM, Salah NY, Abozeid NEH, Atif HM, Tantawy NM. Changes in early optical coherence tomography angiography among children and adolescents with type 1 diabetes: relation to fibroblast growth factor 21. *Diabetes Metab Res Rev*. 2023;39(2):e3598. <https://doi.org/10.1002/dmrr.3598>
- 33 Samuelsson U, Anderzen J, Åkesson K, Hanberger L. The importance of low HbA1c during childhood on glycaemic control in adulthood and the risk of late complications. *Acta Paediatr*. 2021;110(4):1264–72. <https://doi.org/10.1111/apa.15591>
- 34 Spanakis EK, Golden SH. Race/ethnic difference in diabetes and diabetic complications. *Curr Diab Rep*. 2013;13(6):814–23. <https://doi.org/10.1007/s11892-013-0421-9>
- 35 Rossing P, Groop PH, Singh R, Lawatscheck R, Tuttle KR. Prevalence of chronic kidney disease in type 1 diabetes among adults in the US. *Diabetes Care*. 2024;47(8):1395–99.
- 36 Vazquez L, Vazquez Arreola E, Hanson RL, Sinha M. Glycemic measures in childhood as predictors of future diabetes-related microvascular complications in an indigenous american population. *Diabetes Care*. 2023;46(9):1659–67. <https://doi.org/10.2337/dc23-0343>
- 37 Rasmussen VF, Thrysoe M, Nyegaard JR, Tankisi H, Karlsson P, Hansen J, et al. Neuropathy in adolescents with type 1 diabetes: confirmatory diagnostic tests, bedside tests, and risk factors. *Diabetes Res Clin Pract*. 2023;201:110736. <https://doi.org/10.1016/j.diabres.2023.110736>
- 38 Ibrahim BA, Hussein SA, Abdullah WH. Cognitive functions in children with type 1 diabetes. *Wiad Lek*. 2023;76(5 pt 1):944–50. <https://doi.org/10.36740/WLek202305108>
- 39 Kar Ş, Er E, Ata A, İnal-Kaleli İ, Özcan T, Köse S, et al. Effect of metabolic control on cognitive functions in children and adolescents with type 1 diabetes mellitus. *J Pediatr Endocrinol Metab*. 2023;36(7):636–42. <https://doi.org/10.1515/jpem-2023-0027>
- 40 Mauras N, Buckingham B, White NH, Tsalikian E, Weinzimer SA, Jo B, et al. Impact of type 1 diabetes in the developing brain in children: a longitudinal study. *Diabetes Care*. 2021;44(4):983–92. <https://doi.org/10.2337/dc20-2125>
- 41 Liu S, Kuja-Halkola R, Larsson H, Lichtenstein P, Ludvigsson JF, Svensson AM, et al. Neurodevelopmental disorders, glycemic control, and diabetic complications in type 1 diabetes: a nationwide cohort study. *J Clin Endocrinol Metab*. 2021;106(11):e4459–70. <https://doi.org/10.1210/clinem/dgab467>
- 42 Papadakis JL, Anderson LM, Garza K, Feldman MA, Shapiro JB, Evans M, et al. Psychosocial aspects of diabetes technology use: the child and family perspective. *Endocrinol Metab Clin North Am*. 2020;49(1):127–41. <https://doi.org/10.1016/j.ecl.2019.10.004>
- 43 Lukács A, Mayer K, Sasvári P, Barkai L. Health-related quality of life of adolescents with type 1 diabetes in the context of resilience. *Pediatr Diabetes*. 2018;19(8):1481–6. <https://doi.org/10.1111/pedi.12769>
- 44 Chen CW, Tinsley LJ, Volkening LK, Anderson BJ, Laffel LM. Observed characteristics associated with diabetes device use among teens with type 1 diabetes. *J Diabetes Sci Technol*. 2023;17(1):186–94. <https://doi.org/10.1177/19322968211050069>
- 45 Rosner B, Roman-Urrestarazu A. Health-related quality of life in paediatric patients with type 1 diabetes mellitus using insulin infusion systems. A systematic review and meta-analysis. *PLoS One*. 2019;14(6):e0217655. <https://doi.org/10.1371/journal.pone.0217655>
- 46 Edqvist J, Rawshani A, Adiels M, Björck L, Lind M, Svensson AM, et al. BMI, mortality, and cardiovascular outcomes in type 1 diabetes: findings against an obesity paradox. *Diabetes Care*. 2019;42(7):1297–304. <https://doi.org/10.2337/dc18-1446>
- 47 DuBose SN, Hermann JM, Tamborlane WV, Beck RW, Dost A, DiMeglio LA, et al. Obesity in youth with type 1 diabetes in Germany, Austria, and the United States. *J Pediatr*. 2015;167(3):627–32.e324. <https://doi.org/10.1016/j.jpeds.2015.05.046>
- 48 Maffei C, Birkebaek NH, Konstantinova M, Schwandt A, Vazeou A, Casteels K, et al. Prevalence of underweight, overweight, and obesity in children and adolescents with type 1 diabetes: data from the international SWEET registry. *Pediatr Diabetes*. 2018;19(7):1211–20. <https://doi.org/10.1111/pedi.12730>
- 49 Birkebaek N, Kahlert J, Bjarnason R, Drivvoll AK, Johansen A, Konradsdottir E, et al. Body mass index standard deviation score and obesity in children with type 1 diabetes in the Nordic countries. HbA1c and other predictors of increasing BMISDS. *Pediatr Diabetes*. 2018;19(7):1198–205. <https://doi.org/10.1111/pedi.12693>

- 50 Nansel T, Lipsky L, Iannotti R. Cross-sectional and longitudinal relationships of body mass index with glycemic control in children and adolescents with type 1 diabetes mellitus. *Diabetes Res Clin Pract.* 2013; 100(1):126–32. <https://doi.org/10.1016/j.diabres.2012.12.025>
- 51 Minges KE, Whittemore R, Weinzimer SA, Irwin ML, Redeker NS, Grey M. Correlates of overweight and obesity in 5529 adolescents with type 1 diabetes: the T1D Exchange Clinic Registry. *Diabetes Res Clin Pract.* 2017; 126:68–78. <https://doi.org/10.1016/j.diabres.2017.01.012>
- 52 Phelan H, Foster NC, Schwandt A, Couper JJ, Willi S, Kroschwald P, et al. Longitudinal trajectories of BMI z-score: an international comparison of 11,513 Australian, American and German/Austrian/Luxembourgian youth with type 1 diabetes. *Pediatr Obes.* 2020;15(2):e12582. <https://doi.org/10.1111/ijpo.12582>
- 53 Grönberg A, Carlsson PO. Good glycemic control without exceeding the BMI trajectory during the first 5 years of treatment in children and adolescents with type 1 diabetes. *Pediatr Diabetes.* 2022;23(3):341–50. <https://doi.org/10.1111/peidi.13309>
- 54 Marlow AL, King BR, Phelan HT, Smart CE. Adolescents with type 1 diabetes can achieve glycemic targets on intensive insulin therapy without excessive weight gain. *Endocrinol Diabetes Metab.* 2022;5(4):e352. <https://doi.org/10.1002/edm2.352>
- 55 Seget S, Jarosz-Chobot P, Ochab A, Polanska J, Rusak E, Witoszek P, et al. Body mass index, basal insulin and glycemic control in children with type 1 diabetes treated with the advanced hybrid closed loop system remain stable-1-year prospective, observational, two-center study. *Front Endocrinol.* 2022;13: 1036808. <https://doi.org/10.3389/fendo.2022.1036808>
- 56 Piccini B, Felicioni M, Pessina B, Bertini M, Casalini E, Ceccotti C, et al. Glycemic control, basal/bolus distribution, BMI and meal management in children and adolescents with type 1 diabetes and advanced hybrid closed loop. *Nutrients.* 2023;15(23):4875. <https://doi.org/10.3390/nu15234875>
- 57 Beck RW, Bergenstal RM, Riddlesworth TD, Kollman C, Li Z, Brown AS, et al. Validation of time in range as an outcome measure for diabetes clinical trials. *Diabetes Care.* 2019; 42(3):400–5. <https://doi.org/10.2337/dc18-1444>
- 58 Chehregosha H, Khamseh ME, Malek M, Hosseinpahan F, Ismail-Beigi F. A view beyond HbA1c: role of continuous glucose monitoring. *Diabetes Ther.* 2019;10(3): 853–63. <https://doi.org/10.1007/s13300-019-0619-1>
- 59 Beck RW. Is it time to replace time-in-range with time-in-tight-range? Maybe not. *Diabetes Technol Ther.* 2024;26(3):147–50.
- 60 Petersson J, Åkesson K, Sundberg F, Särnblad S. Translating glycated hemoglobin A1c into time spent in glucose target range: a multicenter study. *Pediatr Diabetes.* 2019;20(3): 339–44. <https://doi.org/10.1111/peidi.12817>
- 61 Sundberg F, deBeaufort C, Krogvold L, Patton S, Piloya T, Smart C, et al. ISPAD clinical practice consensus guidelines 2022: managing diabetes in preschoolers. *Pediatr Diabetes.* 2022;23(8):1496–511. <https://doi.org/10.1111/peidi.13427>
- 62 Dunn TC, Ajjan RA, Bergenstal RM, Xu Y. Is it time to move beyond TIR to TITR? Real-world data from over 20,000 users of continuous glucose monitoring in patients with type 1 and type 2 diabetes. *Diabetes Technol Ther.* 2024;26(3):203–10. <https://doi.org/10.1089/dia.2023.0565>
- 63 Castañeda J, Arrieta A, van den Heuvel T, Battelino T, Cohen O. Time in tight glucose range in type 1 diabetes: predictive factors and achievable targets in real-world users of the MiniMed 780G system. *Diabetes Care.* 2024;47(5):790–7. <https://doi.org/10.2337/dc23-1581>
- 64 Hamidi V, Pettus JH. Time in tight range for patients with type 1 diabetes: the time is now, or is it too soon? *Diabetes Care.* 2024;47(5): 782–4. <https://doi.org/10.2337/dci23-0092>
- 65 Beck RW, Connor CG, Mullen DM, Wesley DM, Bergenstal RM. The fallacy of average: how using HbA1c alone to assess glycemic control can be misleading. *Diabetes Care.* 2017;40(8):994–9. <https://doi.org/10.2337/dc17-0636>
- 66 Bergenstal RM, Gal RL, Connor CG, Gubitosi-Klug R, Kruger D, Olson BA, et al. Racial differences in the relationship of glucose concentrations and hemoglobin A1c levels. *Ann Intern Med.* 2017;167(2):95–102. <https://doi.org/10.7326/M16-2596>
- 67 Basu A, Slama MQ, Nicholson WT, Langman L, Peyser T, Carter R, et al. Continuous glucose monitor interference with commonly prescribed medications: a pilot study. *J Diabetes Sci Technol.* 2017;11(5):936–41. <https://doi.org/10.1177/1932296817697329>
- 68 Almurashi AM, Rodriguez E, Garg SK. Emerging diabetes technologies: continuous glucose monitors/artificial pancreases. *J Indian Inst Sci.* 2023;103(1):205–30. <https://doi.org/10.1007/s41745-022-00348-3>
- 69 Garg SK, Kipnes M, Castorino K, Bailey TS, Akturk HK, Welsh JB, et al. Accuracy and safety of Dexcom G7 continuous glucose monitoring in adults with diabetes. *Diabetes Technol Ther.* 2022;24(6):373–80. <https://doi.org/10.1089/dia.2022.0011>
- 70 Nguyen A, White JR. FreeStyle libre 3. *Clin Diabetes.* 2022;41(1):127–8. <https://doi.org/10.2337/cd22-0102>
- 71 Patton SR. Adherence to glycemic monitoring in diabetes. *J Diabetes Sci Technol.* 2015; 9(3):668–75. <https://doi.org/10.1177/1932296814567709>
- 72 American Diabetes Association. 16. Diabetes advocacy: Standards of medical care in diabetes-2019. *Diabetes Care.* 2019;42(Suppl 1):S182–3. <https://doi.org/10.2337/dc19-S016>