



Ensemble Wind Forecasting Based on the HARMONIE Model and Adaptive Finite Elements in Complex Orography

Albert Oliver (1), Eduardo Rodríguez (1), José María Escobar (1), Gustavo Montero (1), Mariano Hortal (2), Javier Calvo (2), José Manuel Cascón (3), and Rafael Montenegro (1)

(1) University Institute for Intelligent Systems and Numerical Applications in Engineering (SIANI), University of Las Palmas de Gran Canaria (ULPGC) Campus de Tafira, 35017 Las Palmas de Gran Canaria, Spain, (2) Agencia Estatal de Meteorología (AEMET), Leonardo Prieto Castro, 8, 28040 Madrid, Spain, (3) Department of Economics and Economic History, Faculty of Economics and Business, University of Salamanca, 37007 Salamanca, Spain

Ensemble forecasting [1] is a methodology to deal with uncertainties in the numerical wind prediction. In this work we propose to apply ensemble methods to the adaptive wind forecasting model presented in [2].

The wind field forecasting is based on a mass-consistent model and a log-linear wind profile using as input data the resulting forecast wind from Harmonie [3], a Non-Hydrostatic Dynamic model used experimentally at AEMET with promising results[4]. The mass-consistent model parameters are estimated by using genetic algorithms [5]. The mesh is generated using the meccano method [6] and adapted to the geometry.

The main source of uncertainties in this model is the parameter estimation. For this reason, the model parameters are perturbed to achieve a more reliable wind forecast.

This work presents the results of the model applied in Gran Canaria island using Harmonie data for days from 20 to 23 of February 2010, and its comparison with some experimental data obtained by the AEMET in their meteorological stations.

REFERENCES

- [1] M. Leutbecher, T.N. Palmer. Ensemble forecasting. *Journal of Computational Physics*, Vol. 227, 3515–3539, 2008. <http://dx.doi.org/10.1016/j.jcp.2007.02.014>
- [2] A. Oliver, E. Rodríguez, G. Montero, M. Hortal, J. Calvo, R. Montenegro. Wind forecasting over complex terrain. *Mathematics and Geosciences: Global and Local Perspectives*, ICMAT, Campus de Cantoblanco, Madrid (Spain) November 4th-8th, 2013
- [3] J. Simarro, M. Hortal. A semi-implicit non-hydrostatic dynamical kernel using finite elements in the vertical discretization. *Q.J.R. Meteorol. Soc.*, Vol. 138, 826–839, 2012. <http://dx.doi.org/10.1002/qj.952>
- [4] Navascués B, Calvo J, Morales G, Santos C, Callado A, Cansado A, Cuxart J, Díez M, del Río P, Escribà P, García-Colombo O, García-Moya J, Geijo C, Gutiérrez E, Hortal M, Martínez I, Orfila B, Parodi J, Rodríguez E, Sánchez-Arriola J, Santos-Atienza I, Simarro J (2013) Long-term verification of HIRLAM and ECMWF forecasts over southern europe: History and perspectives of numerical weather prediction at AEMET. *Atmospheric Research* 125-126(0):20–33, <http://dx.doi.org/10.1016/j.atmosres.2013.01.010>
- [5] G. Montero, E. Rodríguez, R. Montenegro, J.M. Escobar, J.M. González-Yuste. Genetic algorithms for an improved parameter estimation with local refinement of tetrahedral meshes in a wind model. *Advances in Engineering Software*, Vol. 36, 3–10, 2005. <http://dx.doi.org/10.1016/j.advengsoft.2004.03.011>
- [6] J.M. Cascón, E. Rodríguez, J.M. Escobar, R. Montenegro. Comparison of the meccano method with standard mesh generation techniques. *Engineering with Computers*, 1–14, 2013. <http://dx.doi.org/10.1007/s00366-013-0338-6>