



## RESEARCH ARTICLE

10.1002/2015GB005326

## Key Points:

- Volumetric N<sub>2</sub> fixation rates in aphotic waters were as high as previous euphotic rates
- N<sub>2</sub> fixation was higher in more ventilated and oxygenated waters containing fresher organic matter
- Deep N<sub>2</sub> fixation contributed largely to depth-integrated rates and poorly to total nitrogen inputs

## Supporting Information:

- Table S1

## Correspondence to:

M. Benavides,  
mar.benavides@ird.fr

## Citation:

Benavides, M., et al. (2016), Basin-wide N<sub>2</sub> fixation in the deep waters of the Mediterranean Sea, *Global Biogeochem. Cycles*, 30, 952–961, doi:10.1002/2015GB005326.

Received 9 NOV 2015

Accepted 16 JUN 2016

Accepted article online 20 JUN 2016

Published online 30 JUN 2016

## Basin-wide N<sub>2</sub> fixation in the deep waters of the Mediterranean Sea

Mar Benavides<sup>1,2</sup>, Sophie Bonnet<sup>1</sup>, Nautzet Hernández<sup>2</sup>, Alba María Martínez-Pérez<sup>3</sup>, Mar Nieto-Cid<sup>3</sup>, Xosé Antón Álvarez-Salgado<sup>3</sup>, Isabel Baños<sup>2</sup>, María F. Montero<sup>2</sup>, Ignacio P. Mazuecos<sup>4</sup>, Josep M. Gasol<sup>5</sup>, Helena Osterholz<sup>6</sup>, Thorsten Dittmar<sup>6</sup>, Ilana Berman-Frank<sup>7</sup>, and Javier Arístegui<sup>2</sup>


<sup>1</sup>Mediterranean Institute of Oceanography-IRD/CNRS/Aix-Marseille University, Noumea, New Caledonia, <sup>2</sup>Instituto de Oceanografía y Cambio Global, Universidad de Las Palmas de Gran Canaria, Las Palmas de Gran Canaria, Spain, <sup>3</sup>Instituto de Investigaciones Mariñas, CSIC, Vigo, Spain, <sup>4</sup>Departamento de Ecología, Instituto del Agua, Universidad de Granada, Granada, Spain, <sup>5</sup>Institut de Ciències del Mar, CSIC, Barcelona, Spain, <sup>6</sup>Research Group for Marine Geochemistry, Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, Oldenburg, Germany, <sup>7</sup>Mina and Everard Goodman Faculty of Life Sciences, Bar-Ilan University, Ramat Gan, Israel

**Abstract** Recent findings indicate that N<sub>2</sub> fixation is significant in aphotic waters, presumably due to heterotrophic diazotrophs depending on organic matter for their nutrition. However, the relationship between organic matter and heterotrophic N<sub>2</sub> fixation remains unknown. Here we explore N<sub>2</sub> fixation in the deep chlorophyll maximum and underneath deep waters across the whole Mediterranean Sea and relate it to organic matter composition, characterized by optical and molecular methods. Our N<sub>2</sub> fixation rates were in the range of those previously reported for the euphotic zone of the Mediterranean Sea (up to 0.43 nmol N L<sup>-1</sup> d<sup>-1</sup>) and were significantly correlated to the presence of relatively labile organic matter with fluorescence and molecular formula properties representative for peptides and unsaturated aliphatics and associated with the presence of more oxygenated ventilated water masses. Finally, and despite that the aphotic N<sub>2</sub> fixation contributes largely to total water column diazotrophic activity (>50%), its contribution to overall nitrogen inputs to the basin is negligible (<0.5%).

### 1. Introduction

The Mediterranean Sea is characterized by high nitrogen-to-phosphorus (N:P) ratios in deep water masses, which increase from 20 to 22:1 in the western subbasin to 24–28:1 in the eastern subbasin [Krom and Herut, 2004; Pujo-Pay et al., 2011]. These ratios are well above the canonical Redfield ratio (16:1) and have been claimed to be caused either by external nutrient inputs with high N:P values [Krom and Herut, 2004; Ludwig et al., 2009; Markaki et al., 2010; Pujo-Pay et al., 2011; Rodellas et al., 2015] or alternatively by dinitrogen (N<sub>2</sub>) fixation (the reduction of N<sub>2</sub> to ammonia by prokaryotic microbes named diazotrophs) [Sachs, 1999]. While early geochemical balance and isotope fractionation studies suggested a high contribution of N<sub>2</sub> fixation to new production in the Mediterranean Sea [Bethoux and Copin-Montegut, 1986; Sachs, 1999; Pantoja et al., 2002], direct studies that measured biological N<sub>2</sub> fixation rates through <sup>15</sup>N<sub>2</sub> tracer experiments in sunlit waters found that diazotrophic activity in this basin is very low (typically <0.5 nmol N L<sup>-1</sup> d<sup>-1</sup>) [Jbello et al., 2010; Bonnet et al., 2011; Yogev et al., 2011; Rahav et al., 2013a, 2013b]. These low rates could partly be due to methodological underestimations and to the fact that the majority of studies were performed during the summer, which is the most oligotrophic season (as discussed in Rahav et al. [2013a]), but still the contribution of N<sub>2</sub> fixation to new production is thought to be low [Bonnet et al., 2011; Rahav et al., 2013a].

Diazotrophs include cyanobacteria, other bacteria, and archaea [Zehr et al., 2003]. In the sunlit waters of the Mediterranean Sea, unicellular cyanobacteria (UCYN) and filamentous cyanobacteria in symbiosis with diatoms (diatom-diazotroph associations, DDAs) have been reported as the most abundant diazotrophs [Bar-Zeev et al., 2008; Le Moal et al., 2011]. Recent investigations have, however, shown that heterotrophic non-cyanobacterial diazotrophs, which are thought to outnumber autotrophic cyanobacterial diazotrophs in the world's oceans [Farnelid et al., 2011], are also present in the eastern Mediterranean Sea [Man-Aharonovich et al., 2007; Yogev et al., 2011; Rahav et al., 2013c]. Some heterotrophic diazotrophs are not constrained by the availability of light and are therefore able to live below the euphotic layer. Rahav et al. [2013c] measured significant N<sub>2</sub> fixation rates in the aphotic waters of the Levantine Basin at depths extending from the base of the euphotic zone down to 720 m (up to 0.38 nmol N L<sup>-1</sup> d<sup>-1</sup>) and related this



**Figure 1.** Stations sampled during the HOTMIX cruise superimposed to a satellite image of chlorophyll *a* (Chl *a*) concentrations ( $\text{mg m}^{-3}$ ) obtained from the National Aeronautics and Space Administration (NASA) Goddard Earth Sciences Data and Information Services Center Giovanni (NASA GES DISC) online database for the month of May 2014. Stations sampled for nutrients and DOM are depicted by crosses, while stations where  $\text{N}_2$  fixation rates were measured are depicted by open circles.

fixation to sinking organic material such as transparent exopolymeric particles (TEPs) or other aggregates providing carbon resources for heterotrophic diazotrophs. These authors also reported an enhancement of  $\text{N}_2$  fixation rates upon the addition of TEP and of organic substrates such as amino acids. Similar bioassays have been conducted in the aphotic waters off coastal Peru [Bonnet *et al.*, 2013] and in sunlit waters in coastal Denmark [Severin *et al.*, 2015] and suggest that heterotrophic  $\text{N}_2$  fixation is regulated by organic matter availability, but the relationship between the diversity and activity of heterotrophic diazotrophs with the quantity and composition of organic matter has not been studied in detail [i.e., Riemann *et al.*, 2010]. Moreover, the relative contribution of aphotic  $\text{N}_2$  fixation to whole water column diazotrophic activity at the scale of the Mediterranean Sea basin has never been quantified. In the light of the high N:P ratios observed in deep waters of the Mediterranean Sea, we explore here the magnitude of aphotic  $\text{N}_2$  fixation rates and its contribution to total nitrogen inputs to the basin. Furthermore, we investigate the relationship between  $\text{N}_2$  fixation at the DCM and in deep waters to the quality and quantity of the in situ organic matter pools.

## 2. Methods

### 2.1. Sampling Strategy and Hydrographic Data

The HOTMIX cruise took place between 27 April and 29 May 2014 aboard the R/V *Sarmiento de Gamboa* across the Mediterranean Sea (Figure 1). Seawater was collected with a General Oceanics rosette, equipped with 24 L Niskin bottles. Hydrographic data (temperature, conductivity, depth (CTD)) were recorded with a SBE911+ CTD equipped with fluorescence and oxygen sensors.

Seawater samples were collected from four levels: the deep chlorophyll maximum (DCM, 1% photosynthetically active radiation, ranging between 60 and 130 m along the cruise), the Levantine Intermediate Water (LIW, located at 200–300 m in the eastern basin and at 400–500 m in the western basin), the oxygen minimum (located at 900–1000 m), and the Eastern and Western Mediterranean Deep Water (EMDW and WMDW, located at >1500 m). Two more stations were sampled at the exit of the Strait of Gibraltar (25 and 27; Figure 1), where we included sampling of the Mediterranean Water (MW), which is a mixture of LIW and North Atlantic Central Water (NACW).

Oxygen solubility was calculated using the equations outlined in *United Nations Educational, Scientific and Cultural Organization* [1986], and apparent oxygen utilization (AOU,  $\mu\text{mol O}_2 \text{ kg}^{-1}$ ) was computed by subtracting dissolved oxygen concentrations from oxygen solubility at saturation with respect to the atmosphere.

### 2.2. $\text{N}_2$ Fixation Rates

Samples for  $\text{N}_2$  fixation measurements were collected at stations 1, 4, 7, 10, 15, 18, 21, 25, and 27 (Figure 1). The samples were taken in duplicate 4.3 L polycarbonate bottles (Nalgene) fitted with teflon-lined septa and darkened with black fabric to reproduce aphotic conditions. To assay  $\text{N}_2$  fixation, 5 mL of high-purity  $^{15}\text{N}_2$  gas

(Cambridge Isotopes, 98.9% atom  $^{15}\text{N}$ ) were injected to each bottle. The bottles were transferred to a temperature-controlled room set at 13°C. Samples taken from the DCM were incubated in on-deck incubators connected to the underway system seawater and darkened with layers of mesh gauze to simulate in situ light levels. The temperature of the on-deck incubators ranged between ~14 and 19°C along the cruise. Incubations lasted 24 h.

The samples were filtered through precombusted (450°C, 6 h) GF/F filters (Whatman) and stored at –20°C until being analyzed with an Integra2 Analyzer, calibrated every 10 samples using reference material (IAEA-N1).  $\text{N}_2$  fixation rates were calculated with the equations of *Montoya et al.* [1996]. At every station and sampling depth an extra 4.3 L bottle was collected and immediately filtered to determine the background atom %  $^{15}\text{N}$  in the particulate nitrogen (PN). Considering the PN linearity limit of the mass spectrometer (2.32  $\mu\text{g N}$ ), 3 times the standard deviation of our background (time zero atom %  $^{15}\text{N}$ ) values, our usual filtration volume, and incubation time, our volumetric  $\text{N}_2$  fixation rate detection limit was 0.031  $\text{nmol NL}^{-1} \text{d}^{-1}$ .

Our rates could be somewhat underestimated due to incomplete dissolution of  $^{15}\text{N}_2$  gas [*Großkopf et al.*, 2012]. The alternative dissolved  $^{15}\text{N}_2$  method is prone to trace metal contaminations [*Klawonn et al.*, 2015] and thus was not used here. Moreover, the potential underestimation of the  $^{15}\text{N}_2$  bubble method is minimized in long (24 h) incubations [*Mulholland et al.*, 2012], and the contamination of  $^{15}\text{N}_2$  gas stocks has been shown to be minimal for the brand used in this study [*Dabundo et al.*, 2014].

### 2.3. Heterotrophic Prokaryotic Abundance and Production

Prokaryotic abundance was determined by flow cytometry using a FACSCalibur instrument (BD Biosciences) [*Gasol and del Giorgio*, 2000]. Prokaryotes were stained with SYBR Green I (Molecular Probes, Invitrogen) and detected in side scatter against green fluorescence plots. Yellow green Polysciences 1  $\mu\text{m}$  beads were used as a reference.

Prokaryotic heterotrophic production was assayed by  $^3\text{H}$ -Leucine (specific activity = 112  $\text{Ci mmol}^{-1}$ ) assimilation into proteins [*Kirchman et al.*, 1985], using the microcentrifugation procedure [*Smith and Azam*, 1992]. Leucine was added at a 20 nM concentration. Four replicates and two trichloroacetic acid fixed controls were done per sample.

### 2.4. Microbial Electron Transport System Respiratory Activity

Electron transport system (ETS) determinations were performed on 3–8 L samples filtered onto 47 mm GF/F filters as described in *Aristegui and Montero* [1995].

### 2.5. Inorganic Nutrients

Samples for the analysis of nitrate and phosphate concentrations were collected in 50 mL polyethylene tubes and determined immediately on board using a Skalar autoanalyzer SAN++ following the colorimetric methods of *Grasshoff et al.* [1999].


### 2.6. Particulate Organic Matter

For the quantification of particulate organic carbon (POC) and nitrogen (PON) concentrations 2–4 L seawater samples were filtered onto precombusted GF/F filters and analyzed on a Perkin Elmer 2400 CHN. TEP concentrations were measured using the Alcian blue method [*Passow and Alldredge*, 1995] on triplicate samples (0.4–2 L) filtered through 0.4  $\mu\text{m}$  polycarbonate filters (Poretics).

### 2.7. Dissolved Organic Matter

Dissolved organic matter (DOM) samples were collected in acid-clean 5 L polycarbonate carboys and filtered through precombusted GF/F filters under positive pressure with high-purity  $\text{N}_2$  in an all-glass filtration system. The filtrates were collected in acid-clean Teflon bottles for solid phase extraction prior to molecular-level analysis. Approximately 10 mL of the filtrate were collected in precombusted glass ampoules for dissolved organic carbon (DOC) analysis with a Shimadzu TOC-V analyzer. The precision of the instrument was tested on a daily basis with DOC reference materials (provided by D.A. Hansell, University of Miami, USA).

The absorption spectra of the colored DOM (CDOM) were determined between 250 and 750 nm at 1 nm intervals in 10 cm path length quartz cuvettes using a Perkin Elmer lambda 850 spectrophotometer. The absorbance was converted to Neperian absorption coefficient (per meter) according to *Green and Blough*



**Figure 2.** (a)  $N_2$  fixation rates (sized grey circles, a size reference corresponding to a rate of  $0.5 \text{ nmol N L}^{-1} \text{ d}^{-1}$  is displayed at the lower left corner), superimposed on dissolved oxygen ( $\mu\text{mol kg}^{-1}$ ; color scale) and salinity (isolines) data. Red asterisks indicate sites where  $N_2$  fixation activity was below the detection limit ( $<0.031 \text{ nmol N L}^{-1} \text{ d}^{-1}$ ), numbers over the panel indicate the number of the station as depicted in Figure 1, (b) nitrate to phosphate (N:P) ratio, (c) nitrate concentrations ( $\mu\text{M}$ ), (d) phosphate concentrations ( $\mu\text{M}$ ), (e) Chl *a* concentrations ( $\text{mg m}^{-3}$ ), and (f) POC concentrations ( $\mu\text{M}$ ).

[1994]. The carbon-specific absorption coefficient at 254 nm ( $a_{\text{CDOM}}(254)/\text{DOC}$ ) was used for Spearman correlations. The fluorescence of CDOM (FDOM) was recorded with a Perkin Elmer LS55 luminescence spectrofluorometer. Excitation and emission slit widths were 10 nm. Single-point measurements were performed at the classical peaks defined by Coble [1996] representative of humic- (peak C: 340/440 nm) and protein-like (peak T: 280/350 nm) DOM. Milli-Q water was used as a reference blank, and the analyses were carried out at room temperature in a 1 cm quartz fluorescence cell. Carbon-specific FDOM of peaks C and T were used for Spearman correlations, along with the ratio peak C/peak T.

Samples for ultrahigh-resolution mass spectrometry analyses were extracted (solid phase) with Agilent PPL cartridges as described in Dittmar *et al.* [2008]. After extraction, the cartridges were rinsed with acidified ultrapure water and frozen at  $-20^\circ\text{C}$ . Subsequently, they were dried by flushing with high-purity  $N_2$  and eluted with 6 mL of methanol. The efficiency of the extraction was  $47.3 \pm 3.9\%$  on a carbon basis. Methanol extracts were molecularly characterized with a 15 Tesla Fourier transform ion cyclotron resonance mass spectrometer (Solarix FT-ICR-MS, Bruker Daltonik GmbH) using an electrospray ionization source in negative mode (Bruker Apollo II). Molecular formulae were ascribed to the detected masses as outlined in Seidel *et al.* [2014]. The aromaticity and unsaturation degree of each compound were evaluated according to its molecular formula and were presented as the modified aromaticity index (AI-mod) and double bond equivalents (DBE), respectively

**Table 1.** Depth-Integrated Aphotic N<sub>2</sub> Fixation Rates and Their Contribution to Total Water Column Activity and to Total Nitrogen Inputs to Each Subbasin in the Mediterranean Sea

Subbasin	Aphotic Integrated N <sub>2</sub> Fixation (μmol N m <sup>-2</sup> d <sup>-1</sup> ) <sup>a</sup>		Photic (Surface to 0.1% PAR) Integrated N <sub>2</sub> Fixation (μmol N m <sup>-2</sup> d <sup>-1</sup> ) <sup>b</sup>		Total Water Column Integrated N <sub>2</sub> Fixation (μmol N m <sup>-2</sup> d <sup>-1</sup> )		Percent Contribution of Aphotic N <sub>2</sub> Fixation		Volume Below the DCM (km <sup>3</sup> )	N Inputs From Aphotic N <sub>2</sub> Fixation (× 10 <sup>9</sup> mol yr <sup>-1</sup> ) <sup>a</sup>	Total N Inputs per Subbasin (× 10 <sup>9</sup> mol yr <sup>-1</sup> ) <sup>b</sup>	Percent Contribution of Aphotic N <sub>2</sub> Fixation to Total N Inputs
	Eastern (n = 16)	Western (n = 15)	Eastern (n = 18)	Western (n = 24)	Eastern (n = 81) <sup>c</sup>	Western (n = 72) <sup>c</sup>	Eastern (n = 96)	Western (n = 94)				
Eastern	17.83–91.06	1.0–4.0	0–0.4	0–0.4	17.83–95.06	96–100	0.35–0.70	181 <sup>e</sup> /293 <sup>f</sup>	1.98 × 10 <sup>6</sup>	0.39 <sup>d</sup> /0.24 <sup>e</sup>	0.40–0.48 <sup>d</sup> /0.28–0.31 <sup>e</sup>	
Western	37.95–68.96	2.5–3.7	10–76	10–76	40.45–144.96	48–94	0.23–0.38	78–94 <sup>e</sup> /126–142 <sup>f</sup>	1.31 × 10 <sup>6</sup>	0.40–0.48 <sup>d</sup> /0.28–0.31 <sup>e</sup>		

<sup>a</sup>This study.

<sup>b</sup>Ibello *et al.* [2010].

<sup>c</sup>Bonnet *et al.* [2011].

<sup>d</sup>Considering 6 months of activity per year.

<sup>e</sup>Considering a conservative submarine groundwater discharge (SGD) (only freshwater inputs) [Rodellas *et al.*, 2015].

<sup>f</sup>Considering the median SGD for the whole Mediterranean Sea [Rodellas *et al.*, 2015].

[Koch and Dittmar, 2006]. In addition, we ascribed the identified molecular formulae identified to compound groups according to established molar ratios, AI-mod, DBE, and heteroatom contents [Seidel *et al.*, 2014] (see Table S1 in the supporting information). A detailed discussion of molecular DOM quality will be given elsewhere (Martínez-Pérez *et al.*, manuscript in preparation).

### 3. Results and Discussion

#### 3.1. Does Aphotic N<sub>2</sub> Fixation Contribute Significantly to Total Nitrogen Inputs in the Mediterranean Sea?

Deep N<sub>2</sub> fixation rates during this study ranged from 0.13 to 0.43 nmol N L<sup>-1</sup> d<sup>-1</sup> at the DCM and from undetectable (<0.031 nmol N L<sup>-1</sup> d<sup>-1</sup>) to 0.07 nmol N L<sup>-1</sup> d<sup>-1</sup>, in the mesopelagic and bathypelagic waters below (Figure 2a). The highest rates were observed thus at the DCM, and these were higher in the western as compared to the eastern basin (0.31–0.43 and 0.17–0.24 nmol N L<sup>-1</sup> d<sup>-1</sup>, respectively), which is in agreement with previous west-east decreasing gradients observed for euphotic N<sub>2</sub> fixation in the Mediterranean Sea [Bonnet *et al.*, 2011; Rahav *et al.*, 2013b]. Conversely, slightly higher rates were observed at the LIW level in the eastern subbasin (~0.06 nmol N L<sup>-1</sup> d<sup>-1</sup>) as compared to the western subbasin (0.03–0.05 nmol N L<sup>-1</sup> d<sup>-1</sup>). At the oxygen minimum level (900–1250 m), N<sub>2</sub> fixation was undetectable at stations 4, 7, and 10 (eastern subbasin) and station 15 (western subbasin). At stations where N<sub>2</sub> fixation at the oxygen minimum was detectable, the rates were similar (0.03–0.05 nmol N L<sup>-1</sup> d<sup>-1</sup>) to those detected at the level of the MDW at stations 1, 7 (eastern subbasin), and 21 (western subbasin). In the northeast Atlantic N<sub>2</sub> fixation was only detected at the DCM and at the oxygen minimum level at station 27 (Figure 2a). The eastward decrease in N<sub>2</sub> fixation activity coincided with an increase in the N:P ratios of surface waters (Figure 2b), a general decrease in nitrate and phosphate concentrations (Figures 2c and 2d), and the deepening of the DCM (Figure 2e), with the surface concentrations of POC being higher in the western basin.

Our aphotic N<sub>2</sub> fixation rates are in the same range as those reported previously from the surface Mediterranean Sea [Ibello *et al.*, 2010; Bonnet *et al.*, 2011; Yogeve *et al.*, 2011; Rahav *et al.*, 2013b], and they are also comparable with previous results from the mesopelagic to bathypelagic waters of the Southern California Bight (0.07 nmol N L<sup>-1</sup> d<sup>-1</sup>) [Hamersley *et al.*, 2011], the Levantine Basin and the Gulf of Aqaba (0.01–0.38 nmol N L<sup>-1</sup> d<sup>-1</sup>) [Rahav *et al.*, 2013c], and the eastern tropical South Pacific (<1 nmol N L<sup>-1</sup> d<sup>-1</sup>) [Bonnet *et al.*, 2013]. This fact highlights the relevance of aphotic N<sub>2</sub> fixation, which has been largely disregarded in past studies. The resulting contribution of aphotic N<sub>2</sub> fixation to total water column activity (using previously published data as a reference) is 48–94% and 96–100% in the western and eastern subbasins, respectively (Table 1). This high contribution of the aphotic layer to total water column N<sub>2</sub> fixation is comparable to the values reported

from the South Pacific (80–90%) [Bonnet *et al.*, 2013; Benavides *et al.*, 2015] and the eastern Mediterranean Sea (up to 75%) [Rahav *et al.*, 2013c] and could potentially alleviate nitrogen limitation of aphotic bacterial communities [Rahav *et al.*, 2013c]. It is currently not known whether aphotic N<sub>2</sub> fixation occurs homogeneously throughout the water column or if it takes place in hot spots of microbial activity such as particles or ecotones [Riemann *et al.*, 2010], which gives whole water column integrated rates some uncertainty. Our sampling strategy does not allow us to solve this question.

Earlier geochemical-oriented studies suggested that N<sub>2</sub> fixation provided a significant input of nitrogen to the Mediterranean Sea, accounting for the high N:P ratios observed in deep waters [Sachs, 1999; Pantoja *et al.*, 2002]. Considering the depth-integrated N<sub>2</sub> fixation rates measured in our study (17.83–95.06 and 40.45–144.96  $\mu\text{mol N m}^{-2} \text{d}^{-1}$  for the eastern and western subbasins, respectively; Table 1), a volume of  $1.39 \times 10^6$  and  $2.15 \times 10^6 \text{ km}^3$  for the western and eastern subbasins, respectively [Manca *et al.*, 2004], and an average DCM depth of 100 m [Macías *et al.*, 2014], we estimate yearly total nitrogen inputs associated with aphotic N<sub>2</sub> fixation of 0.35–0.70 and  $0.23\text{--}0.38 \times 10^9 \text{ mol N yr}^{-1}$  for the eastern and western subbasins, respectively (considering a conservative diazotrophic activity during 6 months/yr; Table 1). Taking into account all other inputs of nitrogen to the Mediterranean Sea (atmospheric, riverine, and submarine groundwater discharge (SGW)), aphotic N<sub>2</sub> fixation would contribute for <0.5% of total nitrogen inputs in the basin (Table 1). The atmospheric, riverine, and SGW sources of nitrogen feature high N:P ratios (65–102, 36–47, and 80–430, respectively) [Huertas *et al.*, 2012; Rodellas *et al.*, 2015], well above those of the deep Mediterranean Sea waters (28.84 and 25.73 for the east and west subbasins, respectively—as calculated using nutrient data from our cruise). Hence, these high N:P ratios are the main factors responsible for the recurrently measured phosphorus limitation [Krom and Herut, 2004], and thus, we infer that the contribution of aphotic N<sub>2</sub> fixation to the high N:P ratios of the deep waters of the Mediterranean Sea is negligible.

### 3.2. Nature of Aphotic N<sub>2</sub> Fixation in the Mediterranean Sea

Previous molecular analyses identified UCYN and DDAs as recurrent diazotrophs in the sunlit waters of the Mediterranean Sea [Bar-Zeev *et al.*, 2008; Le Moal *et al.*, 2011]. At the global scale non-cyanobacterial *nifH* genes outnumber cyanobacterial ones [Farnelid *et al.*, 2011], and recent studies suggest that non-cyanobacterial diazotrophs may also dominate N<sub>2</sub> fixation activity at the surface of the Mediterranean Sea [Man-Aharonovich *et al.*, 2007; Yogevev *et al.*, 2011; Rahav *et al.*, 2013a, 2015]. Non-cyanobacterial diazotrophs inhabiting aphotic waters are presumably heterotrophic and rely on organic matter for their nutrition [Riemann *et al.*, 2010]. Absolute organic matter inputs to the aphotic zone accounting for deep prokaryotic activity include the transport of DOM from the surface to deep waters via water mass sinking processes, the settling of organic particles, the release of DOM by migrating zooplankton, and the in situ production of DOM by chemolithoautotrophic bacteria [Aristegui *et al.*, 2009]. In the Mediterranean Sea DOM inputs to aphotic waters are quantitatively important given the high frequency and various sites of deep water formation [Santinelli *et al.*, 2010]. Together with the high temperatures observed in aphotic waters (>12.5°C), these peculiar conditions are thought to promote prokaryotic activity at depth [Santinelli *et al.*, 2010; Boutrif *et al.*, 2011; Luna *et al.*, 2012].

Detailed descriptions of the variability of nutrients, DOM, prokaryotic heterotrophic production, and diversity parameters during this cruise can be found elsewhere (Martínez-Pérez *et al.*; Gasol *et al.*, manuscript in preparation). Because the DCM level is not strictly aphotic, we considered the separate and combined contributions of only aphotic and aphotic and DCM N<sub>2</sub> fixation rates for correlations with dissolved inorganic nutrients, bacterial abundance and activity, and DOM data (Table 2). Despite the lower number of variables that were significantly correlated with N<sub>2</sub> fixation once the DCM was removed from the correlations (Table 2), the recurrent observations of heterotrophic diazotrophs in the DCM waters of the Mediterranean Sea [Man-Aharonovich *et al.*, 2007; Yogevev *et al.*, 2011] suggest that DOM exerts a nonnegligible control of diazotrophic activity in this level. In our study N<sub>2</sub> fixation rates showed a significant negative correlation with N:P ratios, which agrees with the low N:P requirements of diazotrophic activity [Capone and Knapp, 2007]. Significant positive correlations were observed with prokaryotic abundance, prokaryotic heterotrophic production, and ETS, suggesting a relationship between aphotic N<sub>2</sub> fixation and bacterial activity, as has been previously observed before in the Levantine Basin [Rahav *et al.*, 2013c]. When examining the relationship between N<sub>2</sub> fixation and DOM compositional parameters (FDOM, CDOM, and FT-ICR-MS), we observed

**Table 2.** Bonferroni-Corrected Significant Spearman Correlations Between N<sub>2</sub> Fixation Rates, Organic Matter, and Bacterial Activity Variables

Variable	Correlation Coefficient	Significance (Two Tailed)
N:P	−0.924	<0.0005
POC	0.730	0.001
PON	0.785	<0.0005
TEP	0.781	<0.0005
Prokaryotic abundance <sup>a</sup>	0.845	<0.0005
Prokaryotic heterotrophic production	0.850	<0.0005
ETS	0.648	0.004
AOU <sup>a</sup>	−0.829	<0.0005
DOC <sup>a</sup>	0.810	<0.0005
Peak C/DOC <sup>a</sup>	−0.635	0.005
Peak T/DOC	0.627	0.002
Peak C/Peak T <sup>a</sup>	−0.833	<0.0005
<i>a</i> CDOM(254)/DOC	0.655	0.003
Double bond equivalent (DBE <sup>b</sup> )	−0.697	0.001
% peptides <sup>ca</sup>	0.525	0.025
% highly unsaturated compounds <sup>ca</sup>	−0.519	0.027
% unsaturated aliphatics <sup>ca</sup>	0.664	0.003
% oxygen-rich unsaturated aliphatics <sup>ca</sup>	0.558	0.016

<sup>a</sup>Correlations that remain significant after removing DCM N<sub>2</sub> fixation rates from the analysis.

<sup>b</sup>Koch and Dittmar [2006].

<sup>c</sup>Refers to DOM molecular formula information, categorization according to Seidel *et al.* [2014].

significant negative correlations with DBE as a measure for the unsaturation of the DOM, some optical indicators of the refractory nature of CDOM such as the C-specific humic-like substances (peak C/DOC) and the ratio of fluorescence of refractory humic-like to fresh protein-like substances (peak C/peak T), the percentage of highly unsaturated compounds, and AOU (Table 2). This suggests that N<sub>2</sub> fixation rates were higher in oxygenated waters containing presumably fresher organic materials, which is also supported by the positive correlations found with TEP, the carbon-specific fluorescence of protein-like substances (calculated as the ratio of peak T/DOC), the carbon-specific absorption coefficient at 254 nm of Weishaar *et al.* [2003] that indicated the presence of carbon double bonds in DOM, and the percentage of peptides and unsaturated aliphatics (as determined by FT-ICR-MS; Martínez-Pérez *et al.*, manuscript in preparation) (Table 2).

The relationship of heterotrophic N<sub>2</sub> fixation with the quantity and/or quality of organic matter has been seldomly studied. Rahav *et al.* [2013c] found a positive correlation between N<sub>2</sub> fixation and TEP in the meso-pelagic zone of the Levantine Basin. These authors suggested that TEP are utilized as organic carbon rich and low oxygen microniches for N<sub>2</sub> fixation in the water column. Other studies have also observed an enhancement of aphotic N<sub>2</sub> fixation rates upon the addition of amino acids or carbohydrates [Bonnet *et al.*, 2013; Rahav *et al.*, 2013c; Loescher *et al.*, 2014], suggesting that aphotic diazotrophic activity depends on labile organic matter inputs to deep water layers. Most studies on the phylogenetic *nifH* gene composition of aphotic waters have reported a high diversity (usually encompassing all four *nifH* clusters) [Hewson *et al.*, 2007; Hamersley *et al.*, 2011; Bonnet *et al.*, 2013; Rahav *et al.*, 2013c; Benavides *et al.*, 2015]. Although we did not characterize the diazotrophic community in this study, we assume that a diverse diazotroph community can also be expected too in the aphotic waters across the Mediterranean Basin, probably dominated by heterotrophic bacteria as reported in the sole study that has explored aphotic *nifH* diversity in Mediterranean Sea waters [Rahav *et al.*, 2013c]. Although *nifH* sequences from aphotic diazotrophs are becoming increasingly available, the metabolic properties of these phylotypes are largely unknown [Riemann *et al.*, 2010]. On the one hand, non-cyanobacterial diazotrophs may present a versatile metabolism, upregulated/downregulated according to changes in the composition and/or lability of DOM. Alternatively, their abundance and activity may fluctuate according to DOM dynamics. A pioneering genomics study targeting diazotrophic Alphaproteobacteria and Gammaproteobacteria isolated from coastal Denmark waters showed that these organisms devoted an important part of their genome to aromatic hydrocarbon metabolism, although diverse systems related to carbohydrate and fatty acid metabolism were also found [Bentzon-Tilia *et al.*, 2015], suggesting a flexible metabolism. Conversely, Severin *et al.* [2015] found that the taxonomic composition of non-cyanobacterial

diazotrophs in laboratory cultures changed dramatically upon exposure to a labile organic carbon mix and low oxygen conditions (10–16% saturation), indicative of a more specialized metabolism.

In the Mediterranean Sea, the high renewal rate of deep waters and their higher temperature as compared to other oceanic basins likely selects for diazotrophs different to those in low oxygen pelagic zones such as the oxycline of the Baltic Sea or the oxygen minimum zone off Peru [Dekaezemacker *et al.*, 2013; Farnelid *et al.*, 2013]. Modeling studies have found that the loss of fixed nitrogen via anammox and denitrification in oxygen minimum zones creates a deficit of nitrogen related to phosphorus which is replenished by N<sub>2</sub> fixation inputs [Deutsch *et al.*, 2007], which is supported by the significant N<sub>2</sub> fixation rates reported from these areas [Fernández *et al.*, 2011; Dekaezemacker *et al.*, 2013; Loescher *et al.*, 2014]. Because oxygen irreversibly destroys the nitrogenase enzyme system, autotrophic cyanobacterial diazotrophs use temporal and/or spatial segregation strategies to avoid oxygen deactivation [Berman-Frank *et al.*, 2003; Zehr, 2011], while heterotrophic diazotrophs are thought to either thrive in oxygen minimum zones or inhabit other oxygen-poor habitats or microhabitats such as suspended particles [Paerl and Prufert, 1987; Riemann *et al.*, 2010]. For example, aphotic N<sub>2</sub> fixation activity has been previously related to moderate or low dissolved oxygen levels [Hamersley *et al.*, 2011; Bonnet *et al.*, 2013; Loescher *et al.*, 2014]. Nevertheless, significant N<sub>2</sub> fixation rates have also been reported from fully oxygenated mesopelagic waters (>200 μmol L<sup>-1</sup>) [Rahav *et al.*, 2013c]. In this study we found significant negative correlations between AOU and N<sub>2</sub> fixation rates (Table 2), and N<sub>2</sub> fixation rates were undetectable in most oxygen minimum levels sampled along the transect (Figure 2). This suggests that aphotic diazotrophs in the Mediterranean Sea prefer newer more oxygenated waters presumably containing less refractory organic matter, which is characteristic of newly formed deep water masses [Santinelli *et al.*, 2006], as discussed above. Alternatively, and given the positive correlations observed between N<sub>2</sub> fixation and POC and PON concentrations (Table 2), we can also infer that aphotic diazotrophs use particles as oxygen-poor microzones in these oxygenated waters. However, trap-measured organic particle fluxes in the Mediterranean Sea are low given the ultraoligotrophic character of its overlying sunlit waters [Meador *et al.*, 2010], potentially rendering few microniches for aphotic diazotrophs relying on particles to avoid nitrogenase deactivation by oxygen.

#### Acknowledgments

The data included in this study are available from the authors on request. This work was supported by project HOTMIX—“Dark-ocean water mass boundaries and mixing zones as “hot-spots” of biodiversity and biogeochemical fluxes across the Mediterranean Sea and Eastern North Atlantic”—(CTM2011-30010-C02-01 and CTM2011-30010-C02-02) to J.A., X.A.A.S., and J.M. G. M.B. was funded by the Marie Skłodowska-Curie Actions of the European Union’s Seventh Framework Programme (FP7/2007-2013) under grant agreement 625185. S.B. was funded by the Institut de Recherche pour le Développement (IRD). A.M.M.-P. was funded by FPI and short stay fellowships from the Spanish Ministry of Economy and Competitiveness. M.N.-C. was supported by the CSIC Program “Junta para la Ampliación de Estudios” cofinanced by the ESF. I.B.-F. was supported by a collaborative grant with S.B. from MOST Israel and the High Council for Science and Technology (HCST)-France. We are grateful to the captain and crew of the R/V *Sarmiento de Gamboa* and the staff of the Unit of Marine Technology from the Spanish Research Council (UTM-CSIC), as well as O. Grosso for his help with IRMS analyses and K. Klaproth for her help with FT-ICR-MS analysis. The authors declare no conflict of interest.

#### 4. Conclusions

Aphotic N<sub>2</sub> fixation represented >50% of total water column integrated diazotrophic activity across the Mediterranean Sea basin. While deep diazotrophic activity may contribute importantly to the fixed nitrogen stock in mesopelagic to bathypelagic remote open ocean areas where terrigenous and atmospheric inputs are minimal, its contribution to the semienclined basin of the Mediterranean Sea, which presents a high drainage to surface area ratio, is negligible. We thus infer that the high N:P ratios recurrently observed in deep waters of the Mediterranean Sea are not greatly impacted by diazotrophic activity. Nevertheless, the N<sub>2</sub> fixation rates measured in the aphotic zone during this study are in the same range as those reported before for the euphotic zone in the Mediterranean Sea and aphotic N<sub>2</sub> fixation rates reported from the South and the North Pacific Oceans. N<sub>2</sub> fixation rates were higher in oxygenated waters with lower AOU and correlated positively to indicators of the labile nature of the available organic matter, suggesting that diazotrophs take advantage of the presence of the fresher organic material present in these newer, presumably more recently subducted water masses. Although evidence on active N<sub>2</sub> fixation in aphotic depths is accumulating, we currently ignore the metabolic requirements of these diazotrophs. Future investigations coupling genomic and proteomic studies combined with organic matter chemical analyses will further our understanding of the requirements and functioning of deep how deep diazotrophic communities.

#### References

- Aristegui, J., and M. F. Montero (1995), The relationship between community respiration and ETS activity in the ocean, *J. Plankton Res.*, *17*(7), 1563–1571.
- Aristegui, J., J. M. Gasol, C. M. Duarte, and G. J. Herndl (2009), Microbial oceanography of the dark ocean’s pelagic realm, *Limnol. Oceanogr.*, *54*(5), 1501–1529, doi:10.4319/lo.2009.54.5.1501.
- Bar-Zeev, E., T. Yogeve, D. Man-Aharonovich, N. Kress, B. Herut, O. Béjà, and I. Berman-Frank (2008), Seasonal dynamics of the endosymbiotic, nitrogen-fixing cyanobacterium *Richelia intracellularis* in the eastern Mediterranean Sea, *ISME J.*, *2*(9), 911–923, doi:10.1038/ismej.2008.56.
- Benavides, M., P. H. Moisanter, H. Berthelot, T. Dittmar, O. Grosso, and S. Bonnet (2015), Mesopelagic N<sub>2</sub> fixation related to organic matter composition in the Solomon and Bismarck Seas (Southwest Pacific), *PLOS ONE*, *10*(12), e0143775, doi:10.1371/journal.pone.0143775.

- Bentzon-Tilia, M., I. Severin, L. H. Hansen, and L. Riemann (2015), Genomics and ecophysiology of heterotrophic nitrogen-fixing bacteria isolated from estuarine surface water, *mBio*, 6(4), e00929-15, doi:10.1128/mBio.00929-15.
- Berman-Frank, I., P. Lundgren, and P. Falkowski (2003), Nitrogen fixation and photosynthetic oxygen evolution in cyanobacteria, *Res. Microbiol.*, 154(3), 157–164, doi:10.1016/S0923-2508(03)00029-9.
- Bethoux, J. P., and G. Copin-Montegut (1986), Biological fixation of atmospheric nitrogen in the Mediterranean Sea, *Limnol. Oceanogr.*, 31(6), 1353–1358, doi:10.4319/lo.1986.31.6.1353.
- Bonnet, S., O. Grosso, and T. Moutin (2011), Planktonic dinitrogen fixation along a longitudinal gradient across the Mediterranean Sea during the stratified period (BOUM cruise), *Biogeosciences*, 8(8), 2257–2267, doi:10.5194/bg-8-2257-2011.
- Bonnet, S., J. Dekaezemaeker, K. A. Turk-Kubo, T. Moutin, R. M. Hamersley, O. Grosso, J. P. Zehr, and D. G. Capone (2013), Aphotic N<sub>2</sub> fixation in the eastern tropical South Pacific Ocean, *PLOS ONE*, 8(12), e81265, doi:10.1371/journal.pone.0081265.s001.
- Boutrif, M., M. Garel, M. T. Cottrell, and C. Tamburini (2011), Assimilation of marine extracellular polymeric substances by deep-sea prokaryotes in the NW Mediterranean Sea, *Environ. Microbiol. Rep.*, 3(6), 705–709, doi:10.1111/j.1758-2229.2011.00285.x.
- Capone, D. G., and A. N. Knapp (2007), Oceanography: A marine nitrogen cycle fix?, *Nature*, 445(7124), 159–160, doi:10.1038/445159a.
- Coble, P. G. (1996), Characterization of marine and terrestrial DOM in seawater using excitation-emission matrix spectroscopy, *Mar. Chem.*, 51(6), 325–346, doi:10.1016/0304-4203(95)00062-3.
- Dabundo, R., M. F. Lehmann, L. Treibergs, C. R. Tobias, M. A. Altabet, P. H. Moisaner, and J. Granger (2014), The contamination of commercial <sup>15</sup>N<sub>2</sub> gas stocks with <sup>15</sup>N-labeled nitrate and ammonium and consequences for nitrogen fixation measurements, *PLOS ONE*, 9(10), e110335, doi:10.1371/journal.pone.0110335.
- Dekaezemaeker, J., S. Bonnet, O. Grosso, T. Moutin, M. Bressac, and D. G. Capone (2013), Evidence of active dinitrogen fixation in surface waters of the eastern tropical South Pacific during El Niño and La Niña events and evaluation of its potential nutrient controls, *Global Biogeochem. Cycles*, 27, 768–779, doi:10.1002/gbc.20063.
- Deutsch, C., J. L. Sarmiento, D. M. Sigman, N. Gruber, and J. P. Dunne (2007), Spatial coupling of nitrogen inputs and losses in the ocean, *Nature*, 445(7124), 163–167, doi:10.1038/nature05392.
- Dittmar, T., B. Koch, N. Hertkorn, and G. Kattner (2008), A simple and efficient method for the solid-phase extraction of dissolved organic matter (SPE-DOM) from seawater, *Limnol. Oceanogr. Methods*, 6, 230–235, doi:10.4319/lom.2008.6.230.
- Farnelid, H., A. F. Andersson, S. Bertilsson, W. A. Al-Soud, L. H. Hansen, S. Sørensen, G. F. Steward, Å. Hagström, and L. Riemann (2011), Nitrogenase gene amplicons from global marine surface waters are dominated by genes of non-cyanobacteria, *PLOS ONE*, 6(4), e19223, doi:10.1371/journal.pone.0019223.t001.
- Farnelid, H., M. Bentzon-Tilia, A. F. Andersson, S. Bertilsson, G. Jost, M. Labrenz, K. Jürgens, and L. Riemann (2013), Active nitrogen-fixing heterotrophic bacteria at and below the chemocline of the central Baltic Sea, *ISME J.*, 7(7), 1413–1423, doi:10.1038/ismej.2013.26.
- Fernández, C., L. Fariás, and O. Ulloa (2011), Nitrogen fixation in denitrified marine waters, *PLOS ONE*, 6(6), e20539, doi:10.1371/journal.pone.0020539.t001.
- Gasol, J. M., and P. del Giorgio (2000), Using flow cytometry for counting natural planktonic bacteria and understanding the structure of planktonic bacterial communities, *Sci. Mar.*, 64(2), 197–224.
- Grasshoff, K., M. Ehrhardt, K. Kremling, and M. Ehrhardt (Eds.) (1999), *Methods of Seawater Analysis*, 3rd ed., Verlag Chemie, Weinheim, Germany.
- Green, S. A., and N. V. Blough (1994), Optical absorption and fluorescence properties of chromophoric dissolved organic matter in natural waters, *Limnol. Oceanogr.*, 39(8), 1903–1916, doi:10.4319/lo.1994.39.8.1903.
- Großkopf, T., W. Mohr, T. Baustian, H. Schunck, D. Gill, M. M. M. Kuypers, G. Lavik, R. A. Schmitz, D. W. R. Wallace, and J. LaRoche (2012), Doubling of marine dinitrogen-fixation rates based on direct measurements, *Nature*, 488(7411), 361–364, doi:10.1038/nature11338.
- Hamersley, M. R., K. A. Turk, A. Leinweber, N. Gruber, J. P. Zehr, T. Gunderson, and D. G. Capone (2011), Nitrogen fixation within the water column associated with two hypoxic basins in the Southern California Bight, *Aquat. Microb. Ecol.*, 63(2), 193–205, doi:10.3354/ame01494.
- Hewson, I., P. H. Moisaner, K. M. Achilles, C. A. Carlson, B. D. Jenkins, E. A. Mondragon, A. E. Morrison, and J. P. Zehr (2007), Characteristics of diazotrophs in surface to abyssopelagic waters of the Sargasso Sea, *Aquat. Microb. Ecol.*, 46, 15–30, doi:10.3354/ame046015.
- Huertas, I. E., A. F. Ríos, J. García-Lafuente, G. Navarro, A. Makaoui, A. Sánchez-Román, S. Rodríguez-Galvez, A. Orbi, J. Ruiz, and F. F. Pérez (2012), Atlantic forcing of the Mediterranean oligotrophy, *Global Biogeochem. Cycles*, 26, GB2022, doi:10.1029/2011GB004167.
- Ibello, V., C. Cantoni, S. Cozzi, and G. Civitarese (2010), First basin-wide experimental results on N<sub>2</sub> fixation in the open Mediterranean Sea, *Geophys. Res. Lett.*, 37, L03608, doi:10.1029/2009GL041635.
- Kirchman, D., E. K'nees, and R. Hodson (1985), Leucine incorporation and its potential as a measure of protein synthesis by bacteria in natural aquatic systems, *Appl. Environ. Microbiol.*, 49(3), 599–607.
- Klawonn, I., G. LaviK, and P. Böning (2015), Simple approach for the preparation of <sup>15</sup>N<sub>2</sub>-enriched water for nitrogen fixation assessments: Evaluation, application and recommendations, *Front. Microbiol.*, doi:10.3389/fmicb.2015.00769.
- Koch, B. P., and T. Dittmar (2006), From mass to structure: An aromaticity index for high-resolution mass data of natural organic matter, *Rapid Commun. Mass Spectrom.*, 20(5), 926–932, doi:10.1002/rcm.2386.
- Krom, M. D., and B. Herut (2004), Nutrient budget for the Eastern Mediterranean: Implications for phosphorus limitation, *Limnol. Oceanogr.*, 49(5), 1582–1592, doi:10.4319/lo.2004.49.5.1582.
- Le Moal, M., H. Collin, and I. C. Biegala (2011), Intriguing diversity among diazotrophic picoplankton along a Mediterranean transect: A dominance of rhizobia, *Biogeosciences*, 8(3), 827–840, doi:10.5194/bg-8-827-2011.
- Loescher, C. R., et al. (2014), Facets of diazotrophy in the oxygen minimum zone waters off Peru, *ISME J.*, 8, 2180–2192, doi:10.1038/ismej.2014.71.
- Ludwig, W., E. Dumont, M. Meybeck, and S. Heussner (2009), River discharges of water and nutrients to the Mediterranean and Black Sea: Major drivers for ecosystem changes during past and future decades? *Prog. Oceanogr.*, 80(3–4), 199–217, doi:10.1016/j.pocean.2009.02.001.
- Luna, G. M., S. Bianchelli, F. Decembrini, E. De Domenico, R. Danovaro, and A. Dell'Anno (2012), The dark portion of the Mediterranean Sea is a bioreactor of organic matter cycling, *Global Biogeochem. Cycles*, 26, GB2017, doi:10.1029/2011GB004168.
- Macías, D., A. Stips, and E. Garcia-Gorriz (2014), The relevance of deep chlorophyll maximum in the open Mediterranean Sea evaluated through 3D hydrodynamic-biogeochemical coupled simulations, *Ecol. Modell.*, 281, 26–37, doi:10.1016/j.ecolmodel.2014.03.002.
- Man-Aharonovich, D., N. Kress, E. B. Zeev, I. Berman-Frank, and O. Béjà (2007), Molecular ecology of *nifH* genes and transcripts in the eastern Mediterranean Sea, *Environ. Microbiol.*, 9(9), 2354–2363, doi:10.1111/j.1462-2920.2007.01353.x.
- Manca, B., M. Burca, A. Giorgetti, C. Coatanoan, M. J. Garcia, and A. Iona (2004), Physical and biochemical averaged vertical profiles in the Mediterranean regions: An important tool to trace the climatology of water masses and to validate incoming data from operational oceanography, *J. Mar. Syst.*, 48(1–4), 83–116, doi:10.1016/j.jmarsys.2003.11.025.

- Markaki, Z., M. D. Lojçe-Pilot, K. Violaki, L. Benyahya, and N. Mihalopoulos (2010), Variability of atmospheric deposition of dissolved nitrogen and phosphorus in the Mediterranean and possible link to the anomalous seawater N/P ratio, *Mar. Chem.*, 120(1–4), 187–194, doi:10.1016/j.marchem.2008.10.005.
- Meador, T. B., A. Gogou, G. Spyres, G. J. Herndl, E. Krasakopoulou, S. Psarra, T. Yokokawa, D. De Corte, V. Zervakis, and D. J. Repeta (2010), Biogeochemical relationships between ultrafiltered dissolved organic matter and picoplankton activity in the Eastern Mediterranean Sea, *Deep-Sea Res., Part II*, 57(16), 1460–1477, doi:10.1016/j.dsr2.2010.02.015.
- Montoya, J. P., M. Voss, P. Kahler, and D. G. Capone (1996), A simple, high-precision, high-sensitivity tracer assay for N<sub>2</sub> fixation, *Appl. Environ. Microbiol.*, 62(3), 986–993.
- Mulholland, M. R., P. W. Bernhardt, J. L. Blanco-Garcia, A. Mannino, K. Hyde, E. Mondragon, K. Turk, P. H. Moisander, and J. P. Zehr (2012), Rates of dinitrogen fixation and the abundance of diazotrophs in North American coastal waters between Cape Hatteras and Georges Bank, *Limnol. Oceanogr.*, 57(4), 1067–1083, doi:10.4319/lo.2012.57.4.1067.
- Paerl, H. W., and L. E. Prufert (1987), Oxygen-poor microzones as potential sites of microbial N<sub>2</sub> fixation in nitrogen-depleted aerobic marine waters, *Appl. Environ. Microbiol.*, 53(5), 1078–1087.
- Pantoja, S., D. J. Repeta, J. P. Sachs, and D. M. Sigman (2002), Stable isotope constraints on the nitrogen cycle of the Mediterranean Sea water column, *Deep-Sea Res., Part I*, 49(9), 1609–1621, doi:10.1016/S0967-0637(02)00066-3.
- Passow, U., and A. L. Alldredge (1995), A dye-binding assay for the spectrophotometric measurement of transparent exopolymer particles (TEP), *Limnol. Oceanogr.*, 40(7), 1326–1335, doi:10.4319/lo.1995.40.7.1326.
- Pujo-Pay, M., P. Conan, L. Oriol, V. Cornet-Barthaux, C. Falco, J. F. Ghiglione, C. Goyet, T. Moutin, and L. Prieur (2011), Integrated survey of elemental stoichiometry (C, N, P) from the western to eastern Mediterranean Sea, *Biogeosciences*, 8(4), 883–899, doi:10.5194/bg-8-883-2011.
- Rahav, E., B. Herut, A. Levi, M. R. Mulholland, and I. Berman-Frank (2013a), Springtime contribution of dinitrogen fixation to primary production across the Mediterranean Sea, *Ocean Sci.*, 9(3), 489–498, doi:10.5194/os-9-489-2013.
- Rahav, E., B. Herut, M. R. Mulholland, B. Voß, D. Stazic, C. Steglich, W. R. Hess, and I. Berman-Frank (2013b), Contribution of dinitrogen fixation to bacterial and primary productivity in the Gulf of Aqaba (Red Sea), *Biogeosci. Discuss.*, 10(6), 10,327–10,361, doi:10.5194/bgd-10-10327-2013.
- Rahav, E., E. Bar-Zeev, S. Ohayon, H. Elifantz, N. Belkin, B. Herut, M. R. Mulholland, and I. Berman-Frank (2013c), Dinitrogen fixation in aphotic oxygenated marine environments, *Front. Microbiol.*, 4(227), 1–11, doi:10.3389/fmicb.2013.00227/abstract.
- Rahav, E., B. Herut, M. R. Mulholland, N. Belkin, H. Elifantz, and I. Berman-Frank (2015), Heterotrophic and autotrophic contribution to dinitrogen fixation in the Gulf of Aqaba, *Mar. Ecol. Prog. Ser.*, 522, 67–77, doi:10.3354/meps11143.
- Riemann, L., H. Farnelid, and G. F. Steward (2010), Nitrogenase genes in non-cyanobacterial plankton: Prevalence, diversity and regulation in marine waters, *Aquat. Microb. Ecol.*, 61(3), 235–247, doi:10.3354/ame01431.
- Rodellas, V., J. Garcia-Orellana, P. Masqué, M. Feldman, and Y. Weinstein (2015), Submarine groundwater discharge as a major source of nutrients to the Mediterranean Sea, *Proc. Natl. Acad. Sci. U.S.A.*, 112(13), 3926–3930, doi:10.1073/pnas.1419049112.
- Sachs, J. P. (1999), Oligotrophy and nitrogen fixation during eastern Mediterranean sapropel events, *Science*, 286(5449), 2485–2488, doi:10.1126/science.286.5449.2485.
- Santinelli, C., B. B. Manca, G. P. Gasparini, L. Nannicini, and A. Seritti (2006), Vertical distribution of dissolved organic carbon (DOC) in the Mediterranean Sea, *Clim. Res.*, 31, 205–216, doi:10.3354/cr031205.
- Santinelli, C., L. Nannicini, and A. Seritti (2010), DOC dynamics in the meso and bathypelagic layers of the Mediterranean Sea, *Deep-Sea Res., Part II*, 57(16), 1446–1459, doi:10.1016/j.dsr2.2010.02.014.
- Seidel, M., M. Beck, T. Riedel, and H. Waska (2014), Biogeochemistry of dissolved organic matter in an anoxic intertidal creek bank, *Geochim. Cosmochim. Acta*, 140, 418–434, doi:10.1016/j.gca.2014.05.038.
- Severin, I., M. Bentzon-Tilia, P. H. Moisander, and L. Riemann (2015), Nitrogenase expression in estuarine bacterioplankton influenced by organic carbon and availability of oxygen, *FEMS Microbiol. Lett.*, doi:10.1093/femsle/fnv105.
- Smith, D. C., and F. Azam (1992), A simple, economical method for measuring bacterial protein synthesis rates in seawater using <sup>3</sup>H-leucine, *Mar. Microb. Food Webs*, 6(2), 107–114.
- United Nations Educational, Scientific and Cultural Organization (1986), Progress on oceanographic tables and standards 1983–1986: Work and recommendations of the UNESCO/SCOR/ICES/IAPSO Joint Panel.
- Weishaar, J. L., G. R. Aiken, B. A. Bergamaschi, M. S. Fram, R. Fujii, and K. Mopper (2003), Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon, *Environ. Sci. Technol.*, 37(20), 4702–4708, doi:10.1021/es030360x.
- Yogev, T., E. Rahav, E. Bar-Zeev, D. Man-Aharonovich, N. Stambler, N. Kress, O. Béjà, M. R. Mulholland, B. Herut, and I. Berman-Frank (2011), Is dinitrogen fixation significant in the Levantine Basin, East Mediterranean Sea?, *Environ. Microbiol.*, 13(4), 854–871, doi:10.1111/j.1462-2920.2010.02402.x.
- Zehr, J. P. (2011), Nitrogen fixation by marine cyanobacteria, *Trends Microbiol.*, 19(4), 162–173, doi:10.1016/j.tim.2010.12.004.
- Zehr, J. P., B. D. Jenkins, S. M. Short, and G. F. Steward (2003), Nitrogenase gene diversity and microbial community structure: A cross-system comparison, *Environ. Microbiol.*, 5(7), 539–554, doi:10.1046/j.1462-2920.2003.00451.x.