# How to find information on national food and nutrient consumption surveys across Europe: systematic literature review and questionnaires to selected country experts are both good strategies

Maria Blanquer<sup>1</sup>, Alicia García-Álvarez<sup>1</sup>, Lourdes Ribas-Barba<sup>1</sup>, Trudy M. A. Wijnhoven<sup>2</sup>, Garden Tabacchi<sup>3</sup>, Mirjana Gurinovic<sup>4</sup> and Lluís Serra-Majem<sup>1,5</sup>\*

(Received 4 February 2009 - Revised 6 May 2009 - Accepted 1 June 2009)

The present research was conducted within the framework of the EURopean micronutrient RECommendations Aligned project. In order to identify the best practice in assessing nutrient intakes, a search strategy for collecting data from national food consumption surveys/studies in Europe was developed. Systematic literature searches were carried out on twenty-eight European and the four European Free Trade Association countries. A questionnaire was also sent to two to five experts in each country. Systematic reviews using PubMed yielded 12 703 abstracts that were reduced to 200 studies using inclusion and exclusion criteria. Similarly, a search of ministry web sites yielded 3033 hits, and subsequently reduced to nine surveys. Belgium, France, Germany, Ireland, Sweden, Spain and the United Kingdom were the countries with most data and Slovenia and Liechtenstein were those with the least. Seventy-eight expert questionnaires were obtained from all countries except for Liechtenstein, Luxembourg and Slovakia. Detailed results and references are given. A systematic search and questionnaires are equally good at identifying national surveys across countries. Literature searching provides globally accessible and objective information albeit limited, whereas the questionnaire provides information that, depending upon responders, can be more complete. A combination of both strategies is recommended.

National nutrition surveys: Search methodology: European projects: Best practice

A specific activity of the EURopean micronutrient RE-Commendations Aligned (EURRECA) project is to identify the best practice in assessing nutrient intakes with the objective of harmonising micronutrient recommendations for population groups in Europe. To achieve this end, it is important to find the existing information about those national surveys of dietary intake in as many countries as possible. Most European countries have carried out such surveys. However, the year they were conducted, the population groups included, the age categories considered, the objectives of the survey and the dietary methods used for data collection differ considerably between countries (1-21). There are several

projects presently working on providing comparable intercountry food and nutrient intake data, namely the Data Food Networking project<sup>(10)</sup>, the European Prospective Investigation into Cancer and Nutrition project<sup>(11)</sup>, the European Food Consumption Survey Method project<sup>(12)</sup>, the European Nutrition and Health Report I (2004)<sup>(13)</sup>, the Healthy Lifestyle in Europe by Nutrition in Adolescence project<sup>(14)</sup>, the European Food Consumption Validation project<sup>(15)</sup>, the European Information Campaign on Diet and Nutrition)<sup>(16)</sup>, the EURODIET project (Nutrition and Diet for Healthy Lifestyles in Europe)<sup>(17)</sup>, the European Food Information

Abbreviation: EURRECA, European micronutrient recommendations aligned.

On behalf of EURRECA's RA 1·1 'Intake Methods' members: Serra-Majem L (Coordinator), Cavelaars A, Dhonukshe-Rutten R, Doreste JL, Frost-Andersen L, García-Álvarez A, Glibetic M, De Groot L, Henríquez-Sánchez P, Naska A, Ngo J, Novakovic R, Ortiz-Andrellucchi A, Øverby NC, Gurinovic M, Pijls L, Ranic M, Ribas-Barba L, Ristic-Medic D, Román-Viñas B, Ruprich J, Saavedra-Santana P, Sánchez-Villegas A, Tabacchi G, Tepsic J, Trichopoulou A, van't Veer P, Vucic V, Wijnhoven TMA.

<sup>&</sup>lt;sup>1</sup>Community Nutrition Research Centre of the Nutrition Research Foundation, University of Barcelona Science Park, Baldiri Reixac 4, 08028 Barcelona, Spain

<sup>&</sup>lt;sup>2</sup>Noncommunicable Diseases and Environment, World Health Organization Regional Office for Europe, Scherfigsvej 8, DK-2100 Copenhagen Ø, Denmark

<sup>&</sup>lt;sup>3</sup>Institute of Physiology and Human Nutrition, University of Palermo, Via Augusto Elia 3, 90127 Palermo, Italy

<sup>&</sup>lt;sup>4</sup>Department for Nutrition and Metabolism, Institute for Medical Research, University of Belgrade, Tadeusa Koscuska 1, Belgrade 11000, Serbia and Montenegro

<sup>&</sup>lt;sup>5</sup>Department of Clinical Sciences, University of Las Palmas de Gran Canaria, PO Box 550, 35080 Las Palmas de Gran Canaria, Spain

<sup>\*</sup> Corresponding author: Professor Lluís Serra-Majem, fax +34 93 403 4543, email lserra@dcc.ulpgc.es

Resource Network project<sup>(18)</sup>, the EURONUT (European Union's Concerted Action on Nutrition and Health) – Survey in Europe on Nutrition and the Elderly: a Concerted Action project<sup>(19)</sup>, the HECTOR project (Eating out: Habits, Determinants, and Recommendations for Consumers and the European Catering Sector)<sup>(20)</sup>, and the WHO MONICA project (Multinational Monitoring of trends and determinants in Cardiovascular disease)<sup>(21)</sup>.

The purpose of the present paper is to design a search strategy that facilitates the finding of the most representative and recent information regarding dietary data collection and nutrient intake adequacy assessment methods in twenty-eight European countries and the four European Free Trade Association countries, and present an overview of the findings. Food consumption surveys/studies carried out on representative samples of an apparently healthy population represent the first target of this search strategy.

# Methodology

The steps followed to find the information needed and to store and present the results are shown in Fig. 1. Three working directions were used to achieve desired objectives: working in cascade (vertical), in parallel and across (horizontal).

## Search strategy

The search for nutritional surveys and/or related papers on the intakes of nutrients, consumption of foods and/or food groups was carried out for the twenty-eight European

countries (Austria, Belgium, Bulgaria, the Czech Republic, Cyprus, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, The Netherlands, Malta, Poland, Portugal, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden and the United Kingdom) and for the four European Free Trade Association countries (Iceland, Liechtenstein, Norway and Switzerland). As such, the total number of countries included in the search was thirty-two.

The search criteria applied included the following guide-

- 1. Limit the review to European countries (twenty-eight) and European Free Trade Association countries.
- Identify the best studies based on external validity (in countries where there are no national data, studies with a smaller sample could be included, but they should be representative, from regions or covering large areas of a country). Convenience and local samples were not considered.
- 3. Limit the review to food consumption studies conducted after 1990 (in those countries not having such data, the period 1985–90 was considered) along with meeting the earlier established minimum criteria. In the case of multiple studies, the newest one meeting search criteria were selected when a similar methodology was applied.
- 4. Choose the highest population level (national, regional).
- Include healthy children, adolescents, adults and elderly populations.
- 6. Carry out searches through PubMed, ministry websites and books/reports/journal supplements by country.




Fig. 1. Methodological framework: sources of information. EU, European Union; EURRECA, European micronutrient recommendations aligned. \*RA 1·1, Research Activity 1·1 of the EURRECA Network of Excellence. \*\*Per expert to whom it was sent.

7. Key words used in the search: 'national nutrition survey'; 'food and nutrition survey'; 'dietary intake'; 'food intake'; 'food habits'; 'meal habits'; 'food consumption pattern'; 'dietary habits'; 'dietary record'; 'dietary survey'; 'nutrient intake'; 'diet questionnaires'; 'diet cross-sectional study'; 'food cross-sectional studies'; 'nutrient cross-sectional studies'; 'diet evaluation'; 'food evaluation'; 'diet assessment'; 'food assessment'; 'nutrient data collection'; 'food data collection'.

Systematic PubMed search. In the systematic PubMed search, twenty-one key words/search terms were used, following the same order for each country. In addition, the limits 'humans' and 'from 1990' were used. No language restrictions were applied.

A first selection was carried out with all the abstracts obtained per country applying the following exclusion criteria:

- 1. Papers evaluating the association between nutritional status and illness or any other health problem.
- 2. Papers about tobacco or drug consumption.
- Papers that evaluated pesticides, food contaminants and food additives.
- 4. Papers that evaluated the nutritional status of a population exclusively through biochemical parameters or alternatively only provided reference values.
- Those addressing nutrition education and interventions or conducting a needs assessment.
- Papers on patients undergoing surgical procedures or those who were hospitalised.
- 7. Studies that evaluated the relationship between food and biological variables or that exclusively measured biological, anthropometric, socio-demographic or functional variables, as well as those related to the utilisation of health services in nutrition studies.
- 8. Papers that described statistical models or sampling techniques.
- 9. Papers not having an abstract.

A second stage of selection was carried out in a step-wise manner as follows:

Firstly, applying exclusion criteria to abstract titles and secondly to the contents of selected abstracts.

The final abstracts list was used to obtain surveys and related papers through (in this order) PubMed, on-line library of the University of Barcelona. All results, surveys, reports or papers were stored separately in files for each country.

Ministry website search. A similar search was carried out in parallel through the websites of ministries (departments of health and food and agriculture) of each country, following the same search criteria as for the PubMed search. In this case, the list of key words/search terms was reduced to thirteen: 'national nutrition survey'; 'national nutrition report'; 'food and nutrition survey'; 'food consumption report'; 'food consumption survey'; 'food consumption patterns'; 'dietary survey'; 'dietary report'; 'diet evaluation'; 'diet assessment'. In ministries of health and food and agriculture websites, the term 'in humans' was added.

The selection of hits from the 'ministries websites' search was carried out doing a pre-selection of all initial results. This pre-selection consisted of keeping only those relevant to food consumption surveys. Subsequently, the same exclusion criteria used for the PubMed search was applied. Finally, only those that complied with the five first search criteria were selected.

European projects. Twelve European projects were reviewed and all relevant information on food consumption surveys in the participating countries extracted. Key contacts were also extracted for completion of the Experts list (see section down mentioned later).

#### Ouestionnaire criteria and design

A questionnaire was designed including forty-nine questions with the objective of obtaining the greatest amount of information in parallel with the systematic search. In addition, the questionnaire was designed to complete study data that were missing due to survey inaccessibility (based on language problems or because studies were not published, etc.). The questionnaire also aimed at obtaining the contact details of the responsible person(s) for the surveys, with the aim of enlisting their assistance in the completion of the missing information. In order to complete a questionnaire on a particular survey, the survey needed to meet the following priority criteria:

- 1. Be the most recent.
- Be the most representative of the country's population or of part of it: i.e. ideally at the national (country) population level; if not available, at the regional/province/ department/county/population level.
- 3. Be conducted after 1990 (in the case that the only available survey had been conducted between 1985 and 1990, exceptions could be made and it could be considered).
- 4. Have the highest external validity: i.e. the survey results were generalisable to the larger population. External validity is low when surveys using human participants have small samples and/or are obtained from a single geographic location.

#### Questionnaire development

The questionnaire was initially designed to have an online format, but due to logistic inconveniences, it was decided that it would be distributed electronically by email as a result from consensus reached by members of EURRECA's Research Activity 1·1 (Intake Methods), other EURRECA partners, and members of the European Health and Nutrition Report II<sup>(13)</sup>, who were sent a preliminary version to review. The response of this massive mailing was tracked, correspondence exchanged, and comments/suggestions were considered and incorporated into the design of a final questionnaire, which was sent to two to five expert contacts from each of the thirty-two countries.

## Experts list

An Experts list was compiled, which was obtained from internal contact lists and from external and public sites (PubMed, ministry websites, etc.). The list included thirty-two countries as agreed upon by consensus with relevant EURRECA partners. The response of the experts was tracked

and filed; for those countries who did not respond, 'reminder' messages were sent and the response was also monitored. A follow-up message was also sent to clarify doubts related to questionnaire responses for certain countries (i.e. some questionnaires received had no answers to certain questions or contradictory answers, and as a result, it was necessary to obtain further clarification, especially regarding questions on nutrient intake adequacy). Additional 'reminder' messages were sent to the four remaining non-responding countries; however, these efforts were fruitless.

All relevant information found on food consumption from surveys, their related papers and other studies per country was included in the 'Country information found' table. This table, which summarises the outcomes of the search, comprises a total of forty-eight fields (columns) by country included (rows). The fields to be completed per country were as follows: study name; realisation period; financing body; organisation conducting study; year of publication; authors; publication type and references; language; scope or level of study; periodicity; study population; study's main objective; study type; representativeness; initial sample population; sampling methods; participation/response rate (%); sex (n); age groups (n); dietary survey methods; survey method for data collection; vitamin and mineral supplements; dietary supplement use; food portions estimation; prepared foods/meals and recipes; databases for data collection; data format for analysis; food codification; food composition table used: functional and fortified foods included; physical activity evaluation/assessment; nutrient data adjusted by; food groups; results; reference nutrient intakes used; nutritional adequacy assessment methods; anthropometry; biochemistry; source of information. Although the majority of surveys/ papers used to complete the table were found through PubMed, some of them were obtained through other sources such as available data from European projects or other EURRECA partners.

Once the table was completed with the available/accessible information for a particular country (paying special attention to the dietary data collection and nutritional adequacy assessment methods as a main objective), it was sent together with the questionnaire to the country experts.

The surveys/studies contained in the 'Country information found' table were then compared with the surveys/studies received as completed questionnaires and duplicates were excluded.

#### Results

Search results

Systematic PubMed search. This search strategy resulted in the identification of 12 703 abstracts. Review of titles and application of exclusion criteria resulted in 439 selected abstracts. Applying exclusion criteria to full abstracts led to a total selection of 200 final abstracts as shown in Fig. 2.

According to these results, the countries with no data available/accessible through PubMed were Slovenia and Liechtenstein, the countries with the least information were Serbia, Estonia, Bulgaria (with one result) and Cyprus, Latvia, Luxembourg; Malta, Portugal and Slovakia (with two results). By contrast, countries such as Belgium, France, Germany,




Fig. 2. Selection process applied to PubMed papers.

Ireland, Sweden, Spain and the United Kingdom had the highest amount of available/accessible data (with approximately twelve to fifteen results).

Ministry website search. This search identified 3033 results. Pre-selection consisted of keeping only those relevant to food consumption surveys. Subsequently, the same exclusion criteria used for the PubMed search was applied and resulted in seventy hits. Finally, only those that complied with the first five search criteria were selected, resulting in a total final selection of nine results.

#### Questionnaire results

Expert's response. The questionnaire was sent to two to five expert contacts from each of the thirty-two countries. A total of eleven countries responded within a month; seventeen other countries responded after the reminder message and four countries did never respond.

Response rate. The majority of countries (twenty-seven out of thirty-two) sent a completed questionnaire, obtaining a total of seventy-eight questionnaires, in other words, a response rate of 92.2%. Eleven countries answered after the first email sent, seventeen countries after the 'reminder' email and no response was ever obtained from countries such as Liechtenstein, Luxembourg and Slovakia. A final table containing 118 surveys/studies (forty plus seventy-eight) from all countries was compiled.

Table 1 summarises data extracted from a database designed for the storage of all results for each keyword per country.

## Discussion

The present review process has involved the combination of several strategies converging into the same objective: to identify available information on food consumption and nutrient intake from each of the twenty-eight European and four European Free Trade Association countries considered. In order to establish recommendations for future reviews of this nature, a previous evaluation of the strengths and limitations of each strategy followed is useful.

The combination of a lack of consensus for protocols on conducting grey literature systematic searches (mainly Google) and the complexity of limiting the extent of irrelevant results led to the elimination of this method as part of the search strategy. Consequently, the search was limited to

Table 1. Number and selection of abstracts found through PubMed, ministries and questionnaires

Country	PubMed results				
	Total abstracts	Selected abstracts	Final abstracts	Ministries	Questionnaires
Austria	126	12	4 <sup>(23–26)</sup>	0	1 <sup>(27)</sup>
Belgium	292	18	12 <sup>(28-39)</sup>	0	1 <sup>(40)</sup>
Bulgaria	39	1	1 <sup>(41)</sup>	0	1 <sup>(42)</sup>
Cyprus	14	4	2(43,44)	0	3 <sup>(45-47)</sup>
Czech Republic	88	11	8 <sup>(48-55)</sup>	0	1 <sup>(56)</sup>
Denmark .	595	41	8 <sup>(57-64)</sup>	0	1 <sup>(65)</sup>
Estonia	45	7	1 <sup>(66)</sup>	0	1 <sup>(67)</sup>
Finland	704	21	7 <sup>(68-74)</sup>	1 <sup>(75)</sup>	1 <sup>(76)</sup>
France	836	56	12 <sup>(77-88)</sup>	0	2 <sup>(89,90)</sup>
Germany	902	39	13 <sup>(91-103)</sup>	0	1 <sup>(104)</sup>
Greece	369	21	9(105-113)	0	5 <sup>(114–118)</sup>
Hungary	174	7	6 <sup>(119–124)</sup>	0	5(125-129)
Iceland	88	5	5(130-134)	0	1 <sup>(135)</sup>
Ireland	295	26	13 <sup>(136-148)</sup>	2 <sup>(149,150)</sup>	1 <sup>(151)</sup>
Italy	1209	17	9(152-160)	0	2(161,162)
Latvia	16	4	2(163,164)	0	1 <sup>(165)</sup>
Liechtenstein	3	0	0	0	0
Lithuania	57	10	7 <sup>(166–172)</sup>	0	1 <sup>(173)</sup>
Luxembourg	11	2	2 <sup>(174,175)</sup>	0	0
Malta	17	2	2 <sup>(176,177)</sup>	0	1 <sup>(178)</sup>
Netherlands	846	25	8 <sup>(179–186)</sup>	2 <sup>(187,188)</sup>	1 <sup>(189)</sup>
Norway	498	16	5 <sup>(190-194)</sup>	0	7 <sup>(195–201)</sup>
Poland	572	16	9(202-210)	0	4 <sup>(211–214)</sup>
Portugal	110	4	2(215,216)	Ö	4 <sup>(217–220)</sup>
Romania	50	4	4 <sup>(221–224)</sup>	0	2(225,226)
Serbia	149	1	1 <sup>(227)</sup>	0	5 <sup>(228-232)</sup>
Slovakia	42	2	2(233,234)	0	0
Slovenia	20	0	0	0	1 <sup>(235)</sup>
Spain	1014	31	15(236-250)	0	8(251-258)
Sweden	989	13	12 <sup>(259–269)</sup>	0	3 <sup>(270–272)</sup>
Switzerland	297	7	5 <sup>(273–277)</sup>	2 <sup>(278,279)</sup>	2(280,281)
United Kingdom	2236	16	14 <sup>(282–295)</sup>	2 <sup>(296,297)</sup>	10 <sup>(298-307)</sup>
Total	12733	439	200	9	78

PubMed and ministry websites. In order to complete the missing information essential for the task, a questionnaire was sent to the country experts.

Examples of the challenges in undertaking this kind of review include, among others, the diversity of languages, incomplete information in the abstracts, difficulty in obtaining full papers as well as delays in questionnaire responses, probably due to the length of the questionnaire. Although assistance from project researchers and external experts resolved part of these problems, generally, such inconveniences reinforce the need for consensus on the methodology to be followed, in order to limit the barriers to an effective process that tend to occur at the organisational stage.

In regard to the positive aspects of the work carried out, it can be said that the literature review provided an extensive overview of the available information coming from four different sources (PubMed, ministry websites, books/reports, European projects and congresses/expert meetings); it also confirmed the importance of consensus and training from the outset of the task. Regarding the questionnaire, it was found to be an excellent and straightforward tool for obtaining expert information, as most of the questions were answered, which thus facilitated data analysis (22) and enhanced the strengthening of networks at the European level.

In conclusion, the systematic search (PubMed in particular) and the questionnaire completed by experts are the two strategies that yielded the most satisfactory results, and which can also be compared among countries. PubMed provided objective, globally accessible, although possibly limited information (i.e. only abstracts and not full papers are available). The expert-completed questionnaire provided more concrete straightforward information, which was however, based on the expert's knowledge or willingness to collaborate, and thus subject to certain limitations<sup>(22)</sup>. These two approaches are useful, although both could be improved. As such, a combination of these two strategies is therefore recommended.

#### Acknowledgements

The studies reported herein have been carried out within the EURRECA Network of Excellence (www.eurreca.org), financially supported by the Commission of the European Communities, specific Research, Technology and Development Programme Quality of Life and Management of Living Resources, within the Sixth Framework Programme, contract no. 036196. The present report does not necessarily reflect the Commission's views or its future policy in this area. M. B. developed and carried out the search strategy, undertook analysis and wrote the first draft of the paper. A. G.-A. developed and carried out the search strategy, contributed to writing of first draft and commented on following drafts of the paper. L. R.-B. participated in the planning of the strategy, supervised the work and commented on drafts of the paper.

T. M. A. W. participated in the planning of the strategy developed and commented on drafts of the paper. G. T. participated in the planning of the strategy and commented on drafts of the paper. M. G. participated in the planning of the strategy and commented on drafts of the paper. L. S.-M. participated in the planning of the strategy, directed and supervised the work and commented on all drafts of the paper. The authors would like to thank the following persons for their interest shown in the present work and the effort made when completing questionnaires and exchanging correspondence to 'fill in the gaps': Elmadfa I; Freisling H (Department of Nutritional Sciences, University of Vienna - Austria); Van Oyen H; Vandevijvere S (Scientific Institute of Public Health -Belgium); Angelova KC (National Center of Public Health Protection - Bulgaria); Panagiotakos DB; Polychronopoulos EA (Harokopio University - Cyprus); Lazarou C (Harokopio University - Cyprus); Jiri Ruprich (National Institute of Public Health - Czech Republic); Groth M (National Food Institute – Denmark); Vaask S (University of Technology – Estonia); Valsta L; Vartiainen E; Laatikainen T (National Public Health Institute - Finland); Hercberg S; Castetbon K (Institut de Veille Sanitaire - France); Kersting M (Research Institute of Child Nutrition, Dortmund – Germany); Krems C (Federal Research Institute of Nutrition – Germany); Roma-Giannikou E; Manios Y; Benetou V; Trichopoulou A (University of Athens - Greece); Gabor Z (National Public Health and Medical Officer Service (Állami Népegészségügyi és Tisztiorvosi Szolgálat) – Hungary); Jonsdottir E; Thorsdottir I (University Hospital – Iceland); Steingrímsdóttir L; Þorgeirsdóttir H (Public Health Institute of Iceland - Iceland); Wall P; Gibney M (Trinity College Dublin, University College Cork - Ireland; University of Ulster - Northern Ireland); D'Amicis A; Cialfa E; Turín A; Saba A (Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione - Italy); La Vecchia C; Lombardy B ('Mario Negri' Institute - Italy); Joffe R; Santare D (Center of Food and Veterinary Service of Latvia - Latvia); Barzda A (National Nutrition Center of the Lithuanian Ministry of Health - Lithuania); Pace Asciak R; Calleja N; Ellul V; Grima A (Department of Health Information and Research - Malta); van Rossum C; Ocke M (Rijksinstituut voor Volksgezondheid en Milieu = National Institute for Public Health and the Environment - The Netherlands); Johansson L; Frost-Andersen L; Lande B (University of Oslo - Norway); Szponar L; Charzewska J; Brzozowska A (National Food and Nutrition Institute - Poland); Rywik S (National Institute of Cardiology - Poland); Barros H; Moreira P; Ramos E; Barros R; Lopes C (University of Porto - Portugal); Zapirtan H; Vlad M (Institute of Public Health -Romania); Gurinovic M (Department of Nutrition, Institute of Medical Research, University of Belgrade - Serbia); Vukmirovic D; Mijakovac N (Statistical Office Republic Serbia – Serbia); Koch V (University of Ljubljana – Slovenia); Ortega RM (University Complutense of Madrid - Spain); Vioque J (University of Alicante – Spain); Tur JA (University of Balearic Islands - Spain); Enghardt Barbier H (National Food Administration and GfK Sweden - Sweden); Becker W (NFA and Statistics Sweden - Sweden); Gaspoz JM (University of Geneva - Switzerland); Cade J (University of Leeds - United Kingdom) and Margetts B (University of Southampton - United Kingdom).

There is no conflict of interest to report.

#### References

- Elmadfa I & König J (editors) (2001) 17th International Congress of Nutrition. Abstracts. Ann Nutr Metab 45, 1–636.
- Gilloteau P, Lemonnier D & Lairon D (editors) (2007) 10th European Nutrition Conference. Abstracts. *Ann Nutr Metab* 51, 1–421.
- Margetts B & Nelson M (editors) (1997) Design Concepts in Nutritional Epidemilogy, 2nd ed. Oxford: Oxford University Press.
- Frank GC (1996) Applied Nutrition Epidemiology. Gaithersburg, MD: Aspen Press.
- Riboli E, Hunt KJ, Slimani N, et al. (2002) European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. Public Health Nutr 5, 1113–1124.
- Slimari N, Kaaks R, Ferrari P, et al. (2002) European Prospective Investigation into Cancer and Nutrition (EPIC) calibration study: rationale, design and population characteristics. Public Health Nutr 5, 1125–1146.
- Leitzmann C & Cannon G (editors) (2005) The New Nutrition Science project. *Public Health Nutr* 8, 724–804.
- 8. Yngve A (2007) Food and drink marketing to children: a continuing scandal. *Public Health Nutr* **10**, 971–972.
- Sarault NA (1997) 16th International Congress of Nutrition. Abstracts. Ottawa: National Research Council Canada.
- Trichopoulou A & Naska A (2001) Special Issue: the DAFNE initiative assessment of dietary patterns across Europe using household budget survey data. *Public Health Nutr* 4, 1135–1141.
- EPIC (1992) European Prospective Investigation into Cancer and Nutrition. http://epic.iarc.fr/ (accessed January 2009).
- EFCOSUM (1999) European Food Consumption Survey Method. http://www.public-health.tu-dresden.de/dotnetnuke3/ eu/Projects/PastProjects/EFCOSUM/tabid/338/Default.aspx (accessed January 2009).
- Elmadfa I & Weichselbaum E (editors) (2005) European Nutrition and Health Report 2004. Forum Nutr 58, 1–223.
- The HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. http://www.helenastudy.com/ (accessed September 2008).
- EFCOVAL (2006) European Food Consumption Validation. http://www.efcoval.eu/ (accessed January 2009).
- EURALIM (1996) Coordination and Evaluation of a European Information Campaign on Diet and Nutrition. http://www.epidemiology.ch/euralim/task.html (accessed January 2009).
- EURODIET (1998) Nutrition and Diet for Healthy Lifestyles in Europe. http://eurodiet.med.uoc.gr/ (accessed January 2009).
- EUROFIR (2005) European Food Information Resource Network. http://www.eurofir.net/index.asp?id = 1 (accessed January 2009).
- EURONUT (European Union's Concerted Action on Nutrition and Health) – SENECA (Survey in Europe on Nutrition and the Elderly: a Concerted Action) (1988). http://www.unu.edu/ unupress/food/V183e/ch05.htm (accessed January 2009).
- HECTOR (2006) Healthy Eating Out (Eating out: Habits, Determinants, and Recommendations for Consumers and the European Catering Sector). http://www.nut.uoa.gr/hector/ (accessed January 2009).
- WHO MONICA (1980) Multinational monitoring of trends and determinants in cardiovascular disease). http://www.ktl.fi/ monica/ (accessed January 2009).
- García-Álvarez A, Blanquer M, Ribas-Barba L, et al. (2009)
   How does the quality of surveys for nutrient intake adequacy
   assessment compare across Europe? A scoring system to rate
   the quality of data in such surveys. EURRECA Network of
   Excellence (NoE). Br J Nutr 101, Suppl. 2, S51–S63.

- 23. Elmadfa I & Freisling H (2005) Fat intake, diet variety and health promotion. *Forum Nutr* **57**, 1–10.
- Koenig J & Elmadfa I (1999) Food-based dietary guidelinesthe Austrian perspective. Br J Nutr 81, Suppl. 2, S31–S35.
- Rust P & Elmadfa I (2005) Attitudes of Austrian adults to the consumption of fruits and vegetables. Forum Nutr 57, 91–99.
- Yngve A, Wolf A, Poortvliet E, et al. (2005) Fruit and vegetable intake in a sample of 11-year-old children in 9 European countries: The Pro Children Cross-sectional Survey.
   Ann Nutr Metab 49, 236–245, Epublication 28 July 2005.
- Elmadfa I (2009) Austrian Study on Nutritional Status (ASNS Österreichische Studie zum Ernährungsstatus, ÖSES), Adult population 18–65 years, 2005–2006. (In the Press).
- Bazelmans C, Dramaix M, Kornitzer M, et al. (2005) Application of a dietary pattern approach in the Belgium context. Rev Epidemiol Sante Publique 53, 182–191.
- De Craene I, De Backer G, Kornitzer M, et al. (1989) Determinants of fat consumption in a general population. Arch Belg 47, 98–100.
- De Henauw S, Wilms L, Mertens J, et al. (1997) Overall and meal-specific macronutrient intake in Belgian primary school children. Ann Nutr Metab 41, 89–97.
- Lenaers S, Goffin I, Vinck J, et al. (2006) The nutrition situation of young children in Flanders. Verh K Acad Geneeskd Belg 68, 33–53.
- 32. Matthys C, De Henauw S, Bellemans M, *et al.* (2006) Sources of saturated fatty acids in Belgian adolescents' diet: implications for the development of food-based dietary guidelines. *Br J Nutr* **95**, 546–554.
- 33. Matthys C, De Henauw S, Devos C, *et al.* (2003) Estimated energy intake, macronutrient intake and meal pattern of Flemish adolescents. *Eur J Clin Nutr* **57**, 366–375.
- Pynaert I, Matthys C, Bellemans M, et al. (2005) Iron intake and dietary sources of iron in Flemish adolescents. Eur J Clin Nutr 59, 826–834.
- Sioen IA, Pynaert I, Matthys C, et al. (2006) Dietary intakes and food sources of fatty acids for Belgian women, focused on n-6 and n-3 polyunsaturated fatty acids. Lipids 41, 415-422.
- Staessen L, De Bacquer D, De Henauw S, et al. (1998) Fatty acid composition of the Belgian diet: estimates derived from the Belgian interuniversity research on nutrition and health. Ann Nutr Metab 42, 151–159.
- Staessen L, De Henauw S, De Bacquer D, et al. (1998) Fat sources in the Belgian diet. Ann Nutr Metab 42, 138–150.
- Vereecken CA, Bobelijn K & Maes L (2005) School food policy at primary and secondary schools in Belgium-Flanders: does it influence young people's food habits? *Eur J Clin Nutr* 59, 271–277.
- Wind M, de Bourdeaudhuij I, te Velde SJ, et al. (2006) Correlates of fruit and vegetable consumption among 11-year-old Belgian-Flemish and Dutch schoolchildren. J Nutr Educ Behav 38, 211–221.
- Van Oyen G (2005) Belgian national food consumption survey in the adult population (15 years and older). Federal Agency for Food Safety. http://www.iph.fgov.be/epidemio/epien/ index5.htm
- Ivanova L, Dimitrov P, Ovcharova D, et al. (2006) Economic transition and household food consumption: a study of Bulgaria from 1985 to 2002. Econ Hum Biol 4, 383–397, Epublication 20 September 2006.
- National Nutrition Monitoring of Bulgaria (2009) National Center of Public Health Protection (NCPHP). (In the Press).
- 43. Noah A & Truswell AS (2001) There are many Mediterranean diets. *Asia Pac J Clin Nutr* **10**, 2–9.
- Panagiotakos DB, Zeimbekis A, Boutziouka V, et al. (2007)
   Long-term fish intake is associated with better lipid profile,

- arterial blood pressure, and blood glucose levels in elderly people from Mediterranean islands (MEDIS epidemiological study). *Med Sci Monit* **13**, CR307–CR312.
- Panagiotakos DB & Polychronopoulos E (2006) MEDIS Study – Mediterranean Islands Epidemiological Study. Harokopio: University of Athens.
- Lazarou C (2009) CYFAMILIES (Cyprus Families study) Kyperounda Elementary School. Research Promotion Foundation. (In the Press).
- 47. Lazarou C, Panagiotakos DB, Panagiotou G, et al. (2008) Overweight and obesity in preadolescent children and their parents in Cyprus: prevalence and associated sociodemographic factors – the CYKIDS study. Obes Rev 9, 185–193, Epublication 19 December 2007.
- Dofkova M, Kopriva V, Resova D, et al. (2001) The development of food consumption in the Czech Republic after 1989.
   Public Health Nutr 4, 999–1003.
- Fiala J & Brázdová Z (2000) A comparison between the lifestyles of men and women-parents of school age children. Cent Eur J Public Health 8, 94–100.
- Hainer V, Kunesova M, Bellisle F, et al. (2006) The eating inventory, body adiposity and prevalence of diseases in a quota sample of Czech adults. Int J Obes 30, 830–836.
- Humeníkova L & Gates GE (2007) Dietary intakes, physical activity, and predictors of child obesity among 4–6th graders in the Czech Republic. Cent Eur J Public Health 15, 232–238.
- Kobzová J, Vignerová J, Bláha P, et al. (2004) The 6th nationwide anthropological survey of children and adolescents in the Czech Republic in 2001. Cent Eur J Public Health 12, 126–130.
- Kovárová M, Vignerová J, Bláha P, et al. (2002) Bodily characteristics and lifestyle of Czech children aged 7-00 to 10-99 years, incidence of childhood obesity. Cent Eur J Public Health 10, 169–173.
- Müllerová D, Aujezdská A, Stozický F, et al. (1998)
   Evaluation of diet in a sample of Czech mothers six months after delivery. Cas Lek Cesk 137, 624–627.
- Wagenknecht M, Hainer V, Kunesová M, et al. (2007) Relationships between the 'eating inventory' factors, socioeconomic status, anthropometric body adiposity indexes and health risks in Czech population. Cas Lek Cesk 146, 284–286, 287–291.
- Ruprich J, Dofkova M, Rehurkova I, et al. (2006) Individual Food Consumption – The National Study SISP04. Prague: CHFCH NIPH, available at http://www.chpr.szu.cz/ spotrebapotravin.htm
- 57. Due P, Holstein BE, Ito H, *et al.* (1991) Diet and health behavior in Danish children aged 11–15 years. *Ugeskr Laeger* **153**, 984–988.
- 58. Haraldsdóttir J (1999) Dietary guidelines and patterns of intake in Denmark. *Br J Nutr* **81**, Suppl. 2, S43–S48.
- Haraldsdóttir J, Astrup AV, Dynesen AW, et al. (2002)
   Steadily changing food consumption of Danes. Clear trends during the period 1995–2001. Ugeskr Laeger 164, 2028–2033.
- Haraldsdóttir J, Halkjaer J, Holm L, et al. (1999) Changes in food habits among the population from 1995 to 1998. Ugeskr Laeger 161, 5028–5033.
- Osler M & Heitmann BL (1997) Food patterns associated with intakes of fat, carbohydrate and dietary fibre in a cohort of Danish adults followed for six years. Eur J Clin Nutr 51, 354–361.
- Osler M, Heitmann BL, Gerdes LU, et al. (2001) Dietary patterns and mortality in Danish men and women: a prospective observational study. Br J Nutr 85, 219–225.
- Osler M, Heitmann BL & Schroll M (1997) Ten year trends in the dietary habits of Danish men and women. Cohort and cross-sectional data. Eur J Clin Nutr 51, 535–541.

- Osler M, Milman N & Heitmann BL (1998) Dietary and nondietary factors associated with iron status in a cohort of Danish adults followed for six years. Eur J Clin Nutr 52, 459–463.
- Lyhne N, Christensen T, Groth MV, et al. (2005) Danskernes Kostvaner 2000–2002. The Danish National Survey of Dietary Habits and Physical Activity. Søborg: National Food Institute.
- Villa I, Yngve A, Poortvliet E, et al. (2007) Dietary intake among under-, normal- and overweight 9- and 15-year-old Estonian and Swedish schoolchildren. Public Health Nutr 10, 311–322.
- Pomerleau J, McKee M, Robertson A, et al. (2000) Nutrition and Lifestyle in the Baltic Republics. Estonian Adults Nutrition Survey. London: London School of Hygiene and Tropical Medicine.
- Haapalahti M, Mykkänen H, Tikkanen S, et al. (2003) Meal patterns and food use in 10- to 11-year-old Finnish children. Public Health Nutr 6, 365–370.
- Keski-Rahkonen A, Kaprio J, Rissanen A, et al. (2003) Breakfast skipping and health-compromising behaviours in adolescents and adults. Eur J Clin Nutr 57, 842–853.
- Kosonen H, Rimpelä A, Rauma AL, et al. (2005) Consumption of special diet among Finnish adolescents in 1979–2001: repeated national cross-sectional surveys. Soz Praventivmed 50, 142–150.
- Lyytikäinen A, Lamberg-Allardt C, Kannas L, et al. (2005)
   Food consumption and nutrient intakes with a special focus
   on milk product consumption in early pubertal girls in Central
   Finland. Public Health Nutr 8, 284–289.
- Ovaskainen ML, Reinivuo H, Tapanainen H, et al. (2006) Snacks as an element of energy intake and food consumption. Eur J Clin Nutr 60, 494–501.
- Prättälä R, Helasoja V & Mykkänen H (2001) The consumption of rye bread and white bread as dimensions of health lifestyles in Finland. *Public Health Nutr* 4, 813–819.
- Roos E & Prättälä R (1997) Meal pattern and nutrient intake among adult Finns. Appetite 29, 11–24.
- Ovaskainen ML & Valsta L (2003) The National FINDIET 2002 Study. Helsinki: Publications of the National Public Health Institute B3/2003, available at http://www.ktl.fi/portal/suomi/osastot/ eteo/yksikot/ravitsemusyksikko/julkaisut/finravinto\_2002\_-tutkimuksen\_raportti/ (accessed January 2009).
- National Public Health Institute (2003) The National FINDIET 2002 Study. Helsinki: Publications of the National Public Health Institute B3/2003.
- Bellisle F & Rolland-Cachera MF (2007) Three consecutive (1993, 1995, 1997) surveys of food intake, nutritional attitudes and knowledge, and lifestyle in 1000 French children, aged 9–11 years. J Hum Nutr Diet 20, 241–251.
- Bertrais S, Polo Luque ML, Preziosi P, et al. (2000) Contribution of ready-to-eat cereals to nutrition intakes in French adults and relations with corpulence. Ann Nutr Metab 44, 249–255.
- Boggio V, Grossiord A, Guyon S, et al. (1999) Food consumption of infants and young children in France in 1997. Arch Pediatr 6, 740–747.
- Carrière I, Delcourt C, Lacroux A, et al. (2007) Nutrient intake in an elderly population in southern France (POLANUT): deficiency in some vitamins, minerals and omega-3 PUFA. Int J Vitam Nutr Res 77, 57–65.
- 81. Deheeger M, Rolland-Cachera MF, Pequignot F, *et al.* (1990) Longitudinal study of the feeding of children of 10 months, 2 and 4 years of age. *Arch Fr Pediatr* **47**, 531–537.
- Galan P, Preziosi P, Durlach V, et al. (1997) Dietary magnesium intake in a French adult population. Magnes Res 10, 321–328.
- 83. Kesse E, Boutron-Ruault MC & Clavel-Chapelon F (2005) Regional dietary habits of French women born between

- 1925 and 1950. Eur J Nutr **44**, 285–292, Epublication 30 August 2004.
- 84. Lafay L, Vray M, Boute D, *et al.* (1998) Food and nutritional data for a population from northern France: the Fleurbaix Laventie Ville Santé (FLVS) Study. *Rev Epidemiol Sante Publique* **46**, 263–275.
- Platat C, Perrin AE, Oujaa M, et al. (2006) Diet and physical activity profiles in French preadolescents. Br J Nutr 96, 501 – 507.
- Preziosi P, Galan P, Deheeger M, et al. (1999) Breakfast type, daily nutrient intakes and vitamin and mineral status of French children, adolescents, and adults. J Am Coll Nutr 18, 171–178.
- Touvier M, Lioret S, Vanrullen I, et al. (2006) Vitamin and mineral inadequacy in the French population: estimation and application for the optimization of food fortification. Int J Vitam Nutr Res 76, 343–351.
- Vol S, Doctoriarena A, Le Clésiau H, et al. (1992) Food habits of adults in France. Epidemiological data. Presse Med 21, 1105–1109.
- 89. Castetbon K & Hercberg S (2007) Etude Nationale Nutrition Santé ENNS, 2006. Situation nutritionnelle en France en 2006 selon les indicateurs d'objectif et les repères du Programme national nutrition santé (PNNS). (National Nutrition Health Survey ENNS, 2006. Nutritional Situation in France in 2006 According to Objective Indicators and Benchmarks of the National Nutrition Health (PNNS)). Paris: Institut de veille sanitaire.
- 90. Agence Française de sécurité sanitaire des aliments (2008) INCA 2 Les premiers résultats d'une enquête d'intérêt général à forte valeur ajoutée. Étude Individuelle Nationale des Consommations Alimentaires, 2007. (INCA 2 Preliminary Results of a Survey of Interest in High Value Added. National Study of Food Consumption, 2007). Paris: Agence Française de sécurité sanitaire des aliments.
- Alexy U & Kersting M (2003) Time trends in the consumption of dairy foods in German children and adolescents. Eur J Clin Nutr 57, 1331–1337.
- 92. Alexy U, Kersting M & Sichert-Hellert W (2003) Trends in dietary intake of vitamins A, C, and E in German children and adolescents results of the DONALD Study. *Int J Vitam Nutr Res* **73**, 335–342.
- Beitz R, Mensink GB, Hintzpeter B, et al. (2004) Do users of dietary supplements differ from nonusers in their food consumption? Eur J Epidemiol 19, 335–341.
- 94. Beitz R, Mensink GB, Fischer B, *et al.* (2002) Vitamins dietary intake and intake from dietary supplements in Germany. *Eur J Clin Nutr* **56**, 539–545.
- 95. Heseker H, Hartmann S, Kübler W, *et al.* (1995) An epidemiologic study of food consumption habits in Germany. *Metabolism* **44**, Suppl. 2, 10–13.
- Jungjohann SM, Lührmann PM, Bender R, et al. (2005) Eightyear trends in food, energy and macronutrient intake in a sample of elderly German subjects. Br J Nutr 93, 361–378.
- Kersting M, Sichert-Hellert W, Alexy U, et al. (1998) Macronutrient intake of 1 to 18 year old German children and adolescents. Z Ernahrungswiss 37, 252–259.
- Mensink GB & Beitz R (2004) Food and nutrient intake in East and West Germany, 8 years after the reunification – The German Nutrition Survey 1998. Eur J Clin Nutr 58, 1000–1010.
- Sichert-Hellert W, Kersting M, Alexy U, et al. (2000) Ten-year trends in vitamin and mineral intake from fortified food in German children and adolescents. Eur J Clin Nutr 54, 81–86.
- Thamm M, Mensink GB & Thierfelder W (1999) Folic acid intake of women in childbearing age. Gesundheitswesen 61, 207-212.
- Volkert D, Kreuel K, Heseker H, et al. (2004) Energy and nutrient intake of young-old, old-old and very-old elderly in Germany. Eur J Clin Nutr 58, 1190–1200.

- Volkert D, Kreuel K & Stehle P (2005) Fluid intake of community-living, independent elderly in Germany a nationwide, representative study. *J Nutr Health Aging* 9, 305–309.
- Wolfram N, Hillger C, Jüttler G, et al. (2007) Nutrition in childhood-demands and reality. Outcomes of the nutrition and catering situation in Saxon day-care centers. Dtsch Med Wochenschr 42, 2201–2205.
- Kroke A, Manz F, Kersting M, et al. (2004) The German DONALD Study (Dortmund Nutritional and Anthropometric Longitudinally Designed Study). Eur J Nutr 43, 45–54.
- Arvaniti F, Panagiotakos DB, Pitsavos C, et al. (2006) Dietary habits in a Greek sample of men and women: the ATTICA study. Cent Eur J Public Health 14, 74–77.
- Hassapidou MN & Fotiadou E (2001) Dietary intakes and food habits of adolescents in northern Greece. *Int J Food Sci Nutr* 52, 109–116.
- Manios Y & Kafatos A (2006) Health and nutrition education in primary schools in Crete: 10 years follow-up of serum lipids, physical activity and macronutrient intake. Br J Nutr 95, 568–575.
- Moschandreas JA & Kafatos A (2002) Calcium intake in relation to diet and health indicators in Cretan primary and high school pupils, Greece. Int J Vitam Nutr Res 72, 264–277.
- Moschandreas J & Kafatos A (1999) Food and nutrient intakes of Greek (Cretan) adults. Recent data for food-based dietary guidelines in Greece. Br J Nutr 81, Suppl. 2, S71–S76.
- Roma-Giannikou E, Adamidis D, Gianniou M, et al. (1997)
   Nutritional survey in Greek children: nutrient intake. Eur J Clin Nutr 51, 273–285.
- Trichopoulou A, Kouris-Blazos A, Vassilakou T, et al. (1995)
   Diet and survival of elderly Greeks: a link to the past. Am J Clin Nutr 61, Suppl. 6, S1346-S1350.
- 112. Trichopoulou A, Psaltopoulou T, Orfanos P, et al. (2007) Low-carbohydrate-high-protein diet and long-term survival in a general population cohort. Eur J Clin Nutr 61, 575-581, Epublication 29 November 2006.
- Trichopoulou A, Toupadaki N, Tzonou A, et al. (1993) The macronutrient composition of the Greek diet: estimates derived from six case-control studies. Eur J Clin Nutr 47, 549–558.
- 114. Roma-Giannikou E, Adamidis D, Gianniou M, et al. (1997) First national Greek Nutritional Survey. Nutritional survey in Greek children: nutrient intake. Eur J Clin Nutr 51, 273–285.
- 115. Manios Y (2006) The Growth, Exercise and Nutrition Epidemiological Study in preSchoolers (GENESIS). Design and descriptive results of the 'Growth, Exercise and Nutrition Epidemiological Study in preSchoolers': The GENESIS Study. BioMed Central Public Health 15, 6–32.
- Naska A, Orfanos P, Chloptios I, et al. (2005) Greek component dietary habits in Greece: The European Prospective Investigation into Cancer and nutrition (the EPIC project).
   Arch Hellenic Med 22, 259–269.
- Arvaniti F, Panagiotakos DB, Pitsavos C, et al. (2006) Dietary habits in a Greek sample of men and women: The ATTICA Study. Cent Eur J Publ Health 14, 74–78.
- Hassapidou MN, Fotiadou E, Maglara E, et al. (2006) Energy intake, diet composition, energy expenditure and obesity of adolescents in northern Greece. Obesity 14, 855–862.
- Antal M, Nagy K, Bíró L, et al. (2003) Cross-sectional survey on the nutritional and life-style habits of secondary school students in Hungary. Orv Hetil 144, 1631–1636.
- 120. Biró G, Antal M & Zajkás G (1996) Nutrition survey of the Hungarian population in a randomized trial between 1992–1994. Eur J Clin Nutr 50, 201–208.
- Biró L, Zajkás G, Greiner E, et al. (2007) Hungarian National Dietary survey, 2003–2004. Micronutrients: mineral salts. Orv Hetil 148, 703–708.
- Rodler I, Bíró L, Greiner E, et al. (2005) Dietary survey in Hungary, 2003–2004. Orv Hetil 146, 1781–1789.

- 123. Rurik I (2004) Evaluation on lifestyle and nutrition among Hungarian elderly. *Z Gerontol Geriatr* **37**, 33–36.
- Rurik I (2006) Nutritional differences between elderly men and women. Primary care evaluation in Hungary. *Ann Nutr Metab* 50, 45–50, Epublication 7 November 2005.
- 125. 3rd National Nutrition Survey (a part of the National Health Survey of the Population; in Hungarian: OLEF). 3rd paper on the results (vitamins) of this study in 2007. (2007) *Hungarian Med J (Orvosi Hetilap)* 148, 1593–1600.
- Rurik I (2006) Nutritional differences between elderly men and women. Ann Nutr Metab 50, 45–50.
- Rurik I (2004) Evaluation on lifestyle and nurition among Hungarian elderly. Z Geront Geriat 37, 33–36.
- 128. Bíró G, Antal M & Zajkás G (1996) Nutrition survey of the Hungarian population in a randomized trial between 1992–1994. Eur J Clin Nutr 50, 201–208.
- Péter S, Regöly-Mérei A, Bíró L, et al. (2007) Lifestyle of school children: representative survey in Metropolitan elementary Schools – Part One. Ann Nutr Metab 51, 448–453.
- Gunnarsson BS, Thorsdottir I & Palsson G (2004) Iron status in 2-year-old Icelandic children and associations with dietary intake and growth. Eur J Clin Nutr 58, 901–906.
- Thórsdóttir I & Gunnarsdóttir I (2005) Vitamin D in nutrition of young Icelandic children. *Laeknabladidb* 91, 581–586.
- 132. Wolf A, Yngve A, Elmadfa I, et al. (2005) Fruit and vegetable intake of mothers of 11-year-old children in nine European countries: The Pro Children Cross-sectional Survey. Ann Nutr Metab 49, 246–254, Epublication 28 July 2005.
- 133. Yngve A, Wolf A, Poortvliet E, *et al.* (2005) Fruit and vegetable intake in a sample of 11-year-old children in 9 European countries: The Pro Children Cross-sectional Survey. *Ann Nutr Metab* **49**, 236–245, Epublication 28 July 2005.
- 134. Steingrímsdóttir L, Þorgeirsdóttir H & Ólafsdóttir AS (2002) The Diet of Icelanders, Dietary Survey of The Icelandic Nutriton Council 2002: PHI Luxembourg, available at http://www. lydheilsustod.is/rannsoknir/matur-mataraedi-holdafar/kannanir-a-mataraedi/nr/302; http://www.lydheilsustod.is/media/manneldi/rannsoknir/skyrsla.pdf (accessed January 2009).
- Cosgrove M, Flynn A & Kiely M (2005) Consumption of red meat, white meat and processed meat in Irish adults in relation to dietary quality. *Br J Nutr* 93, 933–942.
- Gabhainn SN, Nolan G, Kelleher C, et al. (2005) Dieting patterns and related lifestyles of school-aged children in the Republic of Ireland. Public Health Nutr 5, 457–462.
- Galvin MA, Kiely M, Harrington KE, et al. (2001) The North/ South Ireland Food Consumption Survey: the dietary fibre intake of Irish adults. Public Health Nutr 4, 1061–1068.
- 138. Hannon EM, Kiely M, Harrington KE, et al. (2001) The North/ South Ireland Food Consumption Survey: mineral intakes in 18–64-year-old adults. Public Health Nutr 4, 1081–1088.
- Harrington KE, McGowan MJ, Kiely M, et al. (2001) Macronutrient intakes and food sources in Irish adults: findings of the North/South Ireland Food Consumption Survey. Public Health Nutr 4, 1051–1060.
- 140. Harrington KE, Robson PJ, Kiely M, et al. (2001) The North/South Ireland Food Consumption Survey: survey design and methodology. Public Health Nutr 4, 1037–1042.
- Hurson M, Corish C & Sugrue S (1997) Dietary intakes in Ireland of a healthy elderly population. Ir J Med Sci 166, 220–224.
- Hill TR, O'brien MM, Cashman KD, et al. (2004) Vitamin D intakes in 18-64-y-old Irish adults. Eur J Clin Nutr 58, 1509-1517.
- Kiely M, Flynn A, Harrington KE, et al. (2001) Sampling description and procedures used to conduct the North/South Ireland Food Consumption Survey. Public Health Nutr 4, 1029–1035.

- McNulty H, Eaton-Evans J, Cran G, et al. (1996) Nutrient intakes and impact of fortified breakfast cereals in schoolchildren. Arch Dis Child 75, 474–481.
- 145. O'Brien MM, Kiely M, Harrington KE, et al. (2001) The North/South Ireland Food Consumption Survey: vitamin intakes in18–64-year-old adults. Public Health Nutr 4, 1069–1079.
- O'Brien MM, Kiely M, Galvin M, et al. (2003) The importance of composite foods for estimates of vegetable and fruit intakes. Public Health Nutr 6, 711–726.
- Strain JJ, Robson PJ, Livingstone MB, et al. (1994) Estimates of food and macronutrient intake in a random sample of Northern Ireland adolescents. Br J Nutr 72, 343–352.
- Villegas R, Salim A, Collins MM, et al. (2004) Dietary patterns in middle-aged Irish men and women defined by cluster analysis. Public Health Nutr 7, 1017–1024.
- 149. National Nutrition Surveillance Centre (2003) Dietary Habits of the Irish Population: Results from SLÁN. Annual Report. Dublin: National Nutrition Surveillance Centre, available at http://www.healthpromotion.ie/research (accessed January 2009).
- 150. Koski Marjaana L & Mervi S (2004) Nutrition Report 2003. Publications of National Public Health Institute B4/2004. http://www.ktl.fi/portal/english/public\_health\_monitoring\_promotion/monitoring\_interventions/nutrition\_in\_finland/finnish\_national\_nutrition\_surveillance\_system/nutrition\_report\_2003/
- Higgins M (2001) North/South Ireland Food Consumption Survey – by IUNA (Irish Universities Nutrition Alliance).
   Dublin: Food Safety Promotion Board.
- Bellù R, Riva E, Ortisi MT, et al. (1995) Calcium intakes in a sample of 35,000 Italian schoolchildren. J Int Med Res 23, 191–199.
- 153. Corrêa Leite ML, Nicolosi A, Cristina S, et al. (2003) Dietary and nutritional patterns in an elderly rural population in Northern and Southern Italy: (I). A cluster analysis of food consumption. Eur J Clin Nutr 57, 1514–1521.
- 154. Corrêa Leite ML, Nicolosi A, Cristina S, *et al.* (2003) Dietary and nutritional patterns in an elderly rural population in Northern and Southern Italy: (II). Nutritional profiles associated with food behaviours. *Eur J Clin Nutr* **57**, 1522–1529.
- 155. Inelmen EM, Gimenez GF, Gatto MR, et al. (2000) Dietary intake and nutritional status in Italian elderly subjects. J Nutr Health Aging 4, 91–101.
- Lombardi-Boccia G, Aguzzi A, Cappelloni M, et al. (2003)
   Total-diet study: dietary intakes of macro elements and trace elements in Italy. Br J Nutr 90, 1117–1121.
- Piffer S, Kaisermann D & Pasquazzo MT (2003) Dietary habits in primary school children. Role of school meals and parents's social class. *Ann Ig* 15, 1097–1098.
- Turrini A, Saba A, Perrone D, et al. (2001) Food consumption patterns in Italy: the INN-CA Study 1994–1996. Eur J Clin Nutr 55, 571–588.
- Welch AA, Lund E, Amiano P, et al. (2002) Variability of fish consumption within the 10 European countries participating in the European Investigation into Cancer and Nutrition (EPIC) Study. Public Health Nutr 5, 1273–1285.
- Turrini A & Leclercq C (2006) Food Consumption Patterns in Italy, INRAN-SCAI 2005–2006. Rome: Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione.
- 161. Turrini A, Sap A, Operone D, et al. (2001) Food Consumption Patterns in Italy: The INN-CA 1994–1996. Rome: Istituto Nazionale di Recerca per gli Alimenti e la Nutrizione, Istituto Nazionale di Statistica.
- 162. Petkeviciene J, Klumbiene J, Prättälä R, et al. (2007) Educational variations in the consumption of foods containing fat in Finland and the Baltic countries. Public Health Nutr 10, 518–523.

- Pomerleau J, McKee M, Robertson A, et al. (2001) Macronutrient and food intake in the Baltic republics. Eur J Clin Nutr 55, 200–207.
- 164. Joffe R (2009) National Food Consumption Survey. Latvia: Food Centre of Food and Veterinary Service of Latvia.
- Abaravicius JA, Kucinskiene Z, Songaliene J, et al. (1998)
   Study of nutritional behavior in a group of Lithuanian men.
   Z Ernahrungswiss 37, Suppl. 1, 136–138.
- Grabauskas V, Petkeviciene J, Kriaucioniene V, et al. (2004)
   Health inequalities in Lithuania: education and nutrition habits. Medicina 40, 875–883.
- Grabauskas V, Zaborskis A, Klumbiene J, et al. (2004)
   Changes in health behavior of Lithuanian adolescents and adults over 1994–2002. Medicina 40, 884–890.
- 168. Kasmel A, Helasoja V, Lipand A, et al. (2004) Association between health behaviour and self-reported health in Estonia, Finland, Latvia and Lithuania. Eur J Public Health 14, 32–36.
- Kriaucioniene V, Petkeviciene J & Klumbiene J (2008) Dietary patterns and their association with sociodemographic factors in Lithuanian adult population. *Medicina (Kaunas)* 44, 799–804.
- Puska P, Helasoja V, Prättälä R, et al. (2003) Health behaviour in Estonia, Finland and Lithuania 1994–1998. Standardized comparison. Eur J Public Health 13, 11–17.
- Skemiene L, Ustinaviciene R, Piesine L, et al. (2007) Peculiarities of medical students' nutrition. Medicina 43, 145–152.
- 172. Barzda A (2008) Macronutrient, Food Intake & Health Behaviour in Lithuanian Population. Vilnius: National Nutrition Center Lithuanian Ministry of Health.
- Guillaume M, Lapidus L & Lambert A (1998) Obesity and nutrition in children. The Belgian Luxembourg Child Study IV. Eur J Clin Nutr 52, 323–328.
- Paulus D, Saint-Remy A & Jeanjean M (2001) Dietary habits during adolescence – results of the Belgian Adolux Study. Eur J Clin Nutr 55, 130–136.
- Noah A & Truswell AS (2001) There are many Mediterranean diets. Asia Pac J Clin Nutr 10, 2–9.
- Tessier S & Gerber M (2005) Factors determining the nutrition transition in two Mediterranean islands: Sardinia and Malta. *Public Health Nutr* 8, 1286–1292.
- 177. Pace AR (2003) *The First National Health Interview Survey* Malta: Department of Health and Research, available at http://www.sahha.gov.mt/pages.aspx?page=383 (accessed January 2009).
- 178. Feunekes GI, Stafleu A, de Graaf C, et al. (1997) Family resemblance in fat intake in The Netherlands. Eur J Clin Nutr 51, 793–799.
- 179. Giskes K, Lenthe Fv F, Brug HJ, *et al.* (2004) Dietary intakes of adults in the Netherlands by childhood and adulthood socioeconomic position. *Eur J Clin Nutr* **58**, 871–880.
- Hulshof KF, Brussaard JH, Kruizinga AG, et al. (2003) Socioeconomic status, dietary intake and 10 y trends: the Dutch National Food Consumption Survey. Eur J Clin Nutr 57, 128–137
- Hulshof KF, Löwik MR, Kok FJ, et al. (1991) Diet and other life-style factors in high and low socio-economic groups (Dutch Nutrition Surveillance System). Eur J Clin Nutr 45, 441–450.
- Hulshof KF, Wedel M, Löwik MR, et al. (1992) Clustering of dietary variables and other lifestyle factors (Dutch Nutritional Surveillance System). J Epidemiol Community Health 46, 417–424.
- 183. Kromhout D, de Lezenne Coulander C, Obermann-de Boer GL, et al. (1990) Changes in food and nutrient intake in middle-aged men from 1960 to 1985 (the Zutphen Study). Am J Clin Nutr 51, 123–129.
- Löwik MR, Hulshof KF & Brussaard JH (1999) Patterns of food and nutrient intakes of Dutch adults according to intakes

- of total fat, saturated fatty acids, dietary fibre, and of fruit and vegetables. *Br J Nutr* **81**, Suppl. 2, S91–S98.
- Schuit AJ, van Loon AJ, Tijhuis M, et al. (2002) Clustering of lifestyle risk factors in a general adult population. Prev Med 35, 219–224.
- Waijers P & Feskens E (2004) Assessment of vitamin A intake of the Dutch population. http://www.rivm.nl/bibliotheek/ rapporten/350010002.html
- 187. Haveman-Nies A, Bokje E, Ocke M, et al. (2002) MONICA Optional Study on Nutrition: the dietary assessment methodology. http://www.rivm.nl/bibliotheek/ rapporten/261753001.html
- 188. Ocke MC, van Rossum CTM, Fransen HP, et al. (2008) Dutch National Food Consumption Survey – Young Children 2005/ 2006. RIVM, Report 350070001. http://www.rivm.nl/ bibliotheek/rapporten/350070001.pdf
- Andersen LF, Nes M, Sandstad B, et al. (1995) Dietary intake among Norwegian adolescents. Eur J Clin Nutr 49, 555–564.
- Andersen LF, Øverby N & Lillegaard IT (2004) Intake of fruit and vegetables among Norwegian children and adolescents. *Tidsskr Nor Laegeforen* 124, 1396–1398.
- Johansson LR, Solvoll K, Bjørneboe GE, et al. (1998) Intake of very-long-chain n-3 fatty acids related to social status and lifestyle. Eur J Clin Nutr 52, 716–721.
- Moreiras O, van Staveren WA, Cruz JA, et al. (1991) Intake of energy and nutrients. Euronut SENECA investigators. Eur J Clin Nutr 45, Suppl. 3, 105–119.
- Wandel M (1995) Dietary intake of fruits and vegetables in Norway: influence of life phase and socio-economic factors. Int J Food Sci Nutr 46, 291–301.
- 194. Lande B, Andersen LF, Bærug A, et al. (2003) Infant feeding practices and associated factors in the first six months of life: The Norwegian Infant Nutrition Survey. Acta Pædiatr 92, 152–161
- Lande B, Andersen LF, Veierød MB, et al. (2004) Breastfeeding at 12 months of age and dietary habits among breast-fed and non-breast-fed infants. Public Health Nutr 7, 495–503.
- Lande B (2005) Spedkost 24 months. http://www.shdir.no/vp/multimedia/archive/00006/IS-1299 6113a.pdf
- Andersen LF, Nes M, Sandstad B, et al. (1995) Dietary intake among Norwegian adolescents. Eur J Clin Nutr 49, 555–564.
- Andersen LF, Nes M, Bjørneboe G, et al. (1997) Food habits among 13-year old Norwegian adolescents. Scand J Nutr 41, 150–154.
- Johansson L, Solvoll K, Bjørneboe EA, et al. (1997) Dietary habits among Norwegian men and women Scand. J Nutr 47, 63-70.
- Johansson L (1997) Surveillance of dietary intake in the Norwegian adult population NORKOST 2. http://www.shdir.no/vp/multimedia/archive/00003/IS-0168\_3745a.pdf
- Augustyniak U & Brzozowska A (2002) Nutrients intake of the adolescents in Poland – ten year's literature review (1990–2000). Rocz Panstw Zakl Hig 53, 399–406.
- Biela U & Pajak A (2005) Intake of selected nutrients by perceived control over life, over health and general control in men and women aged 45–64, residents of Kraków. *Przegl Lek* 62, 1367–1371.
- Kolarzyk E, Ostachowska-Gasior A & Skop A (2005) The protein participation in daily diet and nutritional status of medical students in Kraków. Rocz Akad Med Bialymst 50, Suppl. 1, 39–41.
- Pardo B, Sygnowska E, Rywik S, et al. (1991) Dietary habits of the middle-aged Warsaw population in 1984 relative to nutritional guidelines. Appetite 16, 1–15.
- Suliga E (2006) Nutritional status and dietary habits of urban and rural Polish adolescents. Anthropol Anz 64, 399–409.

- Szewczyński JA & Ostrowska A (2002) Changes of dietary habits among Warsaw Medical University students in the years 1985–2001. Wiad Lek 55, Suppl. 1, 489–495.
- Szponar L & Rychlik E (2002) Dietary intake elderly subjects in rural and urban area in Poland. *Pol Merkur Lekarski* 13, 490–496.
- Weker H (2000) Evaluation of the nutritional status and feeding patterns of 3-7 year old children - results of a questionnaire. Med Wieku Rozwoj 4, Suppl. 1, 41-52.
- Wierzbicka E, Brzozowska A & Roszkowski W (2001) Energy and nutrients intake of elderly people living in the Warsaw region, Poland. J Nutr Health Aging 5, 248–252.
- 210. Szponar L, Sekula W, Rychlik E, et al. (2003) Household food consumption and anthropometric survey. Publication of National Food and Nutrition Institute no. 101. Warsaw: National Food and Nutrition Institute.
- Broda G (2005) National Health Survey in Poland WOBASZ Project. Warsaw: Department of Human Nutrition, Warsaw Agricultural University.
- Charzewska J (2006) Nutritional Survey Dietary Habits, Nutrient Intake and Nutritional Status were Assessed.
   Warsaw: National Food and Nutrition Institute.
- 213. Charzewska J (2002) Assessment of Nutritional Status of Postmenopausal Women and Dietary Habits of Warsaw Postmenopausal Women Aged 60, 70 Years. Warsaw: National Food and Nutrition Institute.
- Marques-Vidal P, Ravasco P, Dias C, et al. (2006) Trends of food intake in Portugal, 1987–1999: results from the National Health Surveys. Eur J Clin Nutr 60, 1414–1422, Epublication 21 June 2006.
- Moreira P, Padez C, Mourão I, et al. (2005) Dietary calcium and body mass index in Portuguese children. Eur J Clin Nutr 59, 861–867.
- 216. Lopes C, Oliveira A, Santos AC, et al. (2006) Consumo alimentar no Porto. (Food Consumption in Porto). Faculdade de Medicina do Porto, available at http://www. consumoalimentarporto.med.up.pt
- Barros H (2007) EPITeen Cohort Study. Porto: Department of Hygiene and Epidemiology, University of Porto Medical School.
- Barros R, Moreira A, Fonseca J, et al. (2008) Adherence to the Mediterranean diet and fresh fruit intake are associated with improved asthma control. Allergy 63, 917–923.
- Graça MJ, Dias AL, Rodrigues CM, et al. (2001) Inquérito nacional de saúde – Dados Gerais – Continente 98/99. (National Health Survey – General Data – Mainland 98/ 99). Lisboa: Instituto Nacional de Saúde Dr. Ricardo Jorge.
- Albu A, Indrei LL, Gavăt V, et al. (2005) Specifics of food intake by schoolchildren in the rural areas of Moldova. Rev Med Chir Soc Med Nat Iasi 109, 153–157.
- Petrescu C, Avram CL, Suciu OV, et al. (2005) Nutrition in preschool children from Băile-Herculane. Rev Med Chir Soc Med Nat Iasi 109, 365–372.
- Petrescu C, Buleandră CF, Doroftei S, et al. (2004) Nutrients in food intake of medical students, Timişoara. Rev Med Chir Soc Med Nat Iasi 108, 338–342.
- Zapîrtan H, Ionuţ C, Vlad M, et al. (2006) Dietary imbalance in some communities of Transylvania. Rev Med Chir Soc Med Nat Iasi 110, 993–998.
- 224. Albu A, Prejbeanu I, Gavat I, et al. (2004) Aspects of children's nutritional habits in two rural communities. Rev Med Chir Soc Med Nat Iasi 108, 343-347.
- Vlad M (2008) Aging and Nutrition Project, Nutritional status in elderly-Romania. Ann Nutr Metab 52, Suppl. 1, 62–71.
- Pavlović M, Grujić V & Oshaug A (2005) Nutrition and physical activity of the population in Serbia. World Rev Nutr Diet 94, 51–59.

- Ministry of Health Republic of Serbia (2007) National Health Survey 2006. Belgrade: Ministry of Health Republic of Serbia.
- Pavlović M, Majkić SN, Kadvan A, et al. (2001) Nutrition as the risk factor for atherosclerosis. Jugoslov Med Biohem 20, 107–115.
- 229. Pavlović M, Kadvan A, Vukotić M, et al. (2006) The quality of nutrition in schoolchildren and adult members of families. In Yugoslav Study of Atherosclerosis Precursors in Schoolchildren in Serbia from 1998–2003, Monograph 262–285 [S Nedeljkovic, editor]. Belgrade: CIBID Press.
- Pavlović M & Kadvan A (2006) Methodology of the dietary survey in YUSAD study. In Yugoslav Study of Atherosclerosis in Schoolchildren in Serbia from 1998–2003, Monograph 254–261 [S Nedeljkovic, editor]. Belgrade: CIBID Press.
- 231. Pavlović M, Kadvan A & Vukotić M (2005) The quality of schoolchildren's nutrition in Serbia. [B Koletzko , P Dodds, H Akerblom and M Ashwell, editors]. In Early Nutrition and it's Later Consequences: New Opportunities, Perinatal Programming Of Adult Health-EC Supported Research, Advances in Experimental Medicine and Biology: Springer Science + Business Media New York vol. 569, pp. 212–213.
- Stefániková Z, Jurkovicová J, Sevcíková L, et al. (2001) Did the nutrition of Slovak medical students change? A comparison after 15 years. Cent Eur J Public Health 9, 223–227.
- Stefanikova Z, Sevcikova L, Jurkovicova J, et al. (2006) Positive and negative trends in university students' food intake. Bratisl Lek Listy 107, 217–220.
- Koch V (1999) Nutritional Habits of Slovenian Adults in Health Protection Aspect. Ljubljana: University of Ljubljana.
- Aranceta J, Pérez Rodrigo C, Eguileor I, et al. (1998) Food consumption patterns in the adult population of the Basque Country (EINUT-I). Public Health Nutr 1, 185–192.
- Aranceta J, Serra-Majem L, Pérez-Rodrigo C, et al. (2001)
   Vitamins in Spanish food patterns: the eVe Study. Public Health Nutr 4, 1317–1323.
- 237. Aroca MD, Menárguez JF, Luna C, *et al.* (1997) Dietary habits and consumption patterns in a health district. *Aten Primaria* **19**, 72–79.
- 238. de Rufino PM, Muñoz P & Gómez E (1999) Frequency of food intake by adolescent schoolchildren in Cantabria, Spain. Comparison with the consensus document: food guidelines for the Spanish population. *Gac Sanit* 13, 449–455.
- Estévez MD, Doreste JL, Henríquez P, et al. (1994) Food habits among children in a district of the island of Gran Canaria. Aten Primaria 14, 1141–1147.
- López-Frías M, Nestares T, Iañez I, et al. (2005) Nutrient intake adequacy in schoolchildren from a Mediterranean area (southern Spain). Influence of the use of the school canteen. Int J Vitam Nutr Res 75, 312–319.
- Maldonado R & Villalbí JR (1993) Food preferences among students. An Esp Pediatr 39, 10–14.
- Morales O, Ochoa JJ, López-Frías M, et al. (2007) Vitamin intake in the adult population of the Andalusian Autonomous Community. Nutr Hosp 22, 108–111.
- 243. Royo-Bordonada MA, Gorgojo L, de Oya M, et al. (2003) Food sources of nutrients in the diet of Spanish children: the Four Provinces Study. Br J Nutr 89, 105–114.
- Royo-Bordonada MA, Gorgojo L, Martín-Moreno JM, et al. (2003) Investigators of the Four Provinces Study. Spanish children's diet: compliance with nutrient and food intake guidelines. Eur J Clin Nutr 57, 930–939.
- Ruano I & Serra ME (1997) Eating and life style habits in primary school student population from Mataro (Barcelona) associated with consumption of sugar and TV watching. *Rev Esp Salud Publica* 71, 487–498.
- Serra-Majem L, Ribas Barba L, Pérez-Rodrigo C, et al. (2002)
   Determinants of nutrient intake among children and adoles-

- cents: results from the enKid Study. *Ann Nutr Metab* **46**, Suppl. 1, S31–S38.
- 247. Tur JA, Romaguera D & Pons A (2004) Food consumption patterns in a mediterranean region: does the mediterranean diet still exist? *Ann Nutr Metab* 48, 193–201, Epublication 12 July 2004.
- Serra-Majem L, Ribas Barba L, Pérez Rodrigo C, et al. (2003)
   Dietary habits and food consumption in Spanish children and adolescents (1998–2000): socioeconomic and demographic factors. Med Clin 121, 126–131.
- Serra-Majem L, Ribas Barba L, Pérez Rodrigo C, et al. (2006) Nutrient adequacy in Spanish children and adolescents. Br J Nutr 96, Suppl. 1, S49–S57.
- Serra-Majem L & Aranceta J (editors) Estudio EnKid (2000).
   vol. 1: Desayuno y equilibrio alimentario. (2001) vol. 2: Obesidad infantil y juvenil. (2002) vol. 3: Alimentación infantil y juvenil. (2003) vol. 4: Crecimiento y desarrollo. Krece Plus. (2004) vol. 5: Nutrición infantil y juvenil. (2006) vol. 6: Actividad física y salud (vol. 1: Breakfast and Balanced Diet. (2001), vol. 2: Obese Children and Adolescents. (2002), vol. 3: Feeding Children and Adolescents. (2003), vol. 4: Growth and Development. Kreća Plus. (2004), vol. 5: Child and Adolescent Nutrition. (2006), vol. 6: Physical Activity and Health).
- Serra Majem L & Ribas Barba L (editors) (2007) Trends in nutritional status in Catalonia, Spain (1992–2003). Department of Health, Generalitat of Catalonia. 10, 1339–1414, available at http://www.gencat.net/salut/depsalut/pdf/encat 2003semi.pdf (accessed June 2008).
- Serra Majem L (editor) (1999-2000) Encuesta Nutricional de Canarias. 1997-98. Santa Cruz de Tenerife. Servicio Canario de Salud. (1999) (Nutrition Survey of the Canary Islands. 1997-98. Santa Cruz de Tenerife. Canary Health. (1999)). vol. 1: Hábitos alimentarios y consumo de alimentos. (1999), vol. 2: Factores de riesgo cardiovascular. (2000), vol. 3: Ingesta de energía, nutrientes y riesgo de ingestas inadecuadas. (2000), vol. 4: Evaluación bioquímica del estado nutricional. (2000), vol. 5: Conocimientos, opiniones y actitudes, vol. 5: Política de nutrición en Canarias (vol.1: Food habits and Food Consumption. (1999), vol. 2: Cardiovascular Risk Factors. (2000), vol. 3: Intake of Energy, Nutrients and Risk of Inadequate Intakes. (2000), vol. 4: Assessment of Nutritional Biochemistry. (2000), vol. 5: Knowledge, Opinions and Attitudes, vol. 5: Nutrition Policy in the Canaries), http://www.gobiernodecanarias. org/sanidad/scs/1/plansalud/enca/ppal\_enca.htm
- Serra Majem L (2000) Evaluación del estado nutricional de la población canaria (1997–98) (Assessment of nutritional status of the population canaria (1997–98)). Arch Latinoamer Nutr 50, Suppl. 1, S1–S70.
- 254. Vioque J & Quiles J (2003) Encuesta de Nutrición y Salud de la Comunidad Valenciana. (Survey of Nutrition and Health, Valencia). Alicante: Departamento Salud Pública, Universidad Miguel Hernández, available at http://www.dsp.umh.es/ docencia/medprev/encuesta\_cv.pdf
- Requejo AM, Ortega RM & Rivas T (1994) Estado nutritivo en colectivos escolares madrileños. (Nutritional Status in School Collectives Madrileños). Madrid: Ayuntamiento de Madrid.
- 256. Ortega RM, Rodríguez E & López AM (2007) El control de peso corporal como prioridad sanitaria. Resultados de un estudio de intervención encaminado a aproximar la dieta al ideal teórico. In *Nutrición y alimentación en la promoción de la salud* pp. 279–293, [RM Ortega, AM Requejo and RM Martínez, editors]. Cuenca: Universidad Internacional Menéndez y Pelayo.
- Tur JA (2002) Nutritional Survey of the Balearic Islands (ENIB, 1999–2000). Revista de Ciència 27–30.

- Bergström E, Hernell O & Persson LA (1993) Dietary changes in Swedish adolescents. Acta Paediatr 82, 472–480.
- Eiben G, Andersson CS, Rothenberg E, et al. (2004) Secular trends in diet among elderly Swedes – cohort comparisons over three decades. Public Health Nutr 7, 637–644.
- Höglund D, Samuelson G & Mark A (1998) Food habits in Swedish adolescents in relation to socioeconomic conditions. Eur J Clin Nutr 52, 784–789.
- Nydhal M, Andersson J, Sidenvall B, et al. (2003) Food and nutrient intake in a group of self-managing elderly Swedish women. J Nutr Health Aging 7, 67–74.
- Rothenberg E, Bosaeus I & Steen B (1993) Intake of energy, nutrients and food items in an urban elderly population. Aging (Milano) 5, 105–116.
- Samuelson G, Bratteby LE, Berggren K, et al. (1996) Dietary iron intake and iron status in adolescents. Acta Paediatr 85, 1033–1038
- Samuelson G, Bratteby LE, Enghardt H, et al. (1996) Food habits and energy and nutrient intake in Swedish adolescents approaching the year 2000. Acta Paediat Suppl 415, S1-S19, Erratum in: Acta Paediatr Suppl 85, S1392.
- Sjöberg A, Hallberg L, Höglund D, et al. (2003) Meal pattern, food choice, nutrient intake and lifestyle factors in The Göteborg. Adolescence Study. Eur J Clin Nutr 57, 1569–1578.
- 266. Villa I, Yngve A, Poortvliet E, et al. (2007) Dietary intake among under normal- and overweight 9- and 15-year-old Estonian and Swedish schoolchildren. Public Health Nutr 10, 311–322.
- Von Post-Skagegård M, Samuelson G, Karlström B, et al. (2002) Changes in food habits in healthy Swedish adolescents during the transition from adolescence to adulthood. Eur J Clin Nutr 56, 532–538.
- Wolf A, Yngve A, Elmadfa I, et al. (2005) Fruit and vegetable intake of mothers of 11-year-old children in nine European countries: The Pro Children Cross-sectional Survey. Ann Nutr Metab 49, 246–254, Epublication 28 July 2005.
- 269. Becker W (1998) Riskmaten: 1997–1998. Svenskarna äter nyttigare-allt fler väljer grönt. Rismaten 1997–1998. The Swedes eat more healthy (Risk Maten:1997–1998. Swedes eat healthier, more and more choose green. Rismaten 1997–1998. The Swedes eat more healthy). Var Föda 51, 24–27.
- Eiben G, Andersson CS, Rothenberg E, et al. (2004) Secular trends in diet among elderly Swedes cohort comparisons over three decades. Public Health Nutr 7, 637–644.
- 271. Becker W (2002) Assessmeent of food consumption and energy and nutrient intake of adult population in Sweden, Riksmaten 1997–98. http://www.slv.se/templates/SLV\_Page. aspx?id = 12429&epslanguage=SV
- 272. Enghardt H (2006) Assessmeent of food consumption and energy and nutrient intake of children population in Sweden, Riksmaten children 2003. http://www.slv.se/ templates/SLV\_Page.aspx?id=12488&epslanguage=SV
- Beer-Borst S & Amadò R (1996) Dietary survey 'HEUREKA'
   1991: dietary intake of a Swiss collective assessed by a self-administered 24-hour recall questionnaire. Soz Praventivmed
   41, 295–302.
- Cavadini C, Decarli B, Dirren H, et al. (1999) Assessment of adolescent food habits in Switzerland. Appetite 32, 97–106.
- Decarli B, Cavadini C, Grin J, et al. (2000) Food and nutrient intakes in a group of 11 to 16 year old Swiss teenagers. Int J Vitam Nutr Res 70, 39–147.
- Eichholzer M & Bisig B (2000) Daily consumption of (red) meat or meat products in Switzerland: results of the 1992/93 Swiss Health Survey. Eur J Clin Nutr 54, 136–342.

- Eichholzer M & Bisig B (1999) Inadequate fruit and vegetable consumption in Switzerland: results of the Swiss Health Survey 1992/93. Soz Praventivmed 44, 143–151.
- 278. Beer-Borst S, Costanza M & Pechère-Bertschi A (2006)
  Calibration of Geneva adult population dietary sodium (salt)
  intake. Available at http://www.bag.admin.ch/suchen/index.
  html?keywords=food + habits&go\_search=Suchen&lang=
  de#volltextsuche
- 279. Beer M (2006) NANUSS (National Nutrition Survey Switzerland). Bundesamt für Gesundheit BAG (Federal Office of Public Health BAG). Available at http://www.bag.admin.ch/themen/ernaehrung\_bewegung/05190/05297/index.html?lang=de (accessed January 2009).
- 280. Morabia A (2002) Nutritional and Physical Health Survey. Le Bus Santé 2000, instrument de l'observatoire épidémiologique genevois. (Bus Health 2000, Instrument of the Observatory Epidemiological Genevois). Geneva: Médicine et Hygiène.
- Beer-Borst SBUS Santé Surveys (unite mobile du service d'epidemiologie clinique – Observatoire Epidemiologique).
   (BUS Health Surveys (Service Mobile Unit of Clinical Epidemiology – Epidemiologique Observatory 2008)). Available at www.bus-sante.ch/ (accessed January 2009).
- Billson H, Pryer JA & Nichols R (1999) Variation in fruit and vegetable consumption among adults in Britain. An analysis from the dietary and nutritional survey of British adults. *Eur J Clin Nutr* 53, 946–952.
- 283. Cowin I & Emmett P (2007) Diet in a group of 18-month-old children in South West England, and comparison with the results of a national survey. *J Hum Nutr Diet* **20**, 254–267.
- 284. Crozier SR, Robinson SM, Borland SE, *et al.* (2006) Dietary patterns in the Southampton Women's Survey. *Eur J Clin Nutr* **60**, 1391–1399, Epublication 28 June 2006.
- 285. Emmett P, Rogers I & Symes C (2002) Avon Longitudinal Study of Pregnancy and Childhood. Food and nutrient intakes of a population sample of 3-year-old children in the south west of England in 1996. *Public Health Nutr* 5, 55-64.
- Gibson SA & O'Sullivan KR (1995) Breakfast cereal consumption patterns and nutrient intakes of British schoolchildren. J R Soc Health 115, 366–370.
- 287. Hackett AF, Gibbon M, Sratton G, *et al.* (2002) Dietary intake of 9–10-year-old and 11–12-year-old children in Liverpool. *Public Health Nutr* **5**, 449–455.
- Inchley J, Todd J, Bryce C, et al. (2001) Dietary trends among Scottish schoolchildren in the 1990s. J Hum Nutr Diet 14, 207–216
- 289. Johnson B & Hackett AF (2007) Trends in fruit, vegetable and salad intakes in 9–10-year-old schoolchildren living in Liverpool, 2000–2005. *Public Health Nutr* 10, 252–255.
- 290. Nelson M, Lowes K & Hwang V (2007) The contribution of school meals to food consumption and nutrient intakes of young people aged 4–18 years in England. *Public Health Nutr* 10, 652–662, Epublication 6 March 2007.
- 291. Noble S & Emmett P (2001) Food and nutrient intake in a cohort of 8-month-old infants in the south-west of England in 1993. *Eur J Clin Nutr* **55**, 698–707.
- 292. Pryer JA, Cook A & Shetty P (2001) Identification of groups who report similar patterns of diet among a representative national sample of British adults aged 65 years of age or more. *Public Health Nutr* 4, 787–795.
- Pryer JA, Nichols R, Elliott P, et al. (2001) Dietary patterns among a national random sample of British adults. J Epidemiol Community Health 55, 29–37.
- Rogers I & Emmett P (2002) Avon Longitudinal Study of Pregnancy and Childhood. Fat content of the diet among

- pre-school children in Britain; relationship with food and nutrient intakes. *Eur J Clin Nutr* **56**, 252–263.
- Swan G (2004) Findings from the latest National Diet and Nutrition Survey. Proc Nutr Soc 63, 505-512.
- 296. Thane CW, Bates CJ & Prentice A (2004) Zinc and vitamin A intake and status in a national sample of British young people aged 4–18 y. Eur J Clin Nutr 58, 363–375.
- Dykes F (2003) Feeding Initiative: A Report Evaluating the Breastfeeding Practice Projects 1999–2002: University of Central Lancashire Infant Preston. http://www.dh.gov.uk/en/ Publicationsandstatistics/Publications/PublicationsPolicyAnd Guidance/DH\_4084457
- Gregory J, Foster K, Tyler H, et al. (1990) The Dietary and Nutritional Survey of British Adults. London: HMSO.
- Gregory JR, Collins DL, Davies PSW, et al. (1995) National Diet and Nutrition Survey: Children Aged 1½ to 4½ Years.
   vol. 1: Report of the Diet and Nutrition Survey. London: HMSO.
- Finch S, Doyle W, Lowe C, et al. (1998) National Diet and Nutrition Survey: People Aged 65 Years and Over. vol. 1: Report of the Diet and Nutrition Survey. London: The Stationery Office.
- 301. Gregory JR, Lowe S, Bates CJ, et al. (2000) National Diet and Nutrition Survey: Young People Aged 4 to 18 Years.

- vol. 1: Report of the Diet and Nutrition Survey. London: The Stationery Office.
- Henderson L, Gregory J & Swan G (2002) National Diet and Nutrition Survey: Adults Aged 19 to 64 Years. vol. 1: Types and Quantities of Foods Consumed. London: The Stationery Office.
- 303. Henderson L, Gregory J & Irving K (2003) The National Diet and Nutrition Survey: Adults Aged 19 to 64 Years. vol. 2: Energy, Protein, Carbohydrate, Fat and Alcohol Intake. London: The Stationery Office.
- 304. Henderson L, Irving K & Gregory J (2003) The National Diet and Nutrition Survey: Adults Aged 19 to 64 Years. vol. 3: Vitamin and Mineral Intake and Urinary Analytes. London: The Stationery Office.
- Ruston D, Hoare J, Henderson L, et al. (2004) The National Diet and Nutrition Survey: Adults Aged 19 to 64 Years. vol. 4: Nutritional Status (Anthropometry and Blood Analytes), Blood Pressure and Physical Activity. London: The Stationery Office.
- Hoare J & Henderson L (2004) National Diet and Nutrition Survey: Adults Aged 19 to 64 Years. vol. 5: Summary Report. London: The Stationery Office.
- Cade J, Greenwood D, Burley V, et al. (2002) UK Women's Cohort Study. Leeds: University of Leeds.