

A complete asymptotic analysis of an oscillation free nonlinear equation of Bessel type with a pole in the dependent variable

A. B. Mingarelli ⁽¹⁾, J.M. Pacheco-Castelao⁽²⁾, S. Melkonian ⁽³⁾

Abstract

We characterize the solution set of a nonlinear perturbation of Bessel's equation of order zero on a half-line where the nonlinearity is analytic in the independent variable, algebraic in the dependent variable and, indeed, admits a pole in this variable. We show that the equation fails the Painlevé test and that there are no points in $[0, \infty)$ where blow-up occurs. Although we cannot find even one closed-form solution, it is shown that there are only four families of solutions: those that are asymptotically linear and increasing, solutions that are asymptotically linear and decreasing, another set of solutions that are asymptotically constant, and a final set of solutions that admit singularities at finite points on $[0, \infty)$. As a consequence, we deduce that every solution with or without singularities on $[0, \infty)$ is non-oscillatory and, in fact, has at most two zeros. We also show that the plane Π of real initial conditions $(y(0), y'(0))$ can be decomposed into a union of connected regions, in each of which the solutions are exactly one of the types mentioned above. Furthermore, we obtain that the set of those initial conditions leading to asymptotically constant solutions is a piecewise differentiable curve in Π , one that can be estimated theoretically to a high degree of precision. In addition, the asymptotic behavior of solutions near a finite singularity is obtained. Estimates relating the growth of solutions to their initial conditions are also described and numerical examples are presented to illustrate the theory. Finally, we observe that every solution of our equation has finite singularities when viewed as a solution on the whole line.

Résumé

Nous caractérisons l'ensemble de solutions d'une perturbation non-linéaire de l'équation de Bessel d'ordre zéro sur une demi-droite là où la non-linéarité est analytique dans la variable indépendante, algébrique dans la variable dépendante et, en effet, admet un pôle en cette variable. Nous démontrons que l'équation échoue au test de Painlevé et qu'il n'y a aucun point dans l'intervalle $[0, \infty)$ où la solution devient infinie. Bien que nous ne puissions pas trouver aucune solution sous forme fermée, on démontre qu'il existe que quatre familles de solutions; celles qui sont asymptotiquement linéaires et croissantes, les solutions qui sont asymptotiquement linéaires et décroissantes, un ensemble non-dénombrable de solutions

qui sont asymptotiquement constantes, et un ensemble final de solutions qui admettent des singularités à certains points finis de $[0, \infty)$. Par conséquent, l'on déduit que chaque solution avec ou sans singularités sur $[0, \infty)$ est non-oscillante et, en fait, admet tout au plus deux zéros. Nous prouvons également que le plan Π de valeurs initiales $(y(0), y'(0))$ se décompose en union de régions connexes dont les composantes renferment précisément un des types mentionnés ci-dessus. Nous obtenons aussi que l'ensemble de conditions initiales menant aux solutions asymptotiquement constantes est une courbe dérivable par morceaux dans Π , une courbe qui peut être estimée théoriquement à un degré élevé de précision. En outre, le comportement asymptotique de solutions au près d'une singularité finie est obtenu. Des évaluations reliant la croissance de solutions à leurs conditions initiales sont également décrites et des exemples numériques sont présentés pour fins d'illustration. En conclusion, l'on observe que chaque solution de notre équation (vue comme solution sur la droite entière) admet des singularités finies.

2000 Mathematics Subject Classification, 33C10, 34A34, 34C10, 34C15, 34E10.

Keywords and phrases: Bessel's equation of order zero, asymptotically linear, asymptotically constant, Banach-Cacciopoli fixed point theorem, disconjugate, oscillatory, non-oscillatory, Painlevé, movable singularities, movable critical points.

1 Introduction

Motivated by a recent paper of Dubé-Mingarelli [10], we present an in-depth study of the final example in [10] dealing with the existence of asymptotically constant solutions of the nonlinear differential equation

$$y'' + \frac{e^{-x}}{4}y + \frac{e^{-2x}}{1+y} = 0, \quad x \in [0, \infty), \quad (1)$$

where we note the possible presence of a singularity at a point x_0 where $y(x_0) = -1$. By a *classical solution* of (1) on $[0, \infty)$ we mean a function that is of class $C^2(0, \infty)$ which satisfies the equation everywhere. First of all, we note that (1) is actually a nonlinear perturbation of a Bessel equation of order zero in disguise. That is, the transformation $t = e^{-x/2}$, $y(x) = z(t)$, transforms (1) to an equation of the form

$$t^2 z'' + t z' + t^2 z + \frac{4t^4}{1+z} = 0, \quad t \in (0, 1]. \quad (2)$$

Our results for (1) will therefore apply to this equation (2) upon consideration of the quantity $z(t) = y(\ln(t^{-2}))$.

The formulation of general criteria for the existence of positive monotone solutions of second order nonlinear real differential equations of the form $y''(x) + F(x, y(x)) = 0$, defined on $[0, \infty)$, has been a subject of current interest. Methods used vary from using the fixed point theorem of Banach-Cacciopoli, Hale [12], to Schauder's fixed point theorem (see [12]), and other similar methods. The nonlinearity, being of the form $F(x, y(x))$, is not assumed to be variables-separable, nor is it assumed that estimates

of the form $|F(x, y)| \leq f(x)g(y)$ are in force as is usually the case in studies of this character, cf., Constantin [9], Nasr [21], Yin [28], Zhao [31]. In the cited papers, one finds criteria for the existence of at least one positive solution of a class of equations with various properties, as opposed to here, where we actually find a description of every possible solution of (1), positive or not. Next, we note that the asymptotic methods of Hartman-Onuchic [14] and Hale-Onuchic [13] are not applicable here because of the presence of the pole in the dependent variable, that is, given a real second order nonlinear differential equation of the form (in the notation of [14])

$$u'' = h(x, u, u'),$$

one normally requires the continuity of h as a function of the variable u for $u \in \mathbb{R}$, an assumption that is clearly not satisfied for our equation (1). Modern techniques require, at the very least, local integrability of h as a function of u , so that, once again, this fails.

The origin of Eq.(1) in Dubé-Mingarelli [10] is summarized next. In order to describe the novelty and applicability of the results obtained in Dubé-Mingarelli [10], one would need to find an equation for which most, if not all, existing non-oscillation criteria would not be applicable. In order to do this, one would need to construct an equation with at least three terms in order to circumvent the variables-separable case of F mentioned earlier. On the other hand, and to the best of our knowledge, no existing criterion applies in the case where a pole in the dependent variable is introduced. The two additional exponential terms in (1) were introduced in order to satisfy an integrability condition on both our function F , as a function of its second variable, and a function $k(t)$, leading to a generalized Lipschitz condition on F of Perron-Kamke-type (cf., Dubé-Mingarelli [10]). The resulting equation became a nonlinear perturbation of a second-order (non-oscillatory) linear equation with a nonlinearity having a pole in the dependent variable. As a whole, the resulting equation (i.e., (1)) turned out to be one which is rational in the dependent variables and analytic in the independent variable so that, on the surface, Painlevé theory may be applicable. However, such is not the case, as we show in the discussion below, since (1) fails the Painlevé Test (cf., Ablowitz *et al* [1], Ince [16]). Thus, the equation has movable critical points.

We present a brief non-exhaustive review of second-order equations with singularities that are poles in the dependent variable. In the event that the equation is analytic in its independent variable and rational in its dependent variable, this includes the Painlevé equations (50 of them) and the resulting theory (cf., Ince [[16], Chapter 14]) which, in the end, may be reduced to the study of six specific equations (denoted consecutively by $P_I, P_{II}, \dots, P_{VI}$). The study of these six Painlevé equations has gathered momentum in the past 20 years with the monograph of Ablowitz and Clarkson [1] and others and their applicability to inverse scattering, physics, etc. Their importance lies in part with the fact that their general solutions produce new transcendents, that is, solutions that cannot be expressed in terms of known transcendental functions, including the classical special functions of mathematical physics (Bessel, Airy, Hypergeometric, etc). The fact that a second-order equation (such as (1)) can fail the Painlevé test for the non-existence of movable critical points is of interest since we must then apply

methods specific to the equation itself in order to obtain information about its general solution.

Besides this special class of second-order equations bearing Painlevé's name, we find in the literature a study of special classes of nonlinear singular integral equations (albeit on a finite interval) where the singularities are possible poles in the dependent variable. In the event that the kernel is replaced by a suitable Green's function, we see that such equations can include second-order equations with similar nonlinearities as a special case. One of the earliest papers in this respect is Nowosad [22] where results from doubly-stochastic matrices (see also Michelli-Olsen [20]) are used, in particular, to detect the existence of continuous positive solutions on a finite interval of equations of the form

$$\bar{F}(x) \int_0^1 K(x, y) F(y) dy = 1, \quad 0 \leq x \leq 1, \quad (3)$$

where F is the Fourier transform of a complex-valued function $f \in L^2(\mathbb{R})$ and K is a positive kernel with additional properties (cf., Nowosad [22] for more details). An extension of this result to equations of the form

$$v(s) = \int_0^1 K(s, t) \frac{1}{v(t)^\alpha} dt, \quad 0 \leq s \leq 1, \quad (4)$$

is considered in Karlin-Nirenberg [18] where $\alpha > 0$, with a view to obtaining the existence of a continuous positive solution $v(t)$, $t \in [0, 1]$. However, the possibility that (4) may also admit solutions with movable zeros, i.e., points $x_0 > 0$ where $v(x_0) = 0$, is not entertained. A study of Volterra and Fredholm operators K on $\mathcal{C}[0, 1]$ appears in Bushell [8] with an eye to obtaining the existence (and uniqueness) of positive solutions of integral equations of the form $(I - K)x^p = f$, where $-1 < p < 1$, thus allowing for the possibility of a singularity in the dependent variable, x , in the nature of a branch point. An extension of the results in Karlin-Nirenberg [18] can be found in the articles by Parker-Walters [26], [29], once again relating the problem to fundamental results from matrix theory involving the similarity of an irreducible nonnegative square matrix to a doubly-stochastic matrix (Hartfiel's problem, see Parker-Walters [26]). We would not be led too far astray if we were to mention that a significant development of the theory arising from further studies of Nowosad [22] and Karlin-Nirenberg [18] lies in the connection with the so-called DAD problem, once again motivated by matrix theory (cf., Nussbaum [23], [24], [25] and Borwein *et al* [7]). More recently, a two-point boundary value problem on a finite interval in boundary layer theory appeared in Wang *et al* [30] where, once again, the nonlinearity involves a pole in the dependent variable, and the problem addresses the question of the existence and non-existence of positive solutions on the finite interval. In addition, more direct extensions of the main results in Karlin-Nirenberg [18] are also considered in Agarwal-O'Regan [3], [4] and [5], although the authors appear not to be aware of the extensions of Karlin-Nirenberg [18] via the DAD problem mentioned earlier.

For partial differential equations with singularities of the type we seek, we note Gomes [11] where the Dirichlet problem for $Lu = k(x)u^{-\alpha}$, $\alpha > 0$, and L is elliptic, is considered on a bounded region in \mathbb{R}^N , $N \geq 3$, with a view at obtaining the existence of a positive solution. A result complementing Gomes [11] appears in Jin [17] where, although the nonlinearity is more general than an inverse power of the dependent variable, the operator L is now the Laplacian on \mathbb{R}^n , $n \geq 3$. In addition, Jin [17] considers the Dirichlet problem on a finite domain or, in case the domain is all of \mathbf{R}^n , the solution sought is to be positive, as usual. In the same direction we find the work of Alves *et al* [6] that falls in between Gomes [11] and Jin [17] by allowing a linear perturbation of the Laplacian and a nonlinearity of the type considered by Gomes [11].

In this paper, we consider the problem of determining the asymptotics of every solution of (1). There are two possible approaches to determining the long-term behavior of these solutions of our equation; the first is to consider (1) as a perturbation of a Bessel equation as we have shown above. We can then use the method of variation of parameters and estimate the integrals using known asymptotics of Bessel functions under some assumptions on the nonlinearity. On the other hand, and this is the method we prefer, we use more analytical methods, which include fixed point theorems and Lyapunov functions, thereby allowing for further generality in the scope of the equations that can be studied.

Although (1) is basically a simple-looking differential equation, we will see that it has a rich structure, a structure that does not appear to have been studied in the literature in such generality. In this case, the plane Π of real initial conditions is divided into three connected regions separated by a piecewise differentiable curve along which initial data lead to asymptotically constant solutions. Another piece consists of initial data which produce solutions with finite singularities, while the remaining part gives asymptotically linear solutions. In Section 2, we initiate a study of the qualitative behavior of solutions by defining, in the first instance, a Lyapunov function that will be used extensively in the sequel. In Section 2.1 we connect (1) to the Painlevé program and show that it fails both the Painlevé and the Ablowitz-Ramani-Segur tests, thereby indicating the existence of movable critical points (which manifest themselves as branch points in our case). We also show that finite blow-up points do not exist for a given solution y of (1), but that both derivatives can and do blow up at finite points, the location of which depends upon the particular solution considered.

After introducing the notions of non-oscillation and disconjugacy in Section 2.2, we show that all non-negative solutions are actually disconjugate (i.e., have at most one zero in $[0, \infty)$). Section 3 is devoted to describing the complete set of solutions of (1) (with or without singularities on $[0, \infty)$) via their limits at infinity. In Section 3.1, we show that the solutions obtained in Section 3 are either asymptotically linear and increasing, asymptotically constant, asymptotically linear and decreasing or, finally, solutions with finite blow-up in the first and second derivatives (*finite singularities*). We also describe methods of finding them numerically via the contraction mapping principle. Finally, in Section 4, we connect the solutions of (1) to their initial condi-

tions and show how to find the initial conditions given the asymptotics and vice-versa. In addition, we show that the plane Π of real initial conditions $(y(0), y'(0))$ can be decomposed into a union of connected regions, in each one of which, the solutions are exactly one of the types mentioned above. Furthermore, we obtain that the set of those initial conditions leading to asymptotically constant solutions is a piecewise differentiable curve in Π , one that can be estimated theoretically to a high degree of precision. Estimates relating the growth of solutions to their initial conditions are also described and numerical examples are presented to illustrate the theory.

2 The qualitative behavior of solutions

We state here the required motivating result from Dubé-Mingarelli [10] for reference purposes.

Theorem 1. (See Example 3.5 in Dubé-Mingarelli [10])

Equation (1) admits a positive (on $[0, \infty)$) monotone solution that is asymptotic to one as $x \rightarrow \infty$.

We propose to analyze the remaining solutions of (1) including those with finite singularities. To this end we use the following notation. The initial conditions are assumed to be real and given by $y(0) \equiv y_0$, and $y'(0) \equiv y'_0$ and $y(a) \equiv y_a$, and $y'(a) \equiv y'_a$ where $a > 0$ when appropriate. The fundamental existence and uniqueness theorem guarantees the existence of local solutions of (1) on some small interval $[0, \delta]$, where $\delta > 0$, until a finite singularity is encountered (which generally moves with changes in the initial conditions). In this case the solution ceases to exist in a classical sense. Any interval $\mathcal{I} \subset [0, \infty)$ encountered in the sequel is assumed to be non-empty.

Remark 1. Note that there are three possible types of solution to equation (1): 1) Solutions y for which $y(x) > -1$ for all $x \in [0, \infty)$; 2) Solutions for which $y(x) < -1$ for all $x \in [0, \infty)$; and 3) Solutions for which there is a first point $x_0 \in [0, \infty)$, generally depending on the initial conditions, such that $y(x_0) = -1$ (their existence is considered in Section 2.1).

Lemma 1. Let y denote a solution of (1) defined on an interval $[a, x]$, where $a \geq 0$ and $x > a$. Then there holds the identity

$$\begin{aligned} & \frac{1}{2}y'(x)^2 + \frac{1}{8}e^{-x}y(x)^2 + e^{-2x} \ln |1 + y(x)| + \\ & + \frac{1}{8} \int_a^x e^{-t}y(t)^2 dt + 2 \int_a^x e^{-2t} \ln |1 + y(t)| dt \equiv I_a, \end{aligned} \quad (5)$$

where

$$I_a = \frac{1}{2}y'_a{}^2 + \frac{1}{8}e^{-a}y_a^2 + e^{-2a} \ln |1 + y_a|. \quad (6)$$

The implicit relation (6) defines a curve in Π : (y_a, y'_a) . The case of interest here is where $a = 0$, in which case this curve is bell-shaped and opens to the left. When there is no confusion we will denote I_0 simply by I , for brevity.

Proof. We multiply (1) by $y'(x)$ and integrate over the interval $[a, x]$. \square

The next results follow immediately from (5), thereby defining a Lyapunov function on non-negative solutions.

Corollary 1. If y is any non-negative solution of (1) on $[a, x]$, where $a \geq 0$, the function

$$H(x) = \frac{1}{2}y'(x)^2 + \frac{1}{8}e^{-x}y(x)^2 + e^{-2x} \ln |1 + y(x)|$$

is non-negative and non-increasing.

As a result,

Corollary 2. For any non-negative solution y of (1) over $[a, x]$, we have

$$\frac{1}{2}y'(x)^2 + \frac{1}{8}e^{-x}y(x)^2 + e^{-2x} \ln |1 + y(x)| \leq I_a. \quad (7)$$

Lemma 2. Choose a pair of initial conditions (y_0, y'_0) such that $I \leq 0$. Then the corresponding solution $y(x) \leq 0$ on any interval to the right of $x = 0$ so long as it is defined there (along with its second derivative).

Proof. Note that $y \equiv 0$ is not a solution of (1). Thus, let $y(x)$ be defined on $[0, \delta]$ (along with its second derivative) and suppose, on the contrary, that $y(x) \geq 0$ there. Then using Lemma 1 with $a = 0$ and $x = \delta$, we obtain an immediate contradiction. \square

In particular, we note that whenever $I \leq 0$ then $-2 \leq y_0 \leq 0$, $y_0 \neq -1$.

2.1 On the existence of movable critical points and movable singularities

We recall that a **critical point** of a differential equation is any singularity other than a pole (see Ince [[16], p.317] and Ablowitz-Clarkson [[1], p.348]). Such a critical point is said to be *movable* if its location depends upon the choice of the initial conditions y_0, y'_0 . A second-order nonlinear equation

$$y'' = R(z, y, y'), \quad z \in \mathbb{C},$$

where R is rational in y, y' and analytic in z (such as the complexification of (1)), is said to have the *Painlevé property* or to be of *Painlevé type* if it is free of movable critical points. We now show that (1) is not of Painlevé type (see Ablowitz-Clarkson [1] or Ince [16] for more information on such equations).

Theorem 2. *Eq. (1) is not of Painlevé type.*

Proof. Clearly (1) is rational in y, y' and analytic in x . Let $w(x) = 1 + y(x)$ in order to place the singularity at $w = 0$. Then (1) becomes

$$w'' = \frac{1}{4}e^{-x}(w - 1) + \frac{e^{-2x}}{w},$$

which has the form $w'' = L(x, w)(w')^2 + M(x, w)w' + N(x, w)$ with $L \equiv 0$ and $M \equiv 0$. According to the analysis of equations of this type given by Ince [[16], Sections 14.21 and 14.22, pp.321-325], since $L \equiv 0$, the first necessary condition for the absence of movable branch points is satisfied. However, since $N(x, w)$ has a pole at $w = 0$, the second necessary condition, which requires that the poles of M and N be included among those of L , is violated. Thus, (1) has movable branch points, and is therefore not of Painlevé type. \square

Remark 2. In addition, the ARS test [2] for the Painlevé property requires that the general solution be expressible in a series about a movable pole x_0 of order p , i.e.,

$$y = \sum_{n=0}^{\infty} a_n(x - x_0)^{n+p}, \quad p \in \mathbb{Z}^-,$$

containing two arbitrary constants, one of which is x_0 . Our analysis reveals that the leading-order term for (1) includes a logarithm, so that the equation admits movable branch points, which is further confirmation that it is not of Painlevé type. The asymptotic behavior of the solutions of (1) near such singularities is given by Theorem 5 below. Finally, our calculations show that (1) does not possess a nontrivial symmetry group, so that reduction or integration of the equation by such methods is also not possible.

Remark 3. We make the convention that a solution with a singularity at x_0 will be assumed to exist only up to x_0 and not past it. It follows that any solution of (1) without a singularity is a classical solution on $[0, \infty)$ and it must satisfy either $y(x) < -1$ for all $x \in [0, \infty)$ or $y(x) > -1$ for all $x \in [0, \infty)$ (cf., Remark 1).

Theorem 3. *Let y denote a solution of (1) with the property that for some $x_0 \in (0, \infty)$, $y(x)$ exists on $[0, x_0]$. If $y(x_0) = -1$, then both $|y'|$ and $|y''|$ become infinite at x_0 .*

Proof. For a given solution y there is no loss of generality in assuming that the x_0 in question is the smallest such point. Thus, let $y(x) > -1$ for every x in $[0, x_0)$. Observe that for a given x_0 , a function $f(x_0)$ can be defined by considering only the first, third, and fifth terms of the left side of (5). Its explicit form is

$$\frac{1}{2}y'(x_0)^2 + e^{-2x_0} \ln(1 + y(x_0)) + 2 \int_0^{x_0} e^{-2t} \ln(1 + y(t)) dt = f(x_0). \quad (8)$$

Let $x_0 > 0$ and let y be a solution with initial data set (y_0, y'_0) such that $y(x_0) = -1$. Then f in (8) can be expressed via (5) as

$$f(x_0) = \frac{1}{2}y_0'^2 + \frac{1}{8}y_0^2 + \ln(1 + y_0) - \frac{1}{8}e^{-x_0} - \frac{1}{8} \int_0^{x_0} e^{-t} y(t)^2 dt,$$

a quantity that is necessarily finite since y is continuous on $[0, x_0]$, and clearly, $y_0 \neq -1$.

Now, we estimate the integral appearing in (8). Since y is continuous on $[0, x_0]$ and $y(x_0) = -1$ there is a $\delta > 0$ such that $y(x) < -1/2$ for $x \in (x_0 - \delta, x_0]$, a δ which we now fix. Hence $\ln(1 + y(x)) < -\ln 2$ for such x . It follows that

$$\int_0^{x_0} e^{-2t} \ln(1 + y(t)) dt \leq \int_0^{x_0 - \delta} e^{-2t} \ln(1 + y(t)) dt - \ln 2 \int_{x_0 - \delta}^{x_0} e^{-2t} dt, \quad (9)$$

so that the integral on the left of (9) is bounded above. Thus, in order for $f(x_0)$ to remain finite, we must have $y'(x_0)^2 = \infty$ in (8). That $|y''(x_0)|$ is also infinite follows immediately from (1).

In the event that $y(x) < -1$ for every x in $[0, x_0)$, we repeat the above argument with the estimate $\ln(-1 - y(x)) < -\ln 2$ for x chosen so that $y(x) > -3/2$. The remaining argument is similar. \square

The next result complements Theorem 2 above and shows that our equation is actually free of any solutions which become infinite on $(0, \infty)$ (even though it does, in some cases, possess solutions the first and second derivatives of which become infinite).

Theorem 4. *Let y be any solution of (1). Then there is no point $x_0 \in [0, \infty)$ such that $|y(x_0)| = \infty$.*

Proof. If $y(x) \geq 0$ up to such a point x_0 , then its existence contradicts Corollary 2 above at that point. The general situation is similar, although slightly more technical. So assume, on the contrary, that there exists a solution y and $x_0 > 0$ with $|y(x_0)| = \infty$. There does not exist $x_1 \in (0, x_0)$ with $y(x_1) = -1$ since, otherwise, the solution would terminate at x_1 (cf., Remark 3).

Thus, $y \in C[0, x_0)$ and $y \in C[0, b]$ for every $b < x_0$. For such a $b < x_0$ we use (5) over $[0, b]$ in order to estimate the terms near the singularity x_0 . Doing away with the positive terms we have the estimate

$$e^{-2b} \ln |1 + y(b)| + 2 \int_0^b e^{-2t} \ln |1 + y(t)| dt \leq I_0, \quad (10)$$

where, we recall, I_0 is defined in (6). Passing to the limit as $b \rightarrow x_0^-$ in (10), using the fact that $|y(x_0)| = \infty$, and recalling that I_0 is a constant for a given solution, we obtain

$$\int_0^{x_0} e^{-2t} \ln |1 + y(t)| dt = -\infty. \quad (11)$$

Since $|y(x_0)| = \infty$, there is a $b < x_0$ such that $|1 + y(t)| \geq 1$ for $t \in [b, x_0)$. Fixing such a b we find that

$$\begin{aligned} \int_0^{x_0} e^{-2t} \ln |1 + y(t)| dt &= \int_0^b e^{-2t} \ln |1 + y(t)| dt + \int_b^{x_0} e^{-2t} \ln |1 + y(t)| dt, \\ &= C + \int_b^{x_0} e^{-2t} \ln |1 + y(t)| dt, \\ &\geq C, \end{aligned}$$

where C is a constant determined by the boundedness of the first integral on the right, since $y(x) \neq -1$ on $[0, b]$. However, this contradicts (11). \square

Corollary 3. Let y be a solution of (1) and $x_0 > 0$. If $y''(x_0)$ is finite, then both $y(x_0)$ and $y'(x_0)$ exist and are finite.

Proof. Clearly, $y(x_0)$ is finite by Theorem 4. By the identity (5) employed over the interval $[0, x_0]$, y' cannot be infinite since all other quantities therein are finite (as $y(x_0) \neq -1$). \square

Remark 4. The fundamental existence and uniqueness theorem now guarantees that if a solution y is such that $|y''| < \infty$ on $(0, \infty)$, then y exists on $[0, \infty)$.

A glance at the particular solution of (1) defined by $y_0 = 0, y'_0 = 0$ shows that its second derivative becomes unbounded at the value $x_0 \approx 2.366786778$. Thus, combining Theorems 3-4, we see that the particular solution defined by these zero boundary conditions at $x = 0$ is one which is finite, but with first and second derivatives which become infinite. The next result describes the behavior of such solutions near a singularity.

Theorem 5. *The asymptotic behavior of a solution y of (1) near a finite singularity $x_0 > 0$ is given by*

$$y(x) \sim -1 + \sqrt{2} e^{-x_0} (x - x_0) [-\ln(x - x_0)]^{1/2}, \quad x \rightarrow x_0^+,$$

and

$$y(x) \sim -1 + \sqrt{2} e^{-x_0} (x_0 - x) [-\ln(x_0 - x)]^{1/2}, \quad x \rightarrow x_0^-,$$

where the positive square root of the logarithmic term is taken if $y > -1$, and the negative if $y < -1$.

Proof. Let $w(x) = \frac{1}{1+y(x)}$ in order that $|w| \rightarrow \infty$ at the singularity, and transform (1) into

$$ww'' - 2(w')^2 - e^{-2x}w^4 + \frac{1}{4}e^{-x}(w^3 - w^2) = 0.$$

Let $\tau = x - x_0 \in \mathbb{C}$ and $w = a\tau^p$. Then a leading-order balance requires that $p = -1$, but this forces $a = 0$, which indicates that a logarithmic term must be present at leading order. Thus, put $w = a\tau^{-1}(\log \tau)^q$. Then the dominant terms in the equation are

$$\begin{aligned} ww'' &\sim 2a^2\tau^{-4}(\log \tau)^{2q} - 3a^2q\tau^{-4}(\log \tau)^{2q-1} + O(\tau^{-4}(\log \tau)^{2q-2}), \\ (w')^2 &\sim a^2\tau^{-4}(\log \tau)^{2q} - 2a^2q\tau^{-4}(\log \tau)^{2q-1} + O(\tau^{-4}(\log \tau)^{2q-2}), \\ w^4 &\sim a^4\tau^{-4}(\log \tau)^{4q}, \end{aligned}$$

and a balance of these terms requires that $2q - 1 = 4q$, i.e., $q = -1/2$, from which we obtain $a^2 = -e^{2x_0}/2$. Thus,

$$w \sim \frac{ie^{x_0}}{2^{1/2}}\tau^{-1}(\log \tau)^{-1/2},$$

and

$$y = -1 + \frac{1}{w} \sim -1 - 2^{1/2}ie^{-x_0}\tau(\log \tau)^{1/2}.$$

In order to adapt the above to the real solutions considered herein, let $x_0 \in \mathbb{R}$ and $\tau = x - x_0 > 0$. Then

$$y \sim -1 \pm \sqrt{2} e^{-x_0} (x - x_0) \sqrt{-\ln(x - x_0)}, \quad x \rightarrow x_0^+.$$

For solutions with $x < x_0$, the result is not altered if we let $\tau = x_0 - x > 0$, in which case

$$y \sim -1 \pm \sqrt{2} e^{-x_0} (x_0 - x) \sqrt{-\ln(x_0 - x)}, \quad x \rightarrow x_0^-.$$

□

By Theorem 5, it is clear that at a finite singularity x_0 , $y(x_0) = -1$ and that both y' and y'' are infinite at x_0 , in confirmation of Theorems 3-4 above.

2.2 On the non-oscillation of solutions

In this section we assume, unless otherwise specified, that the solutions under consideration have no singularities. As stated in Remark 3, we shall call such solutions *classical* solutions on $[0, \infty)$.

As is the custom in oscillation theory of differential equations, we say that an equation of the second order (whether linear or not) is **oscillatory** on a half-line provided that every (non-trivial) solution has arbitrarily large zeros. An equation is said to be non-oscillatory if it has at least one non-oscillatory solution. In addition, a solution is said to be **disconjugate** on an interval provided that it has at most one zero within it. For more information regarding such solutions in the linear case, the reader may refer to Swanson [27], although we note that a recent comprehensive survey of nonlinear oscillation theory is lacking (cf., Kartsatos [19]). Now, returning to our nonlinear problem (1), according to our convention (see Remark 3), such an oscillating solution will generally belong to the class of solutions such that $y(x) > -1$ for every $x \in [0, \infty)$, unless it is a solution with a critical point.

Lemma 3. Solutions of (1) cannot be identically constant on an interval $\mathcal{I} \subseteq [0, \infty)$.

Proof. Otherwise it is easy to see that its values $y(x)$ must satisfy the quadratic equation $y(x)^2 + y(x) + 4e^{-x} = 0$, in \mathcal{I} , by Eq. (1), which is a contradiction. □

The next result is actually a special case of another result proved in the next section, but it is of independent interest as its proof can be used in more general situations (see the Remark following the Lemma).

Lemma 4. Equation (1) has no non-negative oscillatory solutions. Indeed, every non-negative solution has at most one zero in $[0, \infty)$, and so is disconjugate there.

Proof. We proceed by contradiction. Should $y(x)$ be oscillatory it must have three zeros $x_1 < x_2 < x_3$. A double application of Rolle's theorem to $y(x)$ on $[x_1, x_2]$, $[x_2, x_3]$ gives the existence of points $x'_1 < x'_2$ such that $y'(x'_i) = 0$, for $i = 1, 2$. This, in turn, yields the existence of another point $x''_1 \in [x'_1, x'_2]$ such that $y''(x''_1) = 0$. Using (1) we see that $y(x''_1)$ must be a root of the quadratic equation $y^2 + y + 4e^{-x''_1} = 0$. However, a glance at its discriminant shows that the quantity $y(x''_1) = (-1 \pm \sqrt{1 - 16e^{-x''_1}})/2$

evidently satisfies $-1 < y(x_1'') < 0$, a contradiction.

Next, if a non-negative solution has two zeros in $(0, \infty)$, then its derivative must have two zeros as well. Thus, its second derivative must vanish somewhere, which contradicts the non-negativity hypothesis.

Finally, let $y(0) = 0$ and $y(x_0) = y'(x_0) = 0$, where without loss of generality we can assume that x_0 is the first such zero. Since $y(x) > 0$ in a neighborhood of x_0 (and since y cannot be identically constant there by Lemma 3) it follows that $y''(x_0) \geq 0$. However, a direct calculation using (1) shows that $y''(x_0) < 0$, a contradiction. \square

Remark 5. Of course, non-negative oscillatory solutions cannot exist for second order linear homogeneous equations but they can, indeed, exist for non-homogenous ones. For example, $y(x) = 1 - \sin x$ satisfies such an equation and is non-negative and oscillatory. On the other hand, the proof of this Lemma is easily modified to show that if $g : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$ and $g(x, y) > 0$ whenever $y \geq 0$, then the equation $y''(x) + g(x, y(x)) = 0$ has no non-negative oscillatory solutions among those that are continuable to \mathbb{R}^+ .

In the next section we will prove that all solutions with or without singularities on $[0, \infty)$ are, in fact, non-oscillatory and admit at most two zeros.

3 Asymptotic behavior

In this section we characterize the global asymptotic behavior of all the solutions of (1) in terms of their initial data set $(y_0, y_0') \in \mathbb{R}^2$, using the results of the proof of Theorem 6 above. In so doing we will show the existence of solutions which admit points where their second derivatives (and hence their first, by Theorem 3) become infinite.

Lemma 5. For any set of real initial data $(y_0, y_0') \in \mathbb{R}^2$ with $y_0 \neq -1$ we have $y''(0) \equiv y_0'' \neq 0$.

Proof. Note that when $y_0 = -1$ we cannot define a classical solution since its second derivative is automatically undefined. For $y_0 \neq -1$ we use (1) and see that $y_0'' = 0$ if and only if $y_0^2 + y_0 + 4 = 0$, which is impossible. The result follows. \square

Next, we characterize the solutions of (1) in terms of their asymptotic behavior, the following series of lemmata providing the proof of Theorem 6 below.

Theorem 6. *Equation (1) admits four families of solutions: Specifically, if y is such a solution then either*

1. $y(x) \rightarrow +\infty$, as $x \rightarrow \infty$
2. $y(x) \rightarrow C$, as $x \rightarrow \infty$ where C is a constant, $C \neq -1$
3. $y(x) \rightarrow -\infty$, as $x \rightarrow \infty$

4. $y(x)$ has a finite singularity where $y''(x)$ becomes infinite, so $y(x)$ cannot be continued past this point.

Terminology: In the following Lemmas the reference to the types (1), (2), (3), (4) refer to the types of solutions (1)-(4) respectively in the statement of Theorem 6.

Lemma 6. Let $y_0 \geq 0, y'_0 > 0$. Then the corresponding solutions are of type (1), (2) or (4).

Proof. Given that $y_0 > 0, y'_0 > 0$, then $y''_0 < 0$ and so, by continuity, there exists a point $a > 0$ such that $y(x) > 0, y'(x) > 0, y''(x) < 0$ for $x \in [0, a]$. It follows from this that y is increasing and y' is decreasing on $[0, a]$. There are now two possibilities:

Either there is a (smallest) point $c > a$ such that $y'(c) = 0$, or there is no such point, in which case $y'(x) > 0$ for every $x \in [0, \infty)$. (We note that global existence is obtained in the latter case because of the boundedness of y and so of y' on bounded intervals, cf. Remark 4 and Corollary 3).

In the latter case, since y' is decreasing (it is bounded above) and $y'(x)$ has a limit $\beta \geq 0$ as $x \rightarrow \infty$. If $\beta > 0$ then it is a simple matter to see that $y(x) \rightarrow +\infty$ as $x \rightarrow \infty$ and so (1) holds. On the other hand, $\beta = 0$ is only possible in the case of asymptotically constant solutions ($y(x) \rightarrow C$ for some positive constant C , as $x \rightarrow \infty$) or solutions for which (1) holds. Thus, a solution is either of type (1) or (2) in this special case.

In the former case, let $c > a$ be such that $y'(c) = 0$, and without loss of generality we can assume that c is the first such point. Since $y(c) > 0$ we have $y''(c) < 0$ so that c is a local maximum (recall that $y(x)$ cannot be identically constant past c on account of Lemma 3). Thus, there exists $b > c$ such that $y'(x) < 0$ for $x \in (c, b]$ and since y' is decreasing there it follows that for all $x > b$ such that $y(x) \geq 0$, we must have $y'(x) < y'(b) < 0$. This, however, forces the existence of a point $d > c$ such that $y(d) = 0$. Now we employ (1) to find that $y''(d) < 0$. Because of this there now exists b_1 such that $y(x) < 0, y''(x) < 0$, and so $y'(x)$ is decreasing for $x \in (d, b_1]$. Arguing as before we have $y'(x) < y'(d) < 0$ in a right-neighborhood of $x = d$. However this implies that $y(x) \leq y(d) + y'(d)(x - d)$, for all x in this neighborhood. It follows that if this y is continuable to \mathbb{R}^+ , then $y(x) \rightarrow -\infty$ as $x \rightarrow \infty$. But this forces the existence of a point $x_0 > d$ such that $y(x_0) = -1$ by continuity (since $y_0 > 0$). Hence such a solution is of type (4). It follows that for such initial conditions our solutions are either of type (1), (2) or (4) in Theorem 6.

The case where $y_0 = 0$ is similar to the opening paragraph to this proof and so is omitted. There are then the three possibilities again, type (1), (2) or (4), depending on the size of the values of y'_0 . \square

Lemma 7. Let $y_0 > 0, y'_0 \leq 0$. Then the corresponding solutions all have infinite second (and hence first) derivatives at a finite point (i.e., of type (4)).

Proof. We argue as in Lemma 6 above and thus sketch the proof. Consider first the case where $y_0 > 0, y'_0 = 0$. Then $y''_0 < 0$ so that there exists a point $a > 0$ such that $y(x) > 0, y'(x) < 0, y''(x) < 0$ for $x \in [0, a]$. So, y' is decreasing on $[0, a]$ and the inequality $y'(x) < y'(a) < 0$ is in force whenever the derivative is defined. Thus, there is a point $b > a$ such that $y(b) = 0$ and so $y''(b) < 0$, by (1). It follows that $y(x_0) = -1$ for some $x_0 > b$, i.e., y is of type (4). The proof where $y'(0) < 0$ is similar and so we omit it. \square

Lemma 8. Let $-1 < y_0 < 0, y'_0 > 0$. Then the solutions are either of type (1), (2) or (4).

Proof. Once again we sketch the proof, which in this case parallels that of Lemma 6. For y_0, y'_0 as given there exists a point $a > 0$ such that $y(x) < 0, y'(x) > 0$, for $x \in [0, a]$. Next, Lemma 5 indicates that $y''_0 \neq 0$, and in fact, it is the case that $y''_0 < 0$. This is because the alternative $y''_0 > 0$ is equivalent (by (1)) to the inequality $y_0(1 + y_0) + 4 < 0$ for $-1 < y_0 < 0$. However, this last quadratic inequality is impossible. Thus, we can restrict a further, if need be, to obtain that $y(x) < 0, y'(x) > 0, y''(x) < 0$ in $[0, a]$ (so that y' is decreasing once again). There are now the two usual possibilities (see the proof of Lemma 6): Either there is a (smallest) point $c > a$ such that $y'(c) = 0$, or there is no such point c in which case $y'(x) > 0$ and decreasing on $[0, \infty)$.

Clearly, the existence of such a point c gives a local maximum for y . In addition, since y' is decreasing to the right of c we must have $y(x) = -1$ for some $x = x_0 > c$. Hence, the solution is of type (4).

The alternative that $y'(x) \neq 0$ gives us $y'(x) > 0$. Using standard arguments we can now conclude (see above) that there exists a point b at which $y(b) = 0$ (and again $y''(b) < 0$.) If $y'(x)$ is bounded away from zero it follows that $y(x) \rightarrow \infty$ as $x \rightarrow \infty$. On the other hand, if $y'(x) \rightarrow 0$ as $x \rightarrow \infty$ then either y is of type (1) or type (2). \square

Lemma 9. Let $-1 < y_0 \leq 0, y'_0 \leq 0$. Then the solutions are of type (4).

Proof. For there is an interval of the form $[0, a]$ in which we have $-1 < y(x) < 0, y'(x) < 0, y''(x) < 0$, for $x \in [0, a]$ (see Lemma 8 above). It follows that $y'(x) < 0$ and decreasing for $x > a$ and so we must have $y(x) = -1$ for some $x = x_0 > a$. Once again, the solution is of type (4). The case $y_0 = 0$ is similar, since there is a neighborhood $[0, a]$ in which $y(x) < 0, y'(x) < 0, y''(x) < 0$ for $x \in [0, a]$, etc. \square

Lemma 10. Let $y_0 < -1$. If $y'_0 > 0$, the solution is of type (4). On the other hand, if $y'_0 \leq 0$, the solution is either of type (2), (3) or (4).

Proof. The proof is very similar to that of Lemma 6. For there is an interval of the form $[0, a]$ in which we have either $y(x) < -1, y'(x) < 0, y''(x) > 0$, for $x \in [0, a]$ or $y(x) < -1, y'(x) > 0, y''(x) > 0$ (see Lemma 6 above). In either case it follows that $y'(x)$ is monotone for every $x > a$ and so we may have either $y(x) = -1$ for some $x = x_0 > a$, $y(x) \rightarrow -\infty$ as $x \rightarrow \infty$ (if $y'(x) \rightarrow -\beta < 0$), or $y(x) \rightarrow -\infty$ or $y(x) \rightarrow C$ as $x \rightarrow \infty$ (if $y'(x) \rightarrow 0$), corresponding to the stated initial conditions. Thus, y is either of type (2), (3) or (4). \square

A consequence of the proofs of these Lemmas is that every solution of (1) has at most one point in $(0, \infty)$ where its derivative can vanish, so one can derive that

Corollary 4. Every solution of (1) has at most two zeros on its interval of existence (i.e., every solution is non-oscillatory).

Referring to Fig. 1 the initial conditions for the solutions reckoned from top to bottom are: $(2, 1)$, $(2, 0.744448)$, $(2, 0.6)$, respectively. The top-most follows an increasing linear trend (Type (1) solution) given by the line $y'_0 = 0.200043y_0 + 2.8205$, the middle one is approaching a constant value $C \approx 2.70734$ (Type (2) solution), while the bottom one eventually hits a singularity at $x_0 \approx 32.13$ (Type (4) solution).




Figure 1.

An example of a Type (3) solution is given below in Fig. 2 where the initial conditions are $(-2, -1)$.




Figure 2.

3.1 Asymptotics

Combining Lemmas 6-10 in the previous section we can infer that if y is any solution of (1) defined on the whole half-axis then its derivative $y'(x) \rightarrow \beta$ where $\beta > 0, \beta < 0$ or $\beta = 0$ ($\beta = \pm\infty$ occurs when we have a finite singularity). Next, we show that the technique presented in Dubé-Mingarelli [10] can be modified so that in addition to the particular solution appearing in Theorem 1 there is a continuum of *asymptotically constant* solutions of (1) corresponding to the situation where $y'(x) \rightarrow 0$. We recall that y is said to be asymptotically constant as $x \rightarrow \infty$, and write $y(x) \sim M$, if $y(x) = M + o(1)$ as $x \rightarrow \infty$, where M is a constant. As well, we say that a solution is *asymptotically linear* if it is of the form $y(x) = ax + b + o(1)$ as $x \rightarrow \infty$, for suitable constants a, b . We usually write $y(x) \sim ax + b$ to mean that y is asymptotic to the line $ax + b$ as $x \rightarrow \infty$.

We proceed to show that the family of solutions (types (1), (2) and (3)) described in Theorem 6 consists solely of asymptotically linear solutions (not excluding the asymptotically constant ones). From this we conclude that either a solution of (1) has a singularity (type (4)) or it is asymptotically linear, thus giving a complete characterization of all the solutions of (1).

Lemma 11. Let y be any solution of (1) that is not of type (4) in Theorem 6. Then

$$\lim_{t \rightarrow \infty} \frac{y(t)}{t} = \beta$$

and $y'(t) - \beta \in L^1(0, \infty)$.

Proof. From the proof of Theorem 6 we know that $y(t)$ exists on $[0, \infty)$ and there is a $\beta \in \mathbb{R}$ such that $y'(t) \rightarrow \beta$ as $t \rightarrow \infty$. Without loss of generality we assume that $\beta > 0$ (the case $\beta < 0$ being similar).

Given $\varepsilon > 0$ we choose $T_0 > 0$ so large that $|y'(t) - \beta| < \varepsilon/2$, for all $t > T_0$. Fix such a T_0 . Then

$$|y(t) - y(T_0) - \beta(t - T_0)| \leq \int_{T_0}^t |y'(\tau) - \beta| d\tau < \frac{\varepsilon}{2}(t - T_0),$$

for all $t > T_0$. Hence

$$\left| \frac{y(t)}{t} - \beta \right| \leq \frac{\varepsilon}{2} \left(1 - \frac{T_0}{t} \right) + \left| \frac{y(T_0) - \beta T_0}{t} \right| < \varepsilon,$$

provided we choose $T_1 > T_0$ so large that $\left| \frac{y(T_0) - \beta T_0}{t} \right| < \varepsilon/2$, for all $t > T_1$. We deduce that $y(t)/t \rightarrow \beta$ as $t \rightarrow \infty$.

Note that we can now choose t so large that we have $\beta/2 < |y(t)/t| < 3\beta/2$, for $t > T_3$. If necessary, we can increase t further so that $|1 + y(t)| = |1 + t(y(t)/t)| > 1$, for all $t > T_4 > T_3$. It follows that for every $x > T_4$,

$$\int_x^\infty |t F(t, y(t))| dt < \int_x^\infty \left(\frac{1}{4} t^2 e^{-t} \left| \frac{y(t)}{t} \right| + \frac{t e^{-2t}}{|1 + y(t)|} \right) dt < \int_x^\infty \left(\frac{3}{8} \beta t^2 e^{-t} + t e^{-2t} \right) dt < \infty,$$

where

$$F(t, y) = \frac{1}{4}e^{-t}y + \frac{e^{-2t}}{1+y}.$$

Hence the integral on the left is absolutely convergent on $[0, \infty)$.

Keeping in mind that $y'(t) \rightarrow \beta$ as $t \rightarrow \infty$, we can now integrate (1) over $[x, \infty)$ and so find that

$$y'(x) = \beta + \int_x^\infty F(t, y(t)) dt. \quad (12)$$

From this and Fubini's theorem we can derive that for all $x > T_4$,

$$\begin{aligned} \int_x^\infty |y'(t) - \beta| dt &\leq \int_x^\infty \int_t^\infty |F(s, y(s))| ds dt \\ &= \int_x^\infty (t-x) |F(t, y(t))| dt \\ &< \int_0^\infty t |F(t, y(t))| dt < \infty. \end{aligned}$$

The second conclusion follows (the proof of the case $\beta < 0$ is similar and so is omitted).

Note that when $\beta = 0$ one essential change must occur in the argument. Once again, we derive that $y(t)/t \rightarrow 0$ as $t \rightarrow \infty$. However, we must now choose T so large that $|1 + y(t)| \geq \varepsilon > 0$ for all $t > T$, which is possible since $|y(x)| \rightarrow \infty$ or $|y(x)| \rightarrow C \neq -1$. Thus, $y'(t) \in L^1(T, \infty)$. In addition, since y is not a type (4) solution, it follows that the values $y(t)$ must be bounded away from $t = -1$, for every value of t . So there is a $\delta > 0$ such that $|1 + y(t)| > \delta$ for all $t \in [0, T]$, from which we get $y'(t) \in L^1(0, T)$, and this produces the desired integrability over $[0, \infty)$. \square

Corollary 5. If y is not a type (4) solution of (1) then it is asymptotically linear (or asymptotically constant in the case where $\beta = 0$).

Proof. Since $y' - \beta \in L^1(0, \infty)$,

$$y(x) - y(0) - \beta x = \int_0^x (y'(t) - \beta) dt \rightarrow c \text{ as } x \rightarrow \infty,$$

so that

$$y(x) = \beta x + y(0) + c + \left(\int_0^x (y'(t) - \beta) dt - c \right) = \beta x + y(0) + c + o(1),$$

i.e., $y(x) \sim \beta x + \alpha$. \square

The next four Lemmas provide a means for approximating (and proving the existence of) these asymptotically linear (or asymptotically constant) solutions using the contraction mapping principle of Banach-Cacciopoli, Hale [[12], p.5]. In so doing we obtain estimates for the sizes of the various solutions involved and initial conditions leading to them (more on this in the next section).

Theorem 7. *Let $M, N, M > N \geq 0$ be given numbers such that*

$$(3M - 4N)(N + 1) \geq 1. \tag{13}$$

Then there is a unique asymptotically constant solution of (1) such that $N \leq y(x) \leq M$ and $y(x) \rightarrow M$ as $x \rightarrow \infty$.

Proof. We specialize the proof of Theorem 2.1 in Dubé-Mingarelli [10] to this situation (in conjunction with Example 3.5 on p.6 of Dubé-Mingarelli [10]). We define the space $X = \{u \in C[0, \infty) \mid N \leq u(t) \leq M, t \geq 0\}$, and note that it is a complete metric space with metric induced by the uniform norm on the half-line. We define

$$F(t, u(t)) = \frac{e^{-t}}{4}u(t) + \frac{e^{-2t}}{1 + u(t)},$$

and note that for $u \in X$, we have (cf., Eq. (2.1) in Dubé-Mingarelli [10])

$$\int_0^\infty tF(t, u(t)) dt \leq \int_0^\infty \left(\frac{1}{4}te^{-t}M + \frac{te^{-2t}}{1 + N} \right) dt = \frac{1}{4} \left(M + \frac{1}{N + 1} \right) \leq M - N$$

by (13). This shows that the mapping T defined by (cf., Eq. (2.4) in Dubé-Mingarelli [10])

$$Tu(x) = M - \int_x^\infty (t - x)F(t, u(t)) dt$$

maps $X \rightarrow X$. In addition, we choose $k(t)$ so that

$$k(t) = \frac{1}{4}e^{-t} + \frac{e^{-2t}}{(1 + N)^2}, \tag{14}$$

where we have fixed a typographical error in the corresponding expression for (14) with $N = 0, M = 1$ in Dubé-Mingarelli [[10], Example 3.5]. Then

$$\int_0^\infty tk(t) dt = \frac{1}{4} \left(1 + \frac{1}{(N + 1)^2} \right) < 1$$

by our assumption on N at the outset, as required for the application of the fixed point theorem. It is now an easy matter to verify that T is a contraction on X . Therefore its fixed point corresponds to an asymptotically constant solution of (1). \square

Setting $N = 0$ in this theorem immediately gives the next consequence.

Corollary 6. For any $M \geq 1/3$ there is a unique asymptotically constant solution of (1) such that $0 \leq y(x) \leq M$ and $y(x) \rightarrow M$ as $x \rightarrow \infty$.

The preceding corollary extends the application of Dubé-Mingarelli [[10], p.6] in the case where $M = 1$ and $N = 0$ to a case of general M, N . Now we show that among those solutions with $y'(x) \rightarrow \beta$ where $\beta > 0$ there is a continuum of unique asymptotically linear ones that can also be calculated to an arbitrary degree of precision using the method of successive approximations (which is built-into the contraction mapping principle).

Theorem 8. Let $\mathcal{D} = \{(a, b) \in \mathbb{R}^2 \mid a > 0, b > 0, 3a - 2b < 0, 3b - 2a \geq 1\}$. Then for each $a, b \in \mathcal{D}$ there is a unique solution of (1) such that $0 \leq y(x) \leq ax + b$ and $y(x) \sim ax + b$ as $x \rightarrow \infty$.

Proof. The proof is in the same spirit as that of the aforementioned Theorem 2.1 in Dubé-Mingarelli [10]. We redefine the space

$$X = \{u \in C[0, \infty) \mid 0 \leq \frac{u(t)}{at + b} \leq 1, t \geq 0\},$$

and note that once again, for fixed a, b , X is a complete metric space with metric induced by the norm

$$\|u\|_X = \sup_{t \geq 0} \left| \frac{u(t)}{at + b} \right|.$$

Define the map T on X by

$$Tu(x) = ax + b - \int_x^\infty (t - x)F(t, u(t)) dt.$$

Clearly, for $u \in X$ we have $Tu(x) \in C[0, \infty)$. Defining F as in Theorem 7 above, a simple estimation shows that for $u \in X$ we get

$$\int_0^\infty tF(t, u(t)) dt \leq \int_0^\infty \left(\frac{1}{4}te^{-t}(at + b) + te^{-2t} \right) dt = \frac{1}{4}(1 + b) + \frac{1}{2}a \leq b$$

since $(a, b) \in \mathcal{D}$. Observe that since $3b - 2a \geq 1$ we have $Tu(x) \geq ax + b - b = ax \geq 0$, for all $x \geq 0$. (cf., Eq. (2.6) in Dubé-Mingarelli [10]). Moreover, $Tu(x) \leq ax + b$, since the integral is non-negative on X . Thus T is a self-map. We choose $k(t)$ as before (see (14) above). Next we show that T is a contraction on X . To this end we note that for $u, v \in X$,

$$\begin{aligned} \left| \frac{Tu(x) - Tv(x)}{ax + b} \right| &\leq \frac{1}{ax + b} \int_x^\infty (t - x) |at + b| k(t) \left| \frac{u(t) - v(t)}{at + b} \right| dt \\ &\leq \frac{1}{b} \sup_{t \geq 0} \left| \frac{u(t) - v(t)}{at + b} \right| \int_x^\infty (t - x)(at + b) \left(\frac{1}{4}e^{-t} + e^{-2t} \right) dt \\ &\leq \frac{1}{b} \|u - v\|_X \int_0^\infty t(at + b) \left(\frac{1}{4}e^{-t} + e^{-2t} \right) dt \\ &= \frac{1}{b} \left(\frac{3}{4}a + \frac{1}{2}b \right) \|u - v\|_X \\ &\leq \kappa \|u - v\|_X \end{aligned}$$

where $0 < \kappa < 1$ if and only if $3a < 2b$. Passing to the supremum on the left it follows that T is a contraction on X since $3a - 2b < 0$ by hypothesis. The existence of a fixed point for T is established, and so is the existence of an asymptotically linear solution of (1). \square

Remark 6. The point $(a, b) = (0, 1/3)$ is on the boundary of \mathcal{D} and this gives an asymptotically constant solution as we have seen in Theorem 7. Indeed, all the solutions obtained via Theorem 7 appear as boundary points of \mathcal{D} . As an application of Theorem 8 we conclude that there is a solution of (1) such that $y(x) \sim x + 2$ as $x \rightarrow \infty$ since the point $(1, 2) \in \mathcal{D}$.

A result corresponding to Theorem 7 but for *negative* asymptotically constant solutions is next.

Theorem 9. *Let $\alpha > 1 + 1/\sqrt{3}$, and choose $M \geq \max\{8/3, \alpha\}$ so large that*

$$\alpha \leq \frac{1}{2} + \frac{3}{8}M + \frac{1}{8}\sqrt{3}\sqrt{M(3M-8)}.$$

Then (1) has a unique asymptotically constant solution y such that

$$-M \leq y(x) \leq -\alpha, \quad x \geq 0,$$

and $y(x) \sim -M$ as $x \rightarrow \infty$.

Proof. Given α we fix a value of M as per the assumptions. Then the space

$$X = \{u \in C[0, \infty) \mid -M \leq u(t) \leq -\alpha, t \geq 0\},$$

is a complete metric space with metric d defined by $d(u, v) = \max_{t \geq 0} |u(t) - v(t)|$. We define a map T on X as usual by setting

$$Tu(x) = -M - \int_x^\infty (t-x)F(t, u(t)) dt,$$

where F is defined as before by

$$F(t, u(t)) = \frac{1}{4}u(t)e^{-t} + \frac{e^{-2t}}{1+u(t)}, \quad u \in X.$$

Note that since $u \in X$ we have $F(t, u(t)) < 0$ for every $t \geq 0$. In addition, $Tu(x)$ is continuous for $x \geq 0$ and $Tu(x) + M \geq 0$, for every $x \geq 0$. We now need to estimate T from above in order to show that T is a self-map. Since $u \in X$,

$$\begin{aligned} Tu(x) &= -M - \int_x^\infty (t-x) \left(\frac{1}{4}u(t)e^{-t} + \frac{e^{-2t}}{1+u(t)} \right) dt \\ &\leq -M + \int_x^\infty (t-x) \left(\frac{1}{4}Me^{-t} + \frac{e^{-2t}}{\alpha-1} \right) dt \\ &\leq -M + \int_0^\infty \left(\frac{1}{4}Mte^{-t} + \frac{te^{-2t}}{\alpha-1} \right) dt \\ &= -\frac{3}{4}M + \frac{1}{4(\alpha-1)}. \end{aligned}$$

Thus, $Tu(x) \leq -\alpha$ provided we can make $-\frac{3}{4}M + \frac{1}{4(\alpha-1)} \leq -\alpha$, an expression that leads to the quadratic inequality $4\alpha^2 - (4 + 3M)\alpha + 3M + 1 \leq 0$. The discriminant

being given by $3M(3M - 8)$, we see that it is necessary that $M \geq 8/3$. Furthermore, $M \geq \alpha$ by definition of the set X and so $M \geq \max\{8/3, \alpha\}$ as required. A solution of the quadratic inequality is given by choosing M larger still, if necessary, so that $\alpha \leq \frac{1}{2} + \frac{3}{8}M + \frac{1}{8}\sqrt{3}\sqrt{M(3M - 8)}$. Fixing such a value of M we have $Tu(x) \leq -\alpha$ and so $Tu(x) \in X$. Hence $T : X \rightarrow X$.

In order to show that T is a contraction on X (and hence continuous) we proceed as follows: For $u, v \in X$,

$$\begin{aligned} |Tu(x) - Tv(x)| &\leq \int_x^\infty (t-x) |F(t, u(t)) - F(t, v(t))| dt \\ &\leq \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t}|u(t) - v(t)| + \frac{e^{-2t}|u(t) - v(t)|}{|1+u(t)||1+v(t)|} \right) dt \\ &\leq \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t}|u(t) - v(t)| + \frac{e^{-2t}|u(t) - v(t)|}{(\alpha-1)^2} \right) dt \\ &\leq d(u, v) \int_0^\infty \left(\frac{1}{4}te^{-t} + \frac{te^{-2t}}{(\alpha-1)^2} \right) dt \\ &\leq \kappa d(u, v), \end{aligned}$$

where $\kappa = (1 + (\alpha - 1)^{-2})/4 < 1$ since $\alpha > 1 + 1/\sqrt{3}$ by hypothesis. As a result we find $d(Tu, Tv) \leq \kappa d(u, v)$ for $u, v \in X$. So T is a contraction and thus has a fixed point $u \in X$. This fixed point corresponds to a solution y with the stated properties. \square

Remark 7. These methods give rough estimates to the initial conditions that generate the required solution. For example, it follows from Theorem 9 that the solution $y(x)$ with $y(x) \sim -M$ satisfies $-M \leq y_0 \leq -\alpha$. Thus, in practice it is best to find the smallest interval of length $M - \alpha$.

The final result in this section is an analog of Theorem 8 for negative solutions satisfying the condition $y'(x) \rightarrow -\beta$ as $x \rightarrow \infty$ where $\beta > 0$. It is somewhat more technical than the stated result since we must stay away from a point where $y = -1$. Its proof is a combination of the proofs of both Theorems 8-9.

Theorem 10. *Let $\alpha > 1$, $a > 0$, $b > 0$, and*

$$\mathcal{D} = \left\{ (a, b) \in \mathbb{R}^2 \mid 3b - 2a \geq 4\alpha + \frac{1}{\alpha - 1}, \frac{3a - 2b}{3b - 2a} < \alpha(\alpha - 2) \right\}.$$

Then for each $a, b \in \mathcal{D}$ there is a unique solution of (1) such that $-ax - b \leq y(x) \leq -\alpha$ for $x \geq 0$, and $y(x) \sim -ax - b$ as $x \rightarrow \infty$.

Proof. We define a complete metric space by $X = \{u \in C[0, \infty) \mid -at - b \leq u(t) \leq -\alpha, t \geq 0\}$, with metric $d(u, v)$ defined by the norm

$$d(u, v) = \sup_{t \geq 0} \left| \frac{u(t) - v(t)}{-at - b} \right| = \|u - v\|_X.$$

As is usual we define the map T on X by

$$Tu(x) = -ax - b - \int_x^\infty (t-x)F(t, u(t)) dt.$$

Once again, for $u \in X$ we have $Tu(x) \in C[0, \infty)$. First we show that T is a self-map.

To this end, let $u \in X$. Since $u(t) < -\alpha < 0$ and $\alpha > 1$, $F(t, u(t)) < 0$ for every $t \geq 0$. Hence $Tu(x) + ax + b \geq 0$, for all $x \geq 0$. In addition, for $u \in X$,

$$\begin{aligned}
Tu(x) &= -ax - b - \int_x^\infty (t-x)F(t, u(t)) dt \\
&= -ax - b + \int_x^\infty (t-x)|F(t, u(t))| dt \\
&\leq -ax - b + \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t}(at+b) + \frac{e^{-2t}}{\alpha-1} \right) dt \\
&\leq -ax - b + \int_0^\infty t \left(\frac{1}{4}e^{-t}(at+b) + \frac{e^{-2t}}{\alpha-1} \right) dt \\
&\leq -ax - b + \frac{1}{4(\alpha-1)} + \frac{1}{2}a + \frac{1}{4}b \leq \frac{1}{4(\alpha-1)} + \frac{1}{2}a - \frac{3}{4}b
\end{aligned}$$

for each $x \geq 0$. Since $(a, b) \in \mathcal{D}$, $Tu(x) \leq -\alpha$ and so $T : X \rightarrow X$.

We show that T is a contraction on X . Arguing as in the proof of Theorem 8 we find for $u, v \in X$,

$$\begin{aligned}
\left| \frac{Tu(x) - Tv(x)}{-ax - b} \right| &\leq \frac{1}{ax + b} \int_x^\infty (t-x) |at+b| \left(\frac{1}{4}e^{-t} + \frac{e^{-2t}}{(\alpha-1)^2} \right) \left| \frac{u(t) - v(t)}{at+b} \right| dt \\
&\leq \frac{1}{b} \sup_{t \geq 0} \left| \frac{u(t) - v(t)}{at+b} \right| \int_x^\infty (t-x)(at+b) \left(\frac{1}{4}e^{-t} + \frac{e^{-2t}}{(\alpha-1)^2} \right) dt \\
&\leq \frac{1}{b} d(u, v) \int_0^\infty t(at+b) \left(\frac{1}{4}e^{-t} + \frac{e^{-2t}}{(\alpha-1)^2} \right) dt, \quad \text{so that} \\
d(Tu, Tv) &\leq \kappa d(u, v),
\end{aligned}$$

where the contraction constant

$$\kappa = \frac{(2a+b)\alpha^2 - (4a+2b)\alpha + (3a+2b)}{4b(\alpha-1)^2} < 1$$

if and only if

$$(2a-3b)\alpha^2 - (4a-6b)\alpha + (3a-2b) < 0, \tag{15}$$

a quadratic with discriminant $4(3b^2 + ab - 2a^2) = 4(b+a)(3b-2a) > 0$ because $(a, b) \in \mathcal{D}$. In addition, $2a-3b < 0$, so (15) is equivalent to

$$\alpha(\alpha-2) > \frac{3a-2b}{3b-2a}. \tag{16}$$

Thus, with a, b suitably restricted as in Eq. (16), we get $\kappa < 1$ and so T is a contraction on X . Note that if $\alpha > 2$ and $3a = 2b$, then the first of the two conditions on a, b in \mathcal{D} is sufficient to guarantee the existence of asymptotically linear solutions of this type. \square

Remark 8. We emphasize once again that one advantage in using fixed point methods to prove the existence of asymptotically linear solutions of (1) lies in the fact that the Banach-Cacciopoli fixed point theorem actually implies that the method of “successive approximations” (or *Picard iterations* as they are sometimes called) can be used to approximate the solutions themselves, i.e., the method lends itself to computability (see Hille [[15], Chapter 2]). Although the Schauder fixed point theorem is more general and can be used here, it only gives existence (without uniqueness) and little or no computable features for the solutions themselves.

Example 1. In order to determine the initial conditions that generate the solution in Theorem 1, we simply write

$$y_{n+1}(x) = 1 - \int_x^\infty (t - x)F(t, y_n(t)) dt, \quad n \geq 1,$$

where $0 \leq y_1(t) \leq 1$ is given by say, $y_1(t) = 1/2$. These iterations must converge to the required solution by the stated fixed point theorem.

Finding the first few iterates and substituting the value $x = 0$ in both y_n and its derivative gives the approximate values $y_0 = 0.6236, y'_0 = 0.49148$ (see Fig. 3 below). The more iterations one uses the better the approximation. Of course, any seed (within the required limits) can be used as a starting value and the iterations will all converge to the unique solution sought.




Figure 3.

4 Connecting solutions to initial conditions

In this final section we give information on those asymptotic solutions having prescribed initial conditions and conversely. In so doing we will be in a position to describe the asymptotic behavior of all solutions in a plane of initial conditions (see below). It is

best to refer to Theorem 6 in the sequel. We recall that $y_0 \equiv y(0)$, $y'_0 \equiv y'(0)$ where all the initial conditions are assumed real. In addition, $I \equiv I_0$ is defined as in (6), that is,

$$I = \frac{1}{2}y'_0{}^2 + \frac{1}{8}y_0^2 + \ln |1 + y_0|. \quad (17)$$

The following result gives a sufficient condition for the existence of solutions with finite singularities on $[0, \infty)$.

Lemma 12. Let $I \leq 0$. Then the corresponding solution of (1) has a point where its second derivative is infinite.

Proof. First we note that the region defined by (17) subject to $I \leq 0$ is contained in the strip $-1 < y_0 \leq 0$, of the (y_0, y'_0) -plane. If $y'_0 \leq 0$ then, by Lemma 9, the solution has a singularity as stated.

On the other hand, if $y'_0 > 0$ then $I < 0$. Now, use of Lemma 2 shows that any corresponding solution of (1) must be negative on every interval to the right of $x = 0$ in which it is defined. Hence by combining Lemma 8 and Theorem 6 we deduce that y cannot be of type (1) or (2) (since then the solution must be positive somewhere to the right of zero and this contradicts the stipulated negativity above). \square

We have seen in Theorem 8, Theorem 10 and Corollary 5 that asymptotically linear solutions exist and, in fact, prevail when we assume that the solution is not of type (4). In this spirit, we provide some theoretical (and so exact) relationships between the solution asymptotics and the corresponding initial conditions.

Lemma 13. Let y be a solution of (1) that is asymptotically linear, viz., for some $a, b \in \mathbb{R}$, $y(x) \sim ax + b$ as $x \rightarrow \infty$, where $a = 0$ is allowed. Then

$$y_0 = b - \int_0^\infty t F(t, y(t)) dt \quad (18)$$

and

$$y'_0 = a + \int_0^\infty F(t, y(t)) dt. \quad (19)$$

Conversely, if (y_0, y'_0) are initial conditions leading to an asymptotically linear solution $y(x)$, then the corresponding constants a, b are given above.

Proof. A double integration of (1) along with Fubini's Theorem gives, for any $x \geq 0$,

$$y(x) = y_0 + y'_0 x - \int_0^x (x - t) F(t, y(t)) dt.$$

On the other hand, since this same solution is asymptotically linear we have $y(x) \rightarrow ax + b$ and $y'(x) \rightarrow a$ as $x \rightarrow \infty$. Hence, by the results in the previous section $y(x)$ also satisfies the integral equation

$$y(x) = b + ax - \int_x^\infty (t - x) F(t, y(t)) dt, \quad x \geq 0.$$

Since these last two equations must be identical we can differentiate them and thereby find (19). To get (18) we set $x = 0$ in both of them. The converse is clear. \square

Notation: Next, let \mathcal{S}_0 be the subset of all initial values y_0 where $-1 < y_0 < \infty$, such that there exists a value of $y'_0 > 0$ whose corresponding solution is asymptotically constant.

We undertake a study of this set \mathcal{S}_0 in the sequel. A combination of Lemma 7, Lemma 8 and Theorem 7 in that order indicates that $\mathcal{S}_0 \neq \phi$. As a result, those initial values with $y_0 > -1$ (resp. $y_0 < -1$) leading to asymptotically constant solutions of (1) lie in the strip $y_0 > -1, y'_0 > 0$ (resp. $y_0 < -1, y'_0 < 0$) of the (y_0, y'_0) -plane.

It is interesting to note that if $y_0 = -1$ is a limit point of \mathcal{S}_0 then $y'_0 \rightarrow +\infty$. That is, the closer we are to the singularity the higher we have to *shoot* in order to get asymptotically constant solutions (otherwise the solutions “fall back” and become solutions having points where the second derivative is infinite). This is the content of the next result.

Lemma 14. Let $y_0^n \in \mathcal{S}_0$ be a sequence of initial values with $y_0^n \rightarrow -1$ as $n \rightarrow \infty$. Then the corresponding values $y_0'^n \rightarrow +\infty$ as $n \rightarrow \infty$.

Proof. Let $(y_0^n, y_0'^n)$ be such an initial condition and $y_n(x)$ be the corresponding solution. Refer to (5) with y replaced by y_n and $a = 0$. Note that since $y(x)$ is asymptotically constant and $y'(x) \rightarrow 0$, as $x \rightarrow \infty$, the left side of (5) is finite. However, the right side of (5) is unbounded unless $(y_0'^n)^2 = +\infty$. The result follows. \square

Lemma 15. Let $y(x)$ be any asymptotically linear solution of (1). Then $y \in C^\infty(0, \infty)$ and there is an expansion of y into a series of initial conditions $y(x) = \sum_{n=0}^{\infty} c_n(y_0, y'_0) x^n$ where the coefficients are functions of y_0, y'_0 alone and the expansion is absolutely and uniformly convergent for x in compact subsets of $[0, \infty)$.

Proof. Since F is analytic for $t \in \mathbb{C}$ and also for $y \in \mathbb{C} \setminus \{-1\}$ (the complex plane punctured at $y = -1$), it follows from standard theorems (cf., Hille [[15], Theorem 2.5.2]) that every asymptotically linear solution of (1) defined by initial conditions y_0, y'_0 where $y_0 \neq -1$ is necessarily complex analytic and so infinitely differentiable for all $x > 0$. In addition,

$$y(x) = \sum_{n=0}^{\infty} c_n(y_0, y'_0) x^n,$$

where the power series converges absolutely and uniformly for x in compact subsets of $[0, \infty)$, and the coefficients are functions of the initial conditions (y_0, y'_0) . The coefficients are easily found by appealing to the differential equation (1) and differentiating as required since $y \in C^\infty(0, \infty)$. Indeed, its first few terms are given by $c_n \equiv c_n(y_0, y'_0)$,

$$\begin{aligned} c_0 &= y_0 \\ c_1 &= y'_0 \\ c_2 &= -\frac{1}{8} \left(y_0 + \frac{4}{1 + y_0} \right), \end{aligned} \tag{20}$$

$$\begin{aligned}
c_3 &= \frac{1}{24} \left(y_0 + \frac{8}{1+y_0} + y_0' \frac{(3+y_0)(1-y_0)}{(1+y_0)^2} \right) \\
c_4 &= -\frac{y_0(127 + 73y_0^2 + 9y_0^3 + 3y_0^4) + y_0'(56 + 40y_0 - 24y_0^2 - 8y_0^3) + 76 + 32y_0'^2}{134(1+y_0)^3} \\
\dots &\quad \dots
\end{aligned} \tag{21}$$

□

Remark 9. Eq.(20) can be used to determine the short-term qualitative behavior of any solution of (1). For example, it is easy to see that if y is any solution of (1) with initial conditions (y_0, y_0') , then for all sufficiently small x for which the solution is defined,

$$\operatorname{sgn} y''(x) = -\operatorname{sgn}(1 + y_0).$$

Thus, we can easily derive from this that the graph of y is concave up (resp. down) whenever $y_0 < -1$ (resp. $y_0 > -1$) if x is small enough.

In order to gain an understanding of the initial conditions leading to asymptotically constant solutions we require an estimate of the approach of such a solution to its limit. This growth is provided by the next result.

Lemma 16. Let y be an asymptotically constant solution of (1) with $y(x) \sim M$, where $M \neq 0$, as $x \rightarrow \infty$. Then, for all sufficiently large x ,

$$|y(x) - M| = O(e^{-x}), \tag{22}$$

$$|y'(x)| = O(e^{-x}). \tag{23}$$

Proof. First, we use an induction argument to show that if $y_n(x)$, $n \geq 1$, denote the successive approximations of $y(x)$, then $F(t, y_n(t)) = O(e^{-t})$ as $t \rightarrow \infty$. Setting $y_1(t) = M$, for every $t \geq 0$, it is easy to see that $F(t, y_1(t)) = O(e^{-t})$ for large t , so the statement is true for $n = 1$. Assume that $F(t, y_i(t)) = O(e^{-t})$, for large t , holds for $i = 2, 3, \dots, n$. Since

$$y_{n+1}(x) = M - \int_x^\infty (t-x)F(t, y_n(t)) dt, \quad n \geq 1, \tag{24}$$

necessarily for every $x \geq 0$, we see that

$$|y_{n+1}(x) - M| \leq \int_x^\infty (t-x)|F(t, y_n(t))| dt \leq A \int_x^\infty (t-x)e^{-t} dt = O(e^{-x}),$$

for all sufficiently large x , where A is some positive constant (which can be made independent of n) whose value is immaterial. Passing to the limit we find (22).

On the other hand, since $y'(x) \rightarrow 0$ as $x \rightarrow \infty$ we also have (cf., see the proof of Lemma 13 above),

$$y'_{n+1}(x) = \int_x^\infty F(t, y_n(t)) dt, \quad n \geq 1. \tag{25}$$

Thus, (23) follows from the estimates on F from the first part upon passage to the limit. □

Next we show that corresponding to any initial value $y_0 > -1$ there is an asymptotically constant solution for some appropriate choice of $y'_0 > 0$.

Theorem 11. *For every $\delta > 0$ and $\delta < 1/2$, there is an asymptotically constant solution y of (1) such that $y(x) \sim -1 + 2\delta$ as $x \rightarrow \infty$. For such a solution, its initial values y_0, y'_0 satisfy $-1 < y_0 < -1 + 2\delta$ and $y'_0 > 0$.*

Proof. Write $M = -1 + 2\delta$ and consider the successive approximations (24)

$$\begin{aligned} y_{n+1}(x) &= M - \int_x^\infty (t-x)F(t, y_n(t)) dt, \quad n \geq 1, \\ &= M - \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t}y_n(t) + \frac{e^{-2t}}{1+y_n(t)} \right) dt. \end{aligned} \quad (26)$$

Choose X_0 so large that

$$\begin{aligned} \frac{1}{4}(3\delta - 1)e^{-x} + \frac{1}{4\delta}e^{-2x} &< \delta \quad \text{and} \\ \frac{1}{4}(1 - \delta)e^{-x} - \frac{1}{12\delta}e^{-2x} &< \delta \quad \text{for all } x > X_0, \end{aligned} \quad (27)$$

and

$$\alpha \equiv \frac{1}{4\delta^2} (e^{-2X_0} + \delta^2 e^{-X_0}) < 1. \quad (28)$$

Define $y_0(x) \equiv -1 + \delta$, for every $x \geq 0$. We now show that for $n = 0, 1, 2, \dots$, we have

$$-1 + \delta \leq y_n(x) \leq -1 + 3\delta \quad (29)$$

for all $x > X_0$. The result is clear for $n = 0$. Assuming (29) true for $n = m$, we estimate y_{m+1} to find

$$\begin{aligned} \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t}(\delta - 1) + \frac{e^{-2t}}{3\delta} \right) dt &\leq \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t}y_m(t) + \frac{e^{-2t}}{1+y_m(t)} \right) dt \\ &\leq \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t}(3\delta - 1) + \frac{e^{-2t}}{\delta} \right) dt, \end{aligned}$$

which gives

$$\frac{1}{4}(\delta - 1)e^{-x} + \frac{e^{-2x}}{12\delta} \leq \int_x^\infty (t-x)F(t, y_m(t)) dt \leq \frac{1}{4}(3\delta - 1)e^{-x} + \frac{e^{-2x}}{4\delta},$$

from which it follows that

$$2\delta - 1 - \frac{1}{4}(3\delta - 1)e^{-x} - \frac{e^{-2x}}{4\delta} \leq y_{m+1}(x) \leq 2\delta - 1 - \frac{1}{4}(\delta - 1)e^{-x} - \frac{e^{-2x}}{12\delta}$$

for every $x > X_0$, and the result follows on account of (27). Thus, (29) is true for every $n \geq 0$ and $x > X_0$.

Next, we estimate successive differences in the approximations. Observe that for $m \geq 0$,

$$\begin{aligned} y_{m+2}(x) - y_{m+1}(x) &= \int_x^\infty (t-x)(F(t, y_m(t)) - F(t, y_{m+1}(t))) dt \\ &= \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t}(y_m(t) - y_{m+1}(t)) + \frac{e^{-2t}(y_{m+1}(t) - y_m(t))}{(1+y_m(t))(1+y_{m+1}(t))} \right) dt. \end{aligned}$$

Consequently,

$$|y_{m+2}(x) - y_{m+1}(x)| \leq \sup_{t>x} |y_{m+1}(t) - y_m(t)| \times \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t} + \frac{e^{-2t}}{(1+y_m(t))(1+y_{m+1}(t))} \right) dt. \quad (30)$$

However, from (29) we have $(1+y_m(t))(1+y_{m+1}(t)) \geq \delta^2$ for every $x > X_0$. Using this in (30), we get the estimate

$$\begin{aligned} |y_{m+2}(x) - y_{m+1}(x)| &\leq \sup_{t>x} |y_{m+1}(t) - y_m(t)| \int_x^\infty (t-x) \left(\frac{1}{4}e^{-t} + \frac{e^{-2t}}{\delta^2} \right) dt \\ &= \sup_{t>x} |y_{m+1}(t) - y_m(t)| \frac{1}{4\delta^2} (e^{-2x} + \delta^2 e^{-x}) \\ &\leq \alpha \sup_{t>x} |y_{m+1}(t) - y_m(t)|, \quad (m \geq 0) \end{aligned} \quad (31)$$

where $0 < \alpha < 1$, provided that $x > X_0$, by (28). Taking the supremum of both sides of (31), we see that this establishes the convergence of the successive approximations, at least for $x > X_0$. The continuity of the limiting function $y(x)$ is also easily proved using these estimates.

Finally, by Lemma 7 and Lemma 9, we must have that for such a solution, $y'_0 > 0$. If $y_0 \geq 0$, then the proof of Lemma 6 shows that $M > 0$, which is false. Thus, $-1 < y_0 < 0$. Arguing as in the proof of Lemma 8, the solution must satisfy $y(x) < 0$, $y'(x) > 0$ and $y''(x) < 0$. But $y'(x)$ has a limit at infinity, and if this limit is positive, then we obtain an asymptotically linear solution, which is impossible. Thus, this limit is zero. However, note that $y'(x) > 0$ throughout its domain, so that $y(x)$ is increasing to its limit. Thus, $-1 < y_0 < -1 + 2\delta$. \square

Remark 10. Combining Theorem 7 and Theorem 11, we obtain that for every initial value $y_0 > -1$, there is a value of $y'_0 > 0$ such that the corresponding solution is asymptotically constant at infinity. A similar result holds for the initial conditions $y_0 < -1$, $y'_0 < 0$ and is left to the reader.

We show presently that the collection of points (y_0, y'_0) in Π which lead to an asymptotically constant solution is, in fact, a piecewise differentiable curve whose graph we can approximate.

Theorem 12. *The set $\mathcal{T}_0 = \{(y_0, y'_0) \mid y_0 \in \mathcal{S}_0\}$ is the graph of a differentiable function on $\mathbb{R} \setminus \{-1\}$.*

Proof. Write $y(x, y_0, y'_0)$ for an asymptotically constant solution of (1) whose initial values are defined by $y(0) = y_0, y'(0) = y'_0$. We take it that $y(x) \sim M$, where $M > 0$, and consider the relation $y(x, y_0, y'_0) - M = 0$ (or $y'(x, y_0, y'_0) = 0$), the idea here being an attempt to approximate this solution. Standard continuous dependence on the initial conditions estimates (e.g., Hille [[15], p.78]) give that $y(x, y_0, y'_0)$ is a differentiable function of (y_0, y'_0) . Fixing x and y_0 for the time being, we deduce that y is a

differentiable function of y'_0 . Since

$$y(x, y_0, y'_0) = y_0 + y'_0 x - \int_0^x (x-t)F(t, y(t, y_0, y'_0)) dt, \quad (32)$$

differentiating both sides with respect to y'_0 we find

$$\frac{\partial y}{\partial y'_0}(x, y_0, y'_0) = x - \int_0^x (x-t) \left(\frac{1}{4}e^{-t} - \frac{e^{-2t}}{(1+y)^2} \right) \frac{\partial y}{\partial y'_0}(t, y_0, y'_0) dt.$$

Thus, $\partial y/\partial y'_0(x, y_0, y'_0)$ cannot vanish identically on $[0, x]$, and so there is a point x_0 and a neighborhood $(x_0 - \delta, x_0 + \delta)$ in which $\partial y/\partial y'_0(x, y_0, y'_0) \neq 0$. The implicit function theorem now implies that $y'_0 = f(x, y_0)$ locally. That is, given x , we see that y'_0 is locally a (differentiable) function of y_0 , so that the set \mathcal{T}_0 is the graph of a function, or this set is a differentiable curve in the plane (y_0, y'_0) of initial conditions. \square

Without loss of generality, we restrict ourselves to the case where $y_0 > -1$ and $y'_0 > 0$ (the case $y_0 < -1$ and $y'_0 < 0$ being similar). From Theorem 12, we see that the collection \mathcal{T}_0 of points (y_0, y'_0) in Π such that $y(x)$ is asymptotically constant at infinity is the graph of a function defined on $(-1, \infty)$. We now show how to estimate this curve.

Use of Lemma 16 and Lemma 13 shows that the curve \mathcal{T}_0 has a vertical tangent line at $x = -1$, while Lemma 15 can be used to estimate this curve as follows: Lemma 16 tells us that for large x , the values of $y'(x)$ are exponentially small. Thus, differentiating the representation $y(x) = \sum_{n=0}^{\infty} c_n(y_0, y'_0) x^n$, with respect to x and setting the resulting series equal to zero for suitable x -ranges we get, solving for y'_0 in terms of y_0 and stopping after the third term,

$$y'_0 = \frac{1}{4} \left(y_0 + \frac{4}{1+y_0} \right) x. \quad (33)$$

For $y_0 > -1$ and $y'_0 > 0$, this curve is displayed below.




Figure 4.

Eq. (33) serves as a rough estimate for the curve defined by the set \mathcal{T}_0 . Sharper estimates can be found by collecting more terms from the power series representation given in Lemma 15 and solving for y'_0 (after setting the differentiated sum equal to zero) and piecing together various arcs obtained by varying the x -values. The resulting curve (with $x = 1$) is shown in Fig. 4. Varying the value of x in (33) can be used to “fit” the curve to solutions obtained numerically. Note that the graph of y'_0 as a function of y_0 (as displayed in Fig. 4) has a vertical tangent line at $y = -1$, in conformity with the theoretical investigations of Lemma 14. As well, we note that initial conditions “above” the curve lead to solutions that are asymptotically linear (not constant), while points below lead to solutions with finite singularities. Finally, we note that this analysis also can be extended to the left plane defined by $y_0 < -1$. In this case, we get another piece of the curve above (also given by (33)), shown below alongside the one in Fig. 4.




Figure 5.

The two branches of the curve in Fig. 5 approximating the set \mathcal{T}_0 have the following property:

Theorem 13. *Given $y_0 \neq -1$, there exists a distinguished value y_0^* of y'_0 (depending on whether $y_0 > -1$ or $y_0 < -1$, resp.) such that for all $y'_0 > y_0^*$ (resp. $y'_0 < y_0^*$) the solutions are asymptotically linear while for $y'_0 < y_0^*$ (resp. $y'_0 > y_0^*$) the solutions have finite singularities. Finally, when $y'_0 = y_0^*$ the solutions are asymptotically constant.*

In order to prove this theorem we need a result that allows us to compare solutions of (1). This is the content of the next lemma.

Lemma 17. Let y, z be any two solutions of (1) such that for some $\delta > 0$, $|1 + y(x)| > \delta$ and $|1 + z(x)| > \delta$ for every $x \in [a, b]$. Then

$$[y'(x)z(x) - y(x)z'(x)] \Big|_a^b = \int_a^b \frac{(y(x) - z(x))(1 + y(x) + z(x))}{(1 + y(x))(1 + z(x))} e^{-2x} dx. \quad (34)$$

Proof. The argument is Sturmian in nature. Multiply (1) by z . With y replaced by z we multiply (1) again by y and subtract the two results. Integrating over $[a, b]$ and simplifying we get

$$\int_a^b \frac{(y(x) - z(x))(1 + y(x) + z(x))}{(1 + y(x))(1 + z(x))} e^{-2x} dx = \int_a^b (y''(x)z(x) - y(x)z''(x)) dx.$$

Integrating the right side by parts there follows (34). □

Proof. (Theorem 12) For a given y_0 , the existence of the special initial slope y_0^* (and corresponding solution, $y^*(x)$), is a consequence of Theorem 12.

Case i: Let $y_0 \geq 0$, a quantity that we now fix, and $y_0' > 0$ (the case $y_0' \leq 0$ immediately leads to type (4) solutions by Lemma 9). We first show that whenever $y_0' > y_0^*$, then the corresponding solution $y(x)$ is asymptotically linear.

For $y_0' > y_0^*$, there exists a neighborhood $[0, \eta]$, $\eta > 0$, such that for the resulting solutions, $y'(x) > y^*(x)$ for all $x \in [0, \eta]$.

Assume that the collection of all such η has no finite upper bound. Since $y^*(x)$ is defined for all x the same is true of $y'(x)$. Thus, $y'(x)$ tends to a limit at infinity (recall that all solutions and their derivatives have limits in this case, cf., Theorem 6). However, this limit cannot be zero at infinity as this would violate Theorem 12 (by creating yet another asymptotically constant solution for the same y_0). Thus, using the proofs of Lemma 6 and Lemma 8, $y'(x)$ tends to a non-zero limit at infinity, and so the solution must be asymptotically linear.

If the collection of these η admits a finite upper bound, then there exists a smallest value $\eta \equiv x_0 > 0$ such that $y'(x_0) = y^*(x_0)$ and $y'(x) > y^*(x)$ for all $x \in [0, x_0]$. Since $y(0) = y^*(0)$ by hypothesis, we deduce from the preceding inequality that $y(x) > y^*(x)$ for every $x \in [0, x_0]$. We now use Lemma 17 above with $z(x) \equiv y^*(x)$ over $[a, b] \equiv [0, x_0]$. Recall that $y'(x_0) = y^*(x_0) = z'(x_0)$ and $y'(0) > y^*(0) = z'(0)$ (by hypothesis). Thus,

$$\begin{aligned} \int_0^{x_0} \frac{(y(x) - z(x))(1 + y(x) + z(x))}{(1 + y(x))(1 + z(x))} e^{-2x} dx &= [y'(x)z(x) - y(x)z'(x)] \Big|_0^{x_0} \\ &= y'(x_0)(z(x_0) - y(x_0)) - y(0)(y'(0) - z'(0)) < 0, \end{aligned} \quad (35)$$

since $z(x_0) - y(x_0) < 0$, $y'(x_0) > 0$ and $y(0)(y'(0) - z'(0)) \geq 0$. This, however, contradicts the fact that the left side is positive since $y(x) > z(x)$ over $[0, x_0]$ and all the other terms are positive. Thus x_0 cannot exist and so we conclude, from the preceding paragraph, that $y(x)$ is asymptotically linear.

Let $y_0' < y_0^*$. We wish to show that y is a type (4) solution. Arguing as in the previous case, on the assumption that $y(x)$ is defined for all x , we cannot have $y'(x)$ tend to zero at infinity. Next, we know that the assumption $y_0' \leq 0$ immediately leads to type (4) solutions by Lemma 9. Hence we need only consider the case where $y_0' > 0$.

Should y not be of type (4), then it must be asymptotically linear (by one of Lemma 6 - Lemma 9, as the case may be, depending on the sign of y_0'). We show presently that this is impossible.

For $y'_0 < y_0^*$, the resulting solution $y'(x) < y'^*(x)$ for all $x \in [0, \eta]$ where $\eta > 0$. Since $y(x)$ is asymptotically linear we have $y'(x) \rightarrow \beta$ as $x \rightarrow \infty$ where $\beta > 0$ (note that $\beta < 0$ leads to a type (4) solution since $y_0 > -1$). Writing $g(x) = y'(x) - y'^*(x)$ for all x , we see that g is differentiable and $g(0) < 0$, by hypothesis. In addition, since $y^*(x)$ is asymptotically constant, for $0 < \varepsilon < \beta/4$ there is an X_0 such that $|y'^*(x)| < \beta/4$ for all $x > X_0$. Furthermore, since $y'(x) \rightarrow \beta$ as $x \rightarrow \infty$, there is an $X_1 > X_0$ such that for every $x > X_1$, $y'(x) > \beta/2$. Hence, for $x > X_1$, we have $g(x) > \beta/2 - \beta/4 = \beta/4 > 0$. But this means that $g(x_0) = 0$ for some $x_0 \in [0, X_1]$, that is, $y'(x_0) = y'^*(x_0)$. Letting x_0 be the smallest such point, if necessary, we can assume that $y'(x) < y'^*(x)$ for all $x \in [0, x_0)$. It follows from this that $y(x) < y^*(x)$ for every $x \in [0, x_0)$. As in the previous case we use Lemma 17 over the interval $[0, x_0]$ with $z(x) \equiv y^*(x)$. In this case we note that the left side of (35) is negative while the right side is positive, leading once again to a contradiction. Hence $y(x)$ cannot be asymptotically linear and so must be of type (4).

Case ii: Finally, let $-1 < y_0 < 0$. The case where $y'_0 \leq 0$ leads to type (4) solutions by Lemma 9 and so we can assume that $y'_0 > 0$.

Let $0 < y'_0 < y_0^*$. One must prove that the resulting solution y is a type (4) solution. As in the previous cases, if $y(x)$ is defined for all x , we cannot have $y'(x) \rightarrow 0$ at infinity. On the other hand, $y'(x) \rightarrow \beta$, and $\beta < 0$ is impossible unless this is actually a type (4) solution. If y is not a type (4) solution, then it follows that $y(x)$ must be asymptotically linear at infinity in the sense that, for some $\beta > 0$, we have $y'(x) \rightarrow \beta$ as $x \rightarrow \infty$. Repeating the discussion above regarding the g -function, there exists a smallest point x_0 such that $y'(x_0) = y'^*(x_0)$. Because of this, one can assume that $y'(x) < y'^*(x)$ for all $x \in [0, x_0)$, and $y'(x) > y'^*(x)$ for all $x \in (x_0, \infty)$ since the derivatives are each monotone functions. So $y(x) > y^*(x)$ for $x > x_0$.

Choose $x_1 > x_0$ so large that $y(x_1) > 0$, $y(x_1) + z(x_1) > 0$, and $y'(x_1) - z'(x_1) > 0$. This is always possible because of the asymptotic nature of these solutions. We now use (34), with $z(x) \equiv y^*(x)$ again, over the interval $[a, b] = [x_1, X]$ where X is chosen so that $y'(X) > \beta/2$. Consider the function $w(x) \equiv 1 + y(x) + z(x)$ for $x \geq x_1$. Since y, z are each monotone, so is w . In addition, $w(x) \geq w(x_1) = 1 + y(x_1) + z(x_1) > 0$, for every $x > x_1$. It now follows that the left-side of (34) is positive while the right-side is negative. This contradiction shows that y is necessarily a type (4) solution. The case $y_0 < -1$ is similar and is left to the reader. \square

4.1 Concluding remarks and open questions

For $x < 0$ it is not difficult to show that every solution of (1) with initial conditions $y_0 > -1, y'_0 > 0$ or $y_0 < -1, y'_0 < 0$ must have a finite singularity. We sketch the proof for completeness. The change of variable $x = -t$, $y(x) = u(t)$ sends (1) on $(-\infty, 0]$

into the equation

$$u'' + \frac{e^t}{4}u + \frac{e^{2t}}{1+u} = 0, \quad t \in [0, \infty).$$

Since we are dealing with solutions having finite singularities, it suffices to consider initial conditions for this equation such that $u_0 > -1, u'_0 < 0$ or $u_0 < -1, u'_0 > 0$ since otherwise we already know that the other initial conditions lead to type (4) solutions for $x > 0$ by Lemmas 7-9 in the discussion above. We now argue as in the proofs of Lemmas 6-9.

Let $u_0 > -1, u'_0 < 0$. Then there is a right-neighborhood of $t = 0$ in which $u(t) < 0, u'(t) < 0$ and $u''(t) < 0$. This then leads to a type (4) solution (as in Lemma 9). On the other hand, if $u_0 < -1, u'_0 > 0$, then there is a right-neighborhood of $t = 0$ such that $u(t) < -1, u'(t) > 0, u''(t) > 0$, and this leads to a type (4) solution (by the methods of Lemma 10). Combining these results we obtain

Theorem 14. *Every solution of (1) on \mathbb{R} has a finite singularity.*

Placing the pole in (1) at $y = -a, a > 0$, leads to no further generality, as this amounts to a rescaling with e^{-2x} replaced by $e^{-2x}/2a$. The techniques here can be adapted to the cases where

$$y'' + f(x)y + \frac{g(x)}{1+y} = 0, \quad x \in [0, \infty),$$

on the assumption that f, g are positive, of exponential order, $f(0) < 4g(0)$, in addition to the usual analyticity requirements. Indeed, more general criteria for a corresponding Theorem 6 can be formulated similar to those described in [10] by restricting the classes of F .

We note the asymptotic behavior of those solutions with finite singularities in Section 2.1. Numerical considerations indicate that this approach to the value -1 is actually very slow, in conformity with the square root of the negative of the logarithm seen above. We also note that this reveals the nature of the logarithmic branch point. A thorough study of this equation for complex x may reveal more interesting facts about the general solution.

References

1. M.J. Ablowitz, P.A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge Univ. Press, Cambridge, UK, 1999.
2. M.J. Ablowitz, A. Ramani and H. Segur, *A connection between nonlinear evolution equations and ordinary differential equations of P-type. I* J. Math. Phys. **21** (1980), 715-721.
3. R.P. Agarwal and D. O'Regan *A note on a singular integral equation arising in a problem in communications* Z. Angew. Math. Mech. **81** (2001), 499-504.

4. R.P. Agarwal and D. O'Regan *Volterra integral equations: the singular case*
Hokkaido Math. J. **32** (2003), 371-381.
5. R.P. Agarwal and D. O'Regan *Fredholm and Volterra integral equations with integrable singularities*
Hokkaido Math. J. **33** (2004), 443-456.
6. C.O. Alves, J.V. Goncalves, and L.A. Maia, *Singular nonlinear elliptic equations in \mathbb{R}^n*
Abstract and Applied Anal. **3** (3-4) (1998), 411-423.
7. J.M. Borwein, A.S. Lewis, and R.D. Nussbaum, *Entropy minimization, DAD problems, and doubly stochastic kernels*
J. Funct. Anal., **123** (2) (1994), 264-307.
8. P.J. Bushell, *Class of Volterra and Fredholm nonlinear integral equations*
Math. Proc. Camb. Phil. Soc. **79** (1976), 329-335.
9. A. Constantin, *Positive solutions of quasilinear elliptic equations*,
J. Math. Anal. Appl. **213** (1997), 334-339.
10. S.G.A. Dubé and A.B. Mingarelli, *Note on a non-oscillation theorem of Atkinson*
Electronic J. Differential Equations, Vol. 2004 (22) (2004), 1-6.
11. S.M. Gomes, *On a singular nonlinear elliptic problem*
SIAM J. Math. Anal. **17** (6) (1986), 1359-1369.
12. J.K. Hale, *Ordinary Differential Equations*
Wiley Interscience Publishers, (New York, 1969).
13. J.K. Hale and N. Onuchic *On the asymptotic behavior of solutions of a class of differential equations*
Contributions to Differential Equations **1** (1963), 61-75.
14. P. Hartman and N. Onuchic *On the asymptotic integration of ordinary differential equations*
Pacific J. Math. **13** (1963), 1193-1207.
15. E. Hille, *Lectures on Ordinary Differential Equations*
Addison-Wesley, (New York, 1969).
16. E.L. Ince, *Ordinary Differential Equations*
Dover Publications, (New York, 1956).
17. Z.R. Jin, *Solutions for a class of singular semilinear elliptic equations*
Nonlinear Anal., **31** (3-4) (1998), 475-492.
18. S. Karlin and L. Nirenberg, *On a theorem of P. Nowosad*
J. Math. Anal. Appl. **17** (1967), 61-67.

19. A.G. Kartsatos, *Recent results on oscillation of solutions of forced and perturbed nonlinear differential equations of even order*
in *Stability of Dynamical Systems- Theory and Applications*, J. R. Graef ed., Lecture Notes in Pure and Applied Mathematics Vol. 28, Marcel Dekker, New York, 1977, 17-72.
20. C.A. Michelli and P. Olsen, *Penalized maximum-likelihood estimation, the Baum-Welch algorithm, diagonal balancing of symmetric matrices and applications to training acoustic data*
J. Comp. Applied Math., **119** (1-2) (2000), 301-331
21. A.H. Nasr, *Necessary and sufficient conditions for the oscillation of forced nonlinear second order differential equations with delayed argument*,
J. Math. Anal. Appl. **212** (1997), 51-59.
22. P. Nowosad, *On the integral equation $\kappa f = 1/f$ arising in a problem in communication*
J. Math. Anal. Appl. **14** (1966), 484-492.
23. R.D. Nussbaum, *Iterated nonlinear maps and Hilbert projective metric, 2.*
Mem. Amer. Math. Soc., **79** (401) (1989), iv + 118 pp.
24. R.D. Nussbaum, *Entropy minimization, Hilbert projective metric, and scaling integral-kernels*
J. Funct. Anal., **115** (1) (1993), 45-99.
25. R.D. Nussbaum, *Hilbert's projective metric and iterated nonlinear maps*
Mem. Amer. Math. Soc., **75** (391) (1988), iv+137 pp.
26. G.E. Parker and T.J. Walters, *Constructing solutions to $\int_0^1 f(j)A(j, x)/f(x)dj = \int_0^1 f(x)A(x, j)/f(j) dj$*
SIAM J. Math. Anal. **13** (5) (1982), 856-865.
27. C.A. Swanson, *Comparison and Oscillation Theory of Linear Differential Equations*
Academic Press, New York, 1968.
28. Z. Yin, *Monotone positive solutions of second order nonlinear differential equations*,
Nonlinear Analysis, **54** (2003), 391-403.
29. T.J. Walters and G.E. Parker, *Positive nonlinear integral equations with reciprocals of the solution in the integrand*
Nonlinear Analysis, **5** (11) (1981), 1163-1172.
30. J. Wang, W. Gao, and Z. Zhang, *Singular nonlinear boundary value problems arising in boundary layer theory*
J. Math. Anal. Appl. **233** (1999), 246-256.

31. Z. Zhao, *Positive solutions of nonlinear second order ordinary differential equations*,
Proc. Amer. Math. Soc., **121** (2) (1994), 465-469.

(1). (Corresponding author)
School of Mathematics and Statistics,
Carleton University,
Ottawa, Ontario, Canada, K1S 5B6
e-mail: amingare@math.carleton.ca,
Tel/Fax: 001 (613) 520-3534.

(2). Departamento de Matemáticas,
Universidad de Las Palmas de Gran Canaria,
Campus de Tafira Baja,
35017 Las Palmas de Gran Canaria, Spain.
e-mail: pacheco@dma.ulpgc.es .

(3). School of Mathematics and Statistics,
Carleton University,
Ottawa, Ontario, Canada, K1S 5B6
e-mail: melkonian@math.carleton.ca.