

## NOTE

# Influence of fishing and functional group of algae on sea urchin control of algal communities in the eastern Atlantic

Fernando Tuya<sup>1,\*</sup>, Pablo Sanchez-Jerez<sup>2</sup>, Ricardo J. Haroun<sup>1</sup>

<sup>1</sup>Department of Biology, BIOGES, Campus Tafira, University of Las Palmas de G.C., 35017 Las Palmas, Canary Islands, Spain

<sup>2</sup>Marine Biology Laboratory, University of Alicante, PO Box 99, 03080 Alicante, Spain

**ABSTRACT:** We address the effect of several factors influencing the depth of interfaces between the shallow photophilic brown algal assemblages and *Diadema antillarum* urchin-dominated barrens on rocky reefs of the warm-temperate central-eastern Atlantic. By means of a multifactorial observational approach, we hypothesized that it would be locally influenced by (1) the functional structure of the dominant brown algal assemblages (corticated foliose algae versus large frondose coarsely-branched furoid species), in addition to (2) the fishing status (fished versus unfished sites) and (3) the large-scale topography of the substrate. Our results show that brown algal assemblages reach deeper waters at unfished sites, which is independent of the dominant brown algal functional group. However, monospecific algal facies dominated by large frondose coarsely branched furoid species (mainly the genera *Cystoseira*) reach deeper waters in comparison with corticated foliose algae, at fished sites.

**KEY WORDS:** Sea urchins · *Diadema antillarum* · Phytobenthos · Functional groups · Ecosystem-level effects · Canary Islands

Resale or republication not permitted without written consent of the publisher

## INTRODUCTION

Large, shallow, rocky areas covered by erect photophilic algae are transformed by sea urchins into overgrazed substrates dominated by encrusting coralline algae. These so-called 'urchin-grazed barrens' (e.g. Mann 1982, McClanahan & Sala 1997, Sala et al. 1998, Shears & Babcock 2003, Tuya et al. 2004a) have been widely described in coastal temperate areas (Vukovic 1982, Verlaque 1987, Andrew & Underwood 1989, Francour 1994, Sala & Zabala 1996, Sala et al. 1998, Babcock et al. 1999, Pinnegar et al. 2000, Alves et al. 2001, Shears & Babcock 2003, Tuya et al. 2004a). These authors have highlighted that the transformation of a zone from an algal-dominated system to an urchin-grazed barren is a dynamic and complex interaction of different physical and ecological mechanisms, operat-

ing at different scales of spatial and temporal variability. Processes affecting the sea urchin population structure, and thus involved in the interactions of urchin barrens with algal assemblages, are widely acknowledged in temperate coastal areas (see Shears & Babcock 2003). Fishing pressure on natural sea urchin predators, the topography and complexity of the substrate and the availability of refuges, recruitment, pollution and diseases, harvesting, and the variability of oceanographic events are the most important processes (for reviews, see Sala et al. 1998 and Pinnegar et al. 2000).

Patches of functionally (sensu Steneck & Dethier 1994) different brown algal species form a photophilic macroscopic algal bed, as a continuous and heterogeneous mosaic of different algal species, throughout the shallowest rocky seascape of the Canarian Archi-

\*Email: fernando@canariaporunacostaviva.org

pelago (Haroun et al. 2003). The long-spined sea urchin *Diadema antillarum* (Philippi) has been recently identified as the main key-herbivorous species ('habitat determiner') throughout the sublittoral rocky bottoms of the warm-temperate central-eastern Atlantic, as it controls the macrobenthic algal assemblages with negligible contribution from other grazing species (Alves et al. 2003, Tuya et al. 2004b). Intensive grazing by *D. antillarum* restricts the lower edge of the photophilic algal facies to the first few meters of the subtidal (Alves et al. 2001, 2003, Tuya et al. 2004b). Hence, clear interfaces or borders (sensu Gagnon et al. 2004) between brown algal stands and urchin barrens extend parallel to the coastline. Several ecological and physical processes play a significant role in the structure and abundance of *D. antillarum* populations and, consequently, in the interaction between both habitats across the eastern Atlantic; these are water turbulence (Alves et al. 2001), fishing and the abundance of top-predators (following the classical trophic cascades paradigm, Tuya et al. 2004a), and the small-scale substrate complexity (Tuya et al. 2004a). However, the dominance of photophilic algal stands by functionally different brown algal groups, as determinants of the interplay between photophilic algal assemblages and urchin barrens, still remains to be considered.

In this paper, we address the effect of several factors influencing interfaces between the photophilic algal assemblages and urchin barrens on rocky reefs of the warm-temperate eastern Atlantic by means of a multifactorial, observational (mensurative) approach. Specifically, we hypothesized that the depth-limit between both habitats would be locally influenced by the functional structure of the dominant brown algal assemblages in addition to factors that seem, as reported in the literature, to affect the population structure of the sea urchin *Diadema antillarum*, such as the fishing pressure (unfished versus fished sites) and the physical topography of rocky reefs.

## MATERIALS AND METHODS

**Area of study and experimental design.** The study was carried out on basaltic rocky bottoms between 3 and 30 m depth of the Canary Archipelago (28° N, 15° W, central-eastern Atlantic Ocean), in October 2003. The distribution of benthic communities along the bathymetric axis shows a clear vertical zonation pattern, with the arrangement of the communities in belts with distinct species composition and abundance, as observed in many shallow temperate subtidal habitats (Garrabou et al. 2002 and references therein). Within the shallowest photophilic zone, extensive

stands of perennial brown algal assemblages make up the dominant community (Haroun et al. 2003). Two main groups of photophilic, non-turfing, fleshy brown algae can be distinguished (Haroun et al. 2003), according to the classical algal functional form groups reported in the literature (Littler & Littler 1984, Steeneck & Dethier 1994, Ruitton et al. 2000, Garrabou et al. 2002, McClanahan et al. 2003), occurring as distinct patches. Corticated foliose brown algae (henceforth referred to as Functional Group 1 [FG1]) consist of sheet-shaped, jointed calcareous and thick leathery-shaped species (e.g. *Cutleria multifida*, *Dyctiota* spp., *Lobophora variegata*, *Padina pavonica*, *Stypopodium* spp., *Taonia* spp., *Zonaria* spp. etc.), from 1 to 15 cm in height, which constitute either algal cushions or thin sheets with mixtures of algal species. On the other hand, corticated large brown macrophytes (henceforth referred to as Functional Group 2 [FG2]) are erect, frondose, coarsely branched fucoid species (mainly of the genera *Cystoseira* and *Sargassum*), which generally form monospecific algal stands.


We define the variable  $Z_{<5\%}$  as the depth-limit (the 'edge') from which the percent (%) cover of the overall photophilic brown phytobenthos drops uniformly below 5% cover with increasing depth. This depth, therefore, represents the shallowest upper edge of grazed urchin barrens. This measure was, in all cases, determined on a homogeneous horizontal surface to avoid the local effect of small-scale substrate topography (crevices, holes etc.).

Two categories of large-scale habitat complexity were defined following the criteria of Garcia-Charton et al. (2000): (1) low (rocky platforms with low relief) and (2) high (rocky bottoms with sharp slopes). Finally, fished and unfished sites were selected outside and inside Marine Protected Areas (MPAs) from the Canary Archipelago ([http://www.mapya.es/rmarinas/index\\_rm.htm](http://www.mapya.es/rmarinas/index_rm.htm)), where only capture of fin fish is controlled. On the other hand, sea urchins are not subjected to a fishery across the overall Canary Archipelago.

Our experimental design tested the effect of the 2 functional groups of brown algae (FG1 versus FG2) on the dependent variable  $Z_{<5\%}$  at both fished and unfished sites, with high and low bottom complexity. We selected a total of 24 sites across the Canary Archipelago as spatial replicates of the 8 defined experimental treatments (2 functional groups of brown algae  $\times$  2 categories of fishing pressure  $\times$  2 categories of bottom complexity), with 3 sites per treatment (each MPA provided 4 sites, while unprotected adjacent areas to each MPA another 4 sites).

**Sampling and data analysis.** At each site, a diver swam perpendicular to the coastline until a clear transition between the shallow photophilic brown algal

Fig. 1. *Diadema antillarum*. Non-linear regression between the mean density of total individuals per site and the depth-limit from which the percent cover of the overall brown phyto-benthos drops uniformly below 5% of cover ( $Z_{<5\%}$ ). Empty icons: Functional Group 1; full icons: Functional Group 2;  $\square$ ■: sites with low bottom complexity inside MPAs;  $\diamond$ ◆: sites with high bottom complexity inside MPAs;  $\circ$ ●: sites with low bottom complexity outside MPAs;  $\triangle$ ▲: sites with high bottom complexity outside MPAs



band and the urchin barren was observed, avoiding bare haloes around the shallowest shelters of sea urchins. To ensure that the percent cover of fleshy brown algae was under 5%, we deployed a  $25 \times 25$  cm quadrat and the percent cover was visually estimated *in situ* following standardized procedures (Foster et al. 1991, Dethier et al. 1993, McClanahan et al. 2003). This protocol was replicated 5 times per site ( $n = 5$ ). Tidal corrections were further applied to compare the overall dataset. In addition, the abundance of all adult *Diadema antillarum* individuals (test diameter without spines  $>2$  cm) were visually recorded by SCUBA divers along urchin barrens using 16 haphazardly located  $2 \times 2$  m ( $4 \text{ m}^2$ ) quadrats per sampling site (Ruitton et al. 2000, Tuya et al. 2004a).

The relationship between the lower edge of the photophilic algal assemblages (quantified by means of the  $Z_{<5\%}$ ) and *Diadema antillarum* sea urchin abundance was estimated using non-linear regression analysis.

A mixed 4-factor ANOVA model (Underwood 1997) was applied to test for differences attributable to the 3 defined experimental factors: (1) fishing status (fixed factor with 2 levels: unfished versus fished), (2) the complexity of the habitat (orthogonal and fixed factor with 'low versus high' as levels) and (3) the dominant functional group of algae (orthogonal and fixed factor incorporating the levels corresponding to the 2 categories of algae: FG1 and FG2). The random factor 'Sites' represents the spatial replication of the experimental treatments, and is therefore nested within the second-order interaction term among the 3 experimental factors. When a term of the ANOVA that included a fixed factor was significant, the SNK *a posteriori* test was applied. Before analysis, Cochran's test was used to check for homogeneity of variances. Although no

transformation rendered variances homogeneous, the ANOVA was carried out as it is robust to heterogeneity of variances, particularly for large balanced experiments (Underwood 1997).

## RESULTS

Grazing by the sea urchin *Diadema antillarum* determines the lower edge of photophilic brown algal assemblages, as a strong non-linear decrease with depth (as absolute values of the variable  $Z_{<5\%}$ ) is observed with increasing sea urchin abundance (Fig. 1).

Clear overall differences in the mean depth of interfaces between photophilic algal assemblages and urchin barrens were observed between fished and unfished (MPAs) sites (Figs. 1 & 2), which was corroborated by the ANOVA model (Table 1). The mean value ( $\pm$ SD) of the variable  $Z_{<5\%}$  across unfished sites




Fig. 2. Effect of fishing status (fished versus unfished MPAs) and the functional group of dominant algae (FG1 versus FG2) on interfaces between photophilic brown algal stands and urchin barrens. Error bars represent  $\pm$ SE

Table 1. 4-factor ANOVA on the effects of fishing status, bottom complexity and functional grouping of dominant brown algal assemblages on the mean depth of interfaces between brown algal stands and urchin-dominated barrens across the Canary Archipelago. \* $p < 0.01$

| Source of variation             | df | MS    | F      |
|---------------------------------|----|-------|--------|
| Functional group (=FG)          | 1  | 3.85  | 5.98   |
| Status (=S)                     | 1  | 39.93 | 61.94* |
| Habitat complexity (=C)         | 1  | 0.44  | 0.69   |
| FG $\times$ S                   | 1  | 5.97  | 9.32*  |
| FG $\times$ C                   | 1  | 0.29  | 0.45   |
| S $\times$ C                    | 1  | 0.01  | 0.02   |
| FG $\times$ S $\times$ C        | 1  | 2.07  | 3.21   |
| Site (FG $\times$ S $\times$ C) | 16 | 0.64  | 11.55* |
| Residual                        | 96 |       |        |

( $20.26 \pm 3.66$ ) is, thus, about 2 to 3 times that at fished sites ( $7.65 \pm 5.10$ ) (Fig. 2). At the unfished sites, no significant differences were detected as a function of the dominance of either functional group of brown algae (FG1 versus FG2). However, significant differences were detected, at fished sites by the SNK test between both of the functional groups (Figs. 1 & 2). In fact, this difference detected solely at the fished sites implied a significant first-order interaction between the factors 'Functional group  $\times$  Status, as detected by the ANOVA ( $F = 9.32$ ,  $p < 0.01$ ; Table 1). Brown algal stands consisting of species of the FG2 (mainly monospecific beds of the genera *Cystoseira*) reached deeper waters at the fished sites ( $10.3 \pm 5.37$  m, mean  $\pm$  SD) in comparison with the brown algae belonging to the FG1 ( $5.03 \pm 3.14$  m) (Fig. 1).

The large-scale substrate complexity in the studied sites appears not to play a significant role in the depth of borders between algal stands and urchin barrens. Thus, neither any detectable interaction with other experimental factors nor the main effect of this factor has been involved in any significant term in the ANOVA (Table 1). Finally, it is important to recognize that differences between sites for each treatment explain a significant amount of the variability within our experimental design ( $F = 11.55$ ,  $p < 0.01$ ; Table 1). This inter-site variability within each established treatment is greater across fished sites (right side of Fig. 1) than across unfished sites (left side of Fig. 1).

## DISCUSSION

Depth-related trends in community structure respond to strong changes in community determining factors (Garrabou et al. 2002). Human-induced disturbances through fishing appear to be a significant factor

regulating the structure of shallow phytobenthic communities at opposite levels of fishing pressure (Shears & Babcock 2003 and references therein). Our study has shown that fished sites present shallow interfaces between photophilic brown algal stands and urchin-dominated barrens, whereas this border is deeper in sites protected from fishing (sites inside MPAs). This result is not surprising as sites protected from fishing activities show diverse and highly structured assemblages of large-sized predatory benthophagous organisms (e.g. fish or crustaceans). These populations have the potential of controlling the key-grazing species *Diadema antillarum* throughout infralittoral rocky reefs of the warm-temperate central-eastern Atlantic (Tuya et al. 2004a). This trend has been similarly reported with other key-herbivorous sea urchins at other coastal systems in a large body of scientific literature covering the cold-temperate mid-latitudes in both of the hemispheres (e.g. Andrew & Choat 1982, Bernstein et al. 1983, Sala & Zabala 1996, Estes et al. 1998, Sala et al. 1998, Babcock et al. 1999, Steneck et al. 2002, Shears & Babcock 2003) by following the classical trophic cascades paradigm (as reviewed by Sala et al. 1998, Pinnegar et al. 2000).

Despite the fact that habitat complexity favours the impact of grazers (e.g. sea urchins) governing the structure of erect brown macroalgae (as reviewed by Sala et al. 1998 and Steneck et al. 2002), large-scale ( $\geq 100$  m) habitat complexity does not seem to have an important effect on the depth of the interfaces between brown algal facies and urchin barrens. This lack of significance is attributable to the selection of a descriptor operating at large spatial-scales ( $\geq 100$  m) instead of medium (1 to 10 m) or small ( $< 1$  m) spatial-scales, which would more readily reflect the availability of refuges (crevices, boulders etc.). Marine organisms respond to variations of different descriptors according to their body size and life characteristics (Garcia-Charton et al. 2000).

Unfished sites have shown brown algal stands with a depth-related extension which is independent of the dominant brown algal functional group. In contrast, the functional group to which the dominant brown algal species belong (FG1 versus FG2) may play a significant role in determining the depth of interfaces between shallow photophilic brown algal assemblages and urchin-grazed barrens, at fished sites. Algal stands, dominated by large frondose coarsely branched furoid species (FG2, mainly 4 and 2 species of the genera *Cystoseira* and *Sargassum*, respectively), significantly reached deeper waters in comparison with corticated foliose brown algae (FG1). We propose 2 complementary mechanisms to explain this observation. Firstly, it can be related to the wave-induced sweeping motion of large erect canopy-forming furoid

algae (FG2); algal stands of this functional group restrict both sea urchin movement and suitable space to graze, as has been suggested for large brown macrophytes in temperate waters (Steneck & Dethier 1994, Konar 2000, Gagnon et al. 2004). Secondly, the sea urchin *Diadema antillarum* has shown lower feeding preference for species belonging to FG2 (e.g. *Cystoseira*) than those from FG1 (Tuya et al. 2001). This avoidance is attributable to the rather coarse and tough body of the genera *Cystoseira* (physical factor), as well as to the presence of polyphenols (chemical factor; Winter & Estes 1992, Knoess & Glombitza 1993) as defenses against herbivores (Shunula & Ndibalema 1986). Despite the potentially confounding effects of chemical and morphological traits (Hay 1997), this hierarchy is in agreement with the susceptibility to being eaten as a function of algal morphology (algal functional groups) reported by Steneck & Dethier (1994), mainly for large invertebrates such as sea urchins in temperate seas.

**Acknowledgements.** This research was financially supported by the Spanish 'Ministerio de Medio Ambiente' within the framework of the 'Canarias, por una costa viva' project ([www.canariasporunacostaviva.org](http://www.canariasporunacostaviva.org)). We gratefully thank A. Boyra, E. Falcón, O. Bergasa, T. Sánchez, A. Iglesias, A. López, N. Rodríguez, A. Del Rosario, G. Herrera, R. Herrera and F. Espino for helping us with the underwater data collection and processing. We thank J. M. Fernandez-Palacios for ideas and comments on a first draft of this manuscript.

#### LITERATURE CITED

- Alves FMA, Chicharo LM, Serrao E, Abreu AD (2001) Algal cover and sea-urchin spatial distribution at Madeira Island (NE Atlantic). *Sci Mar* 65:383–392
- Alves FMA, Chicharo LM, Serrao E, Abreu AD (2003) Grazing by *Diadema antillarum* (Philippi) upon algal communities on rocky substrates. *Sci Mar* 67:307–311
- Andrew NL, Choat JH (1982) The influence of predation and conspecific adults on the survivorship of juvenile *Evechinus chloroticus* (Echinoidea: Echinometridae). *Oecologia* 54:80–87
- Andrew NL, Underwood AJ (1989) Patterns of abundance of the sea urchin *Centrostephanus rodgersii* (Agassiz) on the central coast of New South Wales, Australia. *J Exp Mar Biol Ecol* 131:61–80
- Babcock RC, Kelly S, Shears NT, Walker JW, Willis TJ (1999) Changes in community structure in temperate marine reserves. *Mar Ecol Prog Ser* 189:125–134
- Bernstein BB, Schroeter SC, Mann KH (1983) Sea urchin (*Strongylocentrotus droebachiensis*) aggregating behaviour investigated by a subtidal multifactorial experiment. *Can J Fish Aquat Sci* 40:1975–1986
- Dethier MN, Grahlan ES, Cohen S, Tear LM (1993) Visual versus random-point percent cover estimations: 'objective' is not always better. *Mar Ecol Prog Ser* 96:93–100
- Estes JA, Tinker MT, Williams TM, Doak DF (1998) Killer whale predation on sea otters linking oceanic and nearshore ecosystems. *Science* 282:473–476
- Foster MS, Harrold C, Hardin DD (1991) Point vs. photo quadrat estimates of the cover of sessile marine organisms. *J Exp Mar Biol Ecol* 146:193–203
- Francour P (1994) Pluriannual analysis of the reserve effect on fish community in the Scandola natural reserve (Corsica, North-western Mediterranean). *Oceanol Acta* 17:309–317
- Gagnon P, Himmelman JH, Johnson LE (2004) Temporal variation in community interfaces: kelp-bed boundary dynamics adjacent to persistent urchin barrens. *Mar Biol* 144:1191–1203
- García-Charón JA, Pérez-Ruzafa A, Marcos C (2000) Habitat structure and cascade effects. In: Goñi R, Harmelin-Vivien M, Badalamenti F, Le Direach L, Bernard G (eds) *Introductory guide to methods for selected ecological studies in marine reserves*. GIS Posidonie, Marseille, p 94–98
- Garrabou J, Ballesteros E, Zabala M (2002) Structure and dynamics of North-western Mediterranean rocky benthic communities along a depth gradient. *Estuar Coast Shelf Sci* 55:493–508
- Haroun R, Gil-Rodríguez MC, Wildpret de la Torre W (2003) *Plantas marinas de Canarias*. Canseco Press, Talavera
- Hay ME (1997) The ecology and evolution of seaweed-herbivore interactions on coral reefs. *Coral Reefs* 16: S67–S76
- Knoess W, Glombitza KW (1993) A phenolsulphatase from the marine brown alga *Cystoseira tamariscifolia*. *Phytochemistry* 32:1119–1123
- Konar B (2000) Seasonal inhibitory effects of marine plants on sea urchins: structuring communities the algal way. *Oecologia* 125:208–217
- Littler MM, Littler DS (1984) Models of tropical reef biogenesis: the contribution of algae. *Prog Phycol Res* 3:323–364
- Mann KH (1982) Kelp, sea-urchins and predators: a review of strong interactions in rocky subtidal systems of eastern Canada, 1970–1980. *Neth J Sea Res* 16:414–423
- McClanahan TR, Sala E (1997) A Mediterranean rocky-bottom ecosystem fisheries model. *Ecol Model* 104:145–165
- McClanahan TR, Sala E, Stickels PA, Cokos BA, Baker AC, Starger CJ, Jones SH IV (2003) Interaction between nutrients and herbivory in controlling algal communities and coral condition on Glover's Reef, Belize. *Mar Ecol Prog Ser* 261:135–147
- Pinnegar JK, Polunin NVC, Francour P, Badalamenti F and 7 others (2000) Trophic cascades in benthic marine ecosystems: lessons for fisheries and protected-area management. *Environ Conserv* 27:179–200
- Ruitton S, Francour P, Boudouresque CF (2000) Relationships between algae, benthic herbivorous invertebrates and fishes in rocky sublittoral communities of a temperate sea (Mediterranean). *Estuar Coast Shelf Sci* 50:217–230
- Sala E, Zabala M (1996) Fish predation and the structure of the sea urchin *Paracentrotus lividus* population in the NW Mediterranean. *Mar Ecol Prog Ser* 140:71–81
- Sala E, Boudouresque CF, Harmelin-Vivien ML (1998) Fishing, trophic cascades, and the structure of algal assemblages: evaluation of an old but untested paradigm. *Oikos* 83:425–439
- Shears NT, Babcock RC (2003) Continuing trophic cascade effects after 25 years of no-take marine reserve protection. *Mar Ecol Prog Ser* 246:1–16
- Shunula JP, Ndibalema V (1986) Grazing preferences of *Diadema setosum* and *Heliocidaris erythrogramma* (Echinoderms) on an assortment of marine algae. *Aquat Bot* 25:91–95

- Steneck RS, Dethier MN (1994) A functional group approach to the structure of algal-dominated communities. *Oikos* 69:476–498
- Steneck RS, Graham MH, Bourque BJ, Corbett D, Erlandson JM, Estes JA, Tegner MT (2002) Kelp forest ecosystems: biodiversity, stability, resilience and future. *Environ Conserv* 29:436–459
- Tuya F, Martin JA, Reuss GM, Luque A (2001) Feeding preferences of the sea urchin *Diadema antillarum* in Gran Canaria Island (Central-East Atlantic Ocean). *J Mar Biol Assoc UK* 81:1–5
- Tuya F, Boyra A, Sanchez-Jerez P, Barbera C, Haroun RJ (2004a) Relationships between rocky-reef fish assemblages, the sea urchin *Diadema antillarum* and macroalgae throughout the Canarian Archipelago. *Mar Ecol Prog Ser* 278:157–169
- Tuya F, Boyra A, Sanchez-Jerez P, Barbera C, Haroun RJ (2004b) Can one species determine the structure of the benthic community on a temperate rocky reef? The case of the long-spined sea-urchin *Diadema antillarum* (Echinodermata: Echinoidea) in the eastern Atlantic. *Hydrobiologia* 519:211–214
- Underwood AJ (1997) Experiments in ecology: their logical design and interpretation using analysis of variance. Cambridge University Press, Cambridge
- Verlaque M (1987) Relations entre *Paracentrotus lividus* (Lamarck) et le phytobenthos de Méditerranée occidentale. In: Boudouresque CF (ed) Colloque international sur *Paracentrotus lividus* et les oursins comestibles. GIS Posidonie, Marseille, p 5–36
- Vukovic A (1982) Florofaunistic changes in the infralittoral zone after the sea urchin *Paracentrotus lividus* (L.) population explosion. *Acta Adriat* 23:237–241
- Winter FC, Estes JA (1992) Experimental evidence for the effects of polyphenolic compounds from *Dictyonium californicum* Ruprecht (Phaeophyta: Laminaridae) in feeding rate and growth in the red abalone *Haliotis rufescens* Swainson. *J Exp Mar Biol Ecol* 155:263–277

*Editorial responsibility: Otto Kinne (Editor-in-Chief), Oldendorf/Luhe, Germany*

*Submitted: July 6, 2004; Accepted: October 29, 2004  
Proofs received from author(s): February 1, 2005*