

**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

# Labile Fe(II) concentrations in the Atlantic sector of the Southern Ocean along a transect from the subtropical domain to the Weddell Sea Gyre

G. Sarthou<sup>1,2</sup>, E. Bucciarelli<sup>1,2</sup>, F. Chever<sup>1,2,\*</sup>, S. P. Hansard<sup>3</sup>,  
M. Gonzalez-Davila<sup>4</sup>, J. M. Santana-Casiano<sup>4</sup>, F. Planchon<sup>1,2</sup>, and S. Speich<sup>1,5</sup>

<sup>1</sup>Université Européenne de Bretagne, France

<sup>2</sup>Université de Brest, CNRS, IRD, UMR 6539 LEMAR, IUEM; Technopôle Brest Iroise, Place Nicolas Copernic, 29280 Plouzané, France

<sup>3</sup>Florida Geological Survey, 903 W. Tennessee Street, Tallahassee FL 32301, USA

<sup>4</sup>Facultad de Ciencias del Mar. Departamento de Química, Universidad de Las Palmas de Gran Canaria, Campus de Tafira, 35017, Las Palmas de Gran Canaria, Spain

<sup>5</sup>Laboratoire de Physique des Océans, UMR 6523, IFREMER, CNRS, IRD, UBO, 29280 Plouzané, France

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



\*current address: Université de Brest, CNRS, IRD, UBO, UMR 6538 LDO, IUEM ; Technopôle Brest Iroise, Place Nicolas Copernic, 29280 Plouzané, France

Received: 1 April 2011 – Accepted: 8 April 2011 – Published: 29 April 2011

Correspondence to: G. Sarthou (geraldine.sarthou@univ-brest.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**BGD**

8, 4163–4208, 2011

---

**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Labile Fe(II) distributions were investigated in the Sub-Tropical South Atlantic and the Southern Ocean during the BONUS-GoodHope cruise from 34 to 57° S (February–March 2008). Concentrations ranged from below the detection limit (0.009 nM) to values as high as 0.125 nM. In the surface mixed layer, labile Fe(II) concentrations were always higher than the detection limit, with values higher than 0.060 nM south of 47° S, representing between 39% and 63% of dissolved Fe (DFe). Biological production was evidenced. At intermediate depth, local maxima were observed, with the highest values in the Sub-Tropical domain at around 200 m, and represented more than 70% of DFe. Remineralization processes were likely responsible for those sub-surface maxima. Below 1500 m, concentrations were close to or below the detection limit, except at two stations (at the vicinity of the Agulhas ridge and in the north of the Weddell Sea Gyre) where values remained as high as ~0.030–0.050 nM. Hydrothermal or sediment inputs may provide Fe(II) to these deep waters. Fe(II) half life times ( $t_{1/2}$ ) at 4 °C were measured in the upper and deep waters and ranged from 2.9 to 11.3 min, and from 10.0 to 72.3 min, respectively. Measured values compared quite well in the upper waters with theoretical values from two published models, but not in the deep waters. This may be due to the lack of knowledge for some parameters in the models and/or to organic complexation of Fe(II) that impact its oxidation rates. This study helped to considerably increase the Fe(II) data set in the Ocean and to better understand the Fe redox cycle.

## 1 Introduction

Iron (Fe) is an essential micronutrient for all marine organisms, playing a key role in many metabolic processes, such as photosynthesis, respiration, nitrate reduction, and nitrogen fixation (Sunda, 1988, 1989). Its low concentrations have been shown to limit primary production in more than 50% of the ocean (Boyd and Ellwood, 2010). All

**BGD**

8, 4163–4208, 2011

### Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



natural and artificial Fe fertilization experiments unequivocally showed the importance of Fe for the carbon cycle, particularly for the growth and composition of the phytoplanktonic community (Boyd et al., 2000; Coale et al., 1996; Gervais et al., 2002; Tsuda et al., 2003; Boyd, 2004; Coale et al., 2004; Blain et al., 2007; Pollard et al., 2007). Despite numerous studies on Fe cycling the last 25 years, lots of unknowns persist, in particular because Fe chemistry in seawater is very complex. Fe has been observed to occur in two redox states (Fe(III) and Fe(II), Waite and Morel, 1984). In oxic seawater, the thermodynamically most stable state is Fe(III), but is highly insoluble (0.011 nM in 0.7 NaCl solution, Liu and Millero, 2002) and is rapidly hydrolyzed resulting in the formation of various Fe(III) oxyhydroxide (de Baar and de Jong, 2001). These species, with Fe(OH)<sub>3</sub> being the dominant one in seawater at pH ~8, have the tendency to form colloidal Fe (Kuma et al., 1996) which coagulate and form particulate Fe (Johnson et al., 1997). In contrast to Fe(III), Fe(II) is more soluble but is rapidly oxidized by oxygen (O<sub>2</sub>) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) (Millero et al., 1987; Millero and Sotolongo, 1989; Santana-Casiano et al., 2004, 2005; Gonzalez-Davila et al., 2005, 2006). Although Fe(II) in seawater is less stable than Fe(III), recent models of Fe acquisition by eukaryotic phytoplankton suggest that the reduction of Fe(III) to Fe(II), with subsequent re-oxidation to Fe(III), is a possible mechanism by which Fe is made more bioavailable to phytoplankton (Shaked et al., 2004; Salmon et al., 2006; Maldonado et al., 2006; Morel et al., 2008). Numerous studies have investigated the oxidation of Fe(II) by O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> in different aqueous solutions to understand the behavior of Fe(II) in natural waters (Santana-Casiano et al., 2006 and references herein). The most widely accepted mechanism to describe Fe oxidation with O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> is the Haber-Weiss mechanism, with reactions 1 or 3 limiting the overall oxidation rate (King et al., 1995).





The rates for Eq. (1–4) strongly depend on the relative concentrations of the individual Fe(II) species in solution, mainly  $\text{Fe}^{2+}$ ,  $\text{Fe(OH)}^+$ ,  $\text{Fe(OH)}_2$ ,  $\text{FeHCO}_3^+$ ,  $\text{Fe(CO}_3)_2^{2-}$ , and  $\text{FeCO}_3(\text{OH}^-)$  (Millero, 1989; King, 1998; Santana-Casiano et al., 2006; Trapp and Millero, 2007), as well as on the concentrations of  $\text{O}_2$  and  $\text{H}_2\text{O}_2$ , pH, temperature ( $T$ ) and salinity ( $S$ ). In warm oxygenated seawater, the half-life of Fe(II) can be as low as few seconds (King, 1998), whereas in cold surface or suboxic waters it can be on the order of hours to days (Croot et al., 2001, 2008; Hansard et al., 2009; Moffett et al., 2007).

Several mechanisms provide Fe(II) in the dissolved phase and are reviewed by Hansard et al., 2009). They consist of in-situ processes (both abiotic and biotic) and external sources. The abiotic in-situ processes are mainly photochemical reactions. They include photoreduction of dissolved Fe(III) (oxy)hydroxides, and photoreduction or photolysis of organic, colloidal and particulate Fe (Rich and Morel, 1990; Wells et al., 1991; Kuma et al., 1992a, b; King et al., 1993; Barbeau et al., 2001; Rijkenberg et al., 2006). Extracellular reduction of Fe(III) by photochemically-produced superoxide or reductive dissolution of particulate Fe(III) may also occur (Voelker and Sedlak, 1995; Rose and Waite, 2002, 2003; Kustka et al., 2005; Rose et al., 2005; Salmon et al., 2006). Biotic in-situ processes include bioreduction of organic Fe(III) at cell surface (Maldonado and Price, 2001, 1999; Shaked et al., 2004; Morel et al., 2008), reduction by biogenic superoxide (Rose and Waite, 2002, 2003; Salmon et al., 2006), as well as remineralization via microbial activity (Alldredge and Cohen, 1987), cell lysis (Gobler et al., 2002), and grazing (Hutchins and Bruland, 1994; Hutchins et al., 1995; Sarthou et al., 2008). The external sources are atmospheric inputs (Kieber et al., 2001; Journet et al., 2007; Ozsoy and Saydam, 2001), sediment inputs (Elrod et al., 2004; Lohan and Bruland, 2008), submarine groundwater discharge (Windom et al., 2006), and hydrothermal vents (Coale et al., 1991; Chin et al., 1994; Field and Sherrell, 2000; Statham et al., 2005; Bennett et al., 2008). These sources supply Fe(II), which can then be transported by advective and/or diffusive mixing.

---

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Although it is now evident that Fe(II) plays a key role in Fe chemistry and bioavailability in the ocean, there are relatively few open-ocean measurements of Fe(II) available due to the difficulty to measure such an ephemeral species at subnanomolar concentrations (Bruland and Rue, 2001). To our knowledge, the most comprehensive data set of Fe(II) was published by Hansard et al. (2009) in the Pacific Ocean along a zonal transect at 30° N and a meridional one at 152° W, within the CLIVAR/CO<sub>2</sub> Repeat Hydrography Program. In this paper, we present results of the Bonus-GoodHope (BGH) cruise, carried out in February-March 2008 during the International Polar Year in the Sub-Tropical South Atlantic and the Southern Ocean. Fe(II) distributions as well as oxidation rates are presented and results are discussed considering different processes such as photoreduction, oxidation, biological production, and different inputs, such as atmospheric, sediment, and hydrothermal inputs, and/or advection and mixing.

## 2 Materials and methods

### 2.1 Study area

Sampling and ship board measurements were done aboard *R/V Marion Dufresne* from 8 February to 24 March 2008 in the Atlantic sector of the Southern Ocean during the BGH cruise. Figure 1 shows the cruise track together with the main oceanographic fronts and domains crossed during the cruise, from north to south: (i) the subtropical domain and the southern subtropical front (S-STF), (ii) the Antarctic Circumpolar Current (ACC) domain with 3 fronts crossed, the subantarctic front (SAF), the polar front (PF) and the southern ACC front (SACCF), and (iii) the eastern part of the Weddell Sea gyre with the southern boundary (SBdy) separating this domain from the ACC. Twelve stations were sampled for Fe(II), among which seven were sampled between 0 and 2000 m (Large stations L1 to L7) and five between 0 to 4000 m (Super stations S1 to S5). The position of each station is reported in Fig. 1 and Table 1.

**BGD**

8, 4163–4208, 2011

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2 Sample processing and analytical methods

Samples were collected using acid-cleaned 12 L Go-Flo bottles. When not in use, the Go-Flo bottles were stored inside a clean van with plastic bags covering the top and the bottom including the spigots. On station, the Go-Flo bottles were transferred to the sampling deck and mounted on a Kevlar cable. Plastic bags were removed just after attachment to the Kevlar cable. When the expected depths were reached, bottles were tripped by a Teflon® messenger. Once back on board, the bottles were directly transferred to the clean van for sub-sampling. All sub-samples for Fe(II) measurements were immediately collected in previously 60 mL acid-cleaned high density brown polyethylene (HDPE) bottles. The maximum time between sub-sampling time from the Go-Flo bottle and analysis was 3 min. In order to minimize this time, no filtration was carried out, thus avoiding an underestimation of Fe(II) concentrations due to rapid oxidation. However, Fe(II) produced by fast-kinetic processes involving lithogenic or biogenic particles, such as reductive dissolution of particulate Fe(III) (Rich and Morel, 1990) or bioreduction of organic Fe(III) at cell surface (Maldonado and Price, 1999, 2001; Shaked et al., 2004; Morel et al., 2008) is measured and tends to overestimate Fe(II) concentrations. On the other hand, filtration step can also bring some artifacts. A first one is the damage and explosion of the cells that may release Fe(II) in the dissolved phase (Hutchins et al., 1993). A second one is related to the stress of the cells during filtration that may increase the production of superoxide (Godrant et al., 2009). This reactive oxygen species is involved in the redox Fe cycle, but it also initiates the three-step oxidation of luminol (Ussher et al., 2005; Rose and Waite, 2001), thus potentially inducing an overestimation of the Fe(II) concentrations in the dissolved phase. In the following, the term “labile” Fe(II) is then used, since the measurement is operationally defined and the exact speciation of the measured fraction is not known (Ussher et al., 2007).

Labile Fe(II) concentrations were determined by chemiluminescence flow injection analysis following the method of Croot and Laan (2002). As in Croot and Laan (2002),

**BGD**

8, 4163–4208, 2011

### Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



there was no preconcentration prior to reaction with luminol, allowing a minimal analytical time (~80–90 s). The percentage of labile Fe(II) was calculated as the ratio of labile Fe(II) over the dissolved Fe concentration (i.e.  $(\text{Fe(II)}/\text{DFe}) \cdot 100$ ). The instrument was calibrated by standard addition using peak height measurements and freshly prepared acidified Fe(II) standards added to a surface (20–300 m) and a deep (500–2000 m) sample. Samples were stored at 4 °C in the dark for 24 h to enable complete decay of ambient Fe(II). Non-linear calibration curves were observed, due to the kinetics of luminol oxidation and free-radical generation (Rose and Waite, 2001), and a polynomial 2nd degree curve-fitting technique was used to quantify the results. The blank was determined by running an aged seawater sample (4 °C for 24 h). The detection limit was calculated as three times the standard deviation of the blank value and was equal to  $0.009 \pm 0.006$  nM ( $n = 29$ ).

After each Fe(II) spike, the change in Fe(II) signal was recorded over 10 min every 80–90 s, and allowed us to estimate Fe(II) oxidation rates, similarly to Roy et al (2008). The oxidation rates may be overestimated by up to ~10% due to pH decrease after standard addition (~0.1 pH unit/standard addition). In addition to the samples used for the calibrations, for the four super stations S2 to S5, analyses were also performed with deeper samples (2300–3600 m).

### 2.3 Ancillary measurements

Samples for dissolved Fe (Fe(III) + Fe(II), DFe) and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) analyses were collected from the same Go-Flo bottles as for labile Fe(II) measurements in acid-washed low density polyethylene (LDPE) and high density brown polyethylene (HDPE) bottles, respectively. DFe concentrations were determined by FIA with on line preconcentration onto 8-HQ resin and chemiluminescence detection (Obata et al., 1993; modified by Sarthou et al., 2003). The comprehensive data set is published elsewhere (Chever et al., 2010).  $\text{H}_2\text{O}_2$  samples were analyzed on board within 3 h after collection using a flow injection method with chemiluminescent detection (Yuan and Shiller, 1999). A comprehensive data set will be available elsewhere (Bucciarelli

**BGD**

8, 4163–4208, 2011

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





et al., 2011). The other ancillary parameters were measured from the closest (15–50 min) CTD cast. In-situ  $T$  and  $S$  were acquired from a CTD SEABIRD SBE 911+ mounted in a Niskin-rosette. Oxygen ( $O_2$ ) concentrations were measured on board by Winkler titration. The pH was measured in total scale at a constant temperature of 25 °C ( $pH_{T,25}$ ) using an automated spectrophotometric technique with m-cresol purple as indicator (Gonzalez-Dávila et al., 2003). A VINDTA 3C system (Mintrop et al., 2000), with coulometer determination was used for the titration of the total dissolved inorganic carbon ( $C_T$ ) after phosphoric acid addition. Carbonate concentration were estimated from  $pH_{T,25}$ , total alkalinity (potentiometrically titrated, Mintrop et al., 2000) and  $C_T$ , and computed by using CO2sys.xls v12 (Lewis and Wallace, 1998).

### 3 Results

#### 3.1 Hydrography

The hydrography of the area is detailed in Chever et al. (2010), based on Gladyshev et al. (2008) and using the  $S$  and  $T$  data measured during the BGH cruise (Fig. 2). The subtropical domain (STZ) extended southward to the S-STF (about 42° S, between station L2 and S2). Although station S2 is located south of the S-STF, its surface waters exhibit  $S$  and  $T$  signatures of subtropical waters. This station will be considered in the following as a Sub-Tropical station. Further south, the domain of the ACC extended to the Southern Boundary (SBdy) (~42° S to ~55° S, stations S2 to L7). The SAF, PF and SACCF were found at ~44° S, 50° S, and 51° S, respectively. South of the Sbdy (station S5), waters were entrained in the large scale cyclonic flow of the Weddell gyre.

Along the transect, several major water masses were sampled. They are described elsewhere (Arhan et al., 2011; Speich et al., 2011) and briefly summarized here and on Fig. 2. In the subtropical domain, the central water layer was mostly occupied by waters of Indian Ocean origin (Boebel et al., 2003). Below, the Antarctic Intermediate Water (AAIW), the Upper Circumpolar Deep Water (UCDW), the diluted North Atlantic Deep

**BGD**

8, 4163–4208, 2011

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Water (NADW), and finally the Antarctic Bottom Water (AABW) were observed. In the ACC, between the SAF and PF, below the surface mixed layer (SML) were located the AAIW, the Winter Waters (AAWW, marked by a  $T$  minimum), the UCDW, and the Lower Circumpolar Deep Water (LCDW) with, north of the PF, an addition of diluted South West NADW (SW-NADW, Whitworth III and Nowlin, 1987). Deeper, the AABW was observed against the northern flank of the Mid Atlantic Ridge (MAR). Finally, south of the SBdy, the near surface waters were thought to have been in contact with the western continental margin of the Antarctic Peninsula, while the deeper waters might have had a more recent contact with the northern topographic limit of the Weddell Basin (Orsi et al., 1993; Meredith et al., 2000; Klatt et al., 2005).

### 3.2 Labile Fe(II) concentrations

Labile Fe(II) concentrations are reported in Table 2 and plotted on Fig. 3a. Within the whole data set, concentrations ranged from values below the detection limit to values as high as 0.125 nM.

In the SML, labile Fe(II) concentrations were systematically higher than the detection limit. Over the whole transect, the mean value was equal to  $0.039 \pm 0.024$  nM ( $n = 26$ , median value = 0.037) and concentrations ranged from 0.012 nM to 0.116 nM. Both the minimum and maximum values were observed in the ACC (Station L7, 60 m, and station S4, 30 m, respectively). The highest percentages of Fe(II) relative to DFe (40–67%, Fig. 3b) were found at three stations south of 47° S (S3, S4, and S5) and were associated with values of Fe(II) higher than 0.060 nM. At the other stations, percentages were lower, ranging from 3% to 25%.

At intermediate depth (between the SML and 1500 m), values ranged from below the detection limit to 0.125 nM (Station S2, 196 m). The maxima observed around 200–300 m, more pronounced in the sub-tropical zone, corresponded to percentages of labile Fe(II) relative to DFe ranging from 30% to 70%. Elsewhere values varied between less than 1% and 26%.

**BGD**

8, 4163–4208, 2011

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Below 1500 m, except at station S2 where values were around 0.030 nM at ~2000 m and at station S5 where values ranged from 0.035 to 0.050 nM below 2100 m, labile Fe(II) concentrations were close to or below the detection limit (mean value  $0.010 \pm 0.002$  nM,  $n = 11$ ), representing less than 4% of DFe.

### 3.3 Fe(II) oxidation rates

The natural logarithm transformation of Fe(II) chemiluminescence over time showed linear decreases in signal for all surface and deep samples and spike additions, indicating a pseudo first-order kinetics for Fe(II) oxidation during the timescale monitored (Fig. 4). The pseudo-first order rate constants  $k_{ox}$  at 4 °C were experimentally determined as the slope of the ln-transformed chemiluminescence signal vs. time. The half-lives ( $t_{1/2}$ ) were then calculated as  $t_{1/2} = \ln(2)/k_{ox}$ , for the four Fe(II) spikes (Fig. 5, Table 3).

In the upper waters (20–300 m), values ranged from 2.9 to 11.3 min and increased significantly with latitude (linear regression,  $r^2 = 0.65$ , slope =  $0.3 \text{ min}/^\circ \text{S}$ ,  $P < 0.01$ ,  $n = 12$ ). The mean value of all data was equal to  $6.7 \pm 2.6$  min (median value 6.7 min). In the deep waters (500–2000 m),  $t_{1/2}$  varied between 10.0 and 72.3 min. No significant relationship was observed with latitude (linear regression,  $P = 0.57$ ). The mean value of all the data at depth was equal to  $37.0 \pm 19.7$  min (median value 35.2 min). On average,  $t_{1/2}$  was 6 times higher in the deep than in the upper waters. At the four super stations where two different depths were sampled below 500 m, values were not significantly different (paired t-test,  $P = 0.6$ ,  $n = 4$ ).

**BGD**

8, 4163–4208, 2011

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4 Discussion

### 4.1 Comparison with previously reported Fe(II) data

There are relatively few Fe(II) data reported in the literature (Table 4). Previous studies have been either generally geographically restricted, except the one carried out by Hansard et al. (2009), or focused on particular areas, such as the suboxic zones, or performed during artificial Fe experiments in the Southern Ocean or the Subarctic Pacific.

For the open ocean, measurements were done in the Pacific Ocean (Hansard et al., 2009; O'Sullivan et al., 1991), the Atlantic Ocean (Bowie et al., 2002; Boye et al., 2003, 2006), and the Southern Ocean (Croot et al., 2007). Typically, Fe(II) concentrations range from the detection limit to  $\sim 0.050$ – $0.080$  nM, with concentrations that tend to increase in the surface waters due to photoproduction processes or in water masses influenced by continental margins. Indeed, studies carried out in continental or shelf waters showed concentrations up to 0.3–0.5 nM (Boye et al., 2006; Hansard et al., 2009), and even 3 nM (Waite et al., 1995). These high concentrations were likely due to transport of sediment-derived Fe and/or diffusion of Fe(II) directly from pore waters. These processes were even more pronounced in Fe-rich coastal environments, where values of Fe(II) could reach 40 nM (Hong and Kester, 1986). In suboxic zones, low oxygen concentrations slow down the oxidation rates of Fe(II) and maxima of Fe(II) (0.2–0.6 nM) were associated with the oxygen minima and/or nitrite maxima (Hopkison and Barbeau, 2007; Moffett et al., 2007). During the artificial Fe fertilization experiments, concentrations of Fe(II) were shown to remain elevated ( $>0.2$  nM and up to 1 nM) for several days after enrichment (Croot et al., 2001, 2005, 2008; Croot and Laan, 2002), due to a potential combination of slow oxidation rate and organic complexation (Roy et al., 2008).

Our values are within the range of variation of values in the open ocean and/or suboxic regions, although most of the previous studies measured Fe(II) in the dissolved phase. As already mention in Sect. 2.2, the non filtration may induce underestimation

### Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



or overestimation of the Fe(II) signal, but the good consistency between our data set and previous one suggests that it may be a non-issue. Another difference in methodology is the acidification of the samples at pH 6 prior to analysis (Hansard et al., 2009) to reduce the rate of Fe(II) oxidation. Although the acidification step may result in a measurement of “readily reducible Fe(III)” rather than actual Fe(II), the similarity between the two data set suggests that the method of Hansard et al. (2009) adequately corrected for apparent Fe(III) reduction caused by acidification.

## 4.2 Spatial and vertical distribution of labile Fe(II) during the BGD cruise

On the contrary to DFe concentrations (Chever et al., 2010), no systematic decrease in labile Fe(II) concentrations was observed from the north to the south of the section. The high values of DFe in the STZ were suggested to be due to direct dust deposition coming from Patagonia and/or to lateral advection of Indian Ocean water masses enriched by dust inputs. Another potential source of DFe was the African continental margin both in the SML and deeper waters (Chever et al., 2010). The mean advective time of these waters estimated from the AVISO Mean Absolute Dynamic Topography (Ducet et al., 2000) for both the specific cruise period (2007–2008) and for the entire satellite time series (i.e., from 1992 to 2008) was equal to 1–3 months. Given the very short half-life time of Fe(II) in surface and deep waters (3–72 min), this delay is likely to be too long to preserve the signal of a potential enrichment of water masses by the African continental margin, unless stabilization of Fe(II) by organic complexation occurs (Roy et al., 2008). In the following, we will examine more in detail the depth profiles of labile Fe(II) and discuss the potential sources of Fe(II) in our studied area.

### 4.2.1 Surface mixed layer (SML)

Variations in Fe redox speciation along the day in the SML have been observed in numerous studies (e.g. Hong and Kester, 1986; Johnson et al., 1994; O’Sullivan et al., 1991; Waite et al., 1995; Bowie et al., 2002; Roy et al., 2008). In our study, 5 stations were sampled at night (S1, L2, L4, L6 and L7) and 7 during the day (L1, S2, L3,

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



S3, L5, S4 and S5). The daytime and nighttime mean values were not significantly different ( $0.044 \pm 0.029$  nM and  $0.033 \pm 0.012$  nM, respectively, student test,  $P = 0.39$ ). Unexpectedly, daytime values were negatively correlated to the SML-integrated solar radiation (ANOVA,  $F = 9.12$ ,  $P < 0.01$ ). A parameter that appeared to control the labile Fe(II) values in our data set was the time at which samples were taken. Indeed, maximum values of labile Fe(II) as well as %Fe(II)/DFe were observed for the three stations (S3, S4 and S5) sampled between 12:00 and 16:00 (Universal Time, Fig. 6). This is consistent with photochemical reactions producing Fe(II) in the SML (Rich and Morel, 1990; Wells et al., 1991; Kuma et al., 1992a, b; King et al., 1993; Barbeau et al., 2001; Rijkenberg et al., 2006; Voelker and Sedlak, 1995; Rose and Waite, 2002, 2003; Kustka et al., 2005; Rose et al., 2005; Roy et al., 2008). However, at these stations, the time between the tripping of the Go-Flo bottles (removing the sample from the influence of light) and analysis (15–20 min) were higher than the measured half-life times of Fe(II) (9–11 min). When estimating the initial in situ values that Fe(II) would have if photochemical reactions were the only sources of Fe(II), values are ~2-fold the DFe concentrations. This would suggest that photochemical reactions could not be the only sources of Fe(II) and that a biological production occurred. This would be coherent with a higher photosynthetic efficiency around noon (Legendre et al., 1988). Moreover, the biological superoxide production rates have been recently measured in oceanic and coastal waters of the Gulf of Alaska and values were up to  $0.3$  nM min<sup>-1</sup> (Hansard et al., 2010). Considering our labile Fe(II) data and measured  $t_{1/2}$ , and assuming steady-state, the calculated Fe(II) production rate was about  $0.002$ – $0.013$  nM min<sup>-1</sup>. The potential production rates of superoxide were thus likely sufficient to drive Fe(II) production. If a biological production exists in the surface waters, that would explain our values higher than the detection limit even at stations sampled at night. This night time observation of Fe(II) at pM levels was already done during the FeCycle experiment in the HNLC waters of the South West Pacific (Croot et al., 2007). These authors estimated that ~26% of the inorganic Fe(II) biologically produced would be organically complexed and the rest would be oxidized.

---

**Labile Fe(II)  
concentrations in the  
South Atlantic**G. Sarthou et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



During the ANT XXIV/3 cruise, maxima of DFe and dissolved manganese (Mn) were observed in the surface waters in the Bouvet region at 54–55° S (Klunder et al., 2010; Middag et al., 2010), and were ascribed to recent deposition of aeolian dust originating from South America. At our L7 station (~55° S), however, no surface maximum of labile Fe(II) was observed, neither clear surface maxima of DFe in the Bouvet region (Chever et al., 2010). This may be due to the fact that our sampling was performed three weeks later than theirs and/or to our lower sampling resolution, missing some of the key features they observed in the Bouvet region. At station S4 (~52° S), where we observed the highest value of labile Fe(II) in the SML, a combination of atmospheric inputs, biological production and photochemical processes (see above) might explain this high value.

#### 4.2.2 Intermediate depths (between the SML and 1500 m)

In the STZ (stations L1, S1, L2, and S2), large sub-surface maxima of labile Fe(II) centered at 200–300 m were observed. Concentrations started to increase just at or below the maximum of chlorophyll-a (Ras and Clautres, pers. com.) and high values were observed as deep as 700 m (0.09 nM, station S1). Although generally lower than in the STZ (0.023–0.087 nM), sub-surface maxima were also observed in the ACC and north of the Weddell Gyre at depth between 60 and 300 m. As discussed above, due to the relatively long transport time between the African continental margin and the stations in the STZ, an input of Fe(II) from water masses enriched by the African continental margin is unlikely, unless organic complexation stabilized Fe(II) (Roy et al., 2008). Observed trends in labile Fe(II) compare well with depth profiles of dissolved Mn in the same sector of the Southern Ocean during the ANT XXIV/3 expedition (Middag et al., 2010). Sub-surface maxima of Mn were identified at depth around 150 m and coincided with a gradient in the potential density anomaly ( $\sigma_\theta$ ). Similar relationship was evidenced for  $^{234}\text{Th}/^{238}\text{U}$  ratio in the subsurface, with a steep increase towards values near or over 1 at the pycnocline (Rutgers van der Loeff et al., 2011). This trend in  $^{234}\text{Th}/^{238}\text{U}$  ratio indicates no more scavenging of biogenic particles at the base of

**BGD**

8, 4163–4208, 2011

### Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





the mixed layer and may rely on small scale remineralisation of sinking material (Cai et al., 2008, Maiti et al., 2010). During the BGH cruise, high resolution depth profiles were also obtained for  $^{234}\text{Th}/^{238}\text{U}$  ratio and a more detailed data set is available from the surface to 1000 m depth (Planchon et al., 2011). Except at station L6, our labile Fe(II) maxima also coincided with increases in  $^{234}\text{Th}/^{238}\text{U}$  ratios in the subsurface (Fig. 7).  $^{234}\text{Th}/^{238}\text{U}$  ratios ranging from 1 to maximum 1.3 clearly indicated fast and intense remineralization of sinking material in the mesopelagic zone (100–600 m), which was further confirmed by parallel biogenic particulate barium data (Fig. 7). Therefore one of the most likely sources of labile Fe(II) in the subsurface might be remineralization/disaggregation of biogenic particles settling from above. Moreover, the very large sub-surface maxima in the STZ were consistent with a bloom in a senescent stage (Ras and Claustre, pers. com.). The value of %Fe(II)/DFe at these depths can be as high as 50–70%, suggesting that biogenic Fe is mainly regenerated as Fe(II) species, as already observed in other studies (Hutchins and Bruland, 1994; Sarthou et al., 2008). The remineralization of organic matter (OM) is associated with the consumption of oxygen and the apparent oxygen utilization (AOU) can also provide a quantitative estimate of the amount of material that has been remineralized (Sarma et al., 2007). In our study, no coincidence between the sub-surface maxima of labile Fe(II) and AOU was observed (not shown). Indeed, the maxima of AOU were around 1000–1200 m in the STZ and in the ACC north of the PF, and around 300–800 m south of the PF. The very particle reactive  $^{234}\text{Th}$  has a short half-life (24.1 days), and may then help to trace remineralization processes with a much shorter half-life than when using the AOU, and which occur locally.

In the southern part of our transect (stations L6, S4, L7, S5, Fig. 8), the winter waters were strongly visible on the T data and were associated with local minima of labile Fe(II) concentrations. At the end of the winter, the stratification of the water column divides the Antarctic surface waters into two water masses that evolve differently: the Antarctic surface summer waters (AASSW) and the Antarctic winter waters (AAWW). The cold AAWW are less affected by phytoplankton activity and keep the signal of the winter

**BGD**

8, 4163–4208, 2011

---

**Labile Fe(II)  
concentrations in the  
South Atlantic**G. Sarthou et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





time. Thus, when the concentrations of a given element in the AAWW is higher than the ones in the AASSW, the variation of concentrations can be assumed to represent the biological uptake (Sarhou et al., 1997; Blain et al., 2007). In our case, labile Fe(II) concentrations are lower in the AAWW. This could reflect processes involving a biological source of Fe(II) in the AASSW and confirm results observed in the SML.

### 4.2.3 Below 1500 m

Only two stations showed labile Fe(II) concentrations higher than the detection limit below 1500 m. Station S2 was located at the vicinity of the Agulhas Ridge and local maxima of DFe were observed there (Chever et al., 2010), suggesting hydrothermal or sediment inputs (Elrod et al., 2004; Boyle and Jenkins, 2008; Bennett et al., 2008; Tagliabue et al., 2010). %Fe(II)/DFe was not very high (1.4–5%) and Fe(II) half-life was the lowest of the section (Fig. 3b), suggesting that Fe(II) is continuously provided to the deep waters but reoxidized quite fast. During the ANT XXIV/3 expedition, Klunder et al. (2010) and Middag et al. (2010) evidenced hydrothermal inputs of Fe and Mn in the Bouvet region (52–56° S). The hydrothermal signal was not clearly seen on our DFe (Chever et al., 2010) nor on our labile Fe(II) data, likely due to a lower resolution of our sampling. At station S5, concentrations of labile Fe(II) as high as 0.050 nM were observed at 3500 m, with %Fe(II)/DFe equal to 13%. Along the zero meridian, DFe concentrations in the deep waters north of the Weddell Gyre ( $0.47 \pm 0.16$  nM,  $n = 98$ , Klunder et al., 2010, and  $0.42 \pm 0.07$ ,  $n = 4$ , Chever et al., 2010) were higher than south of the Weddell Gyre ( $0.33 \pm 0.14$  nM,  $n = 98$ , Klunder et al., 2010). North of the Weddell Sea Gyre, the deep waters flow eastward, and might have had a recent contact with the northern limit of the Weddell Basin (Orsi et al., 1993; Meredith et al., 2000; Klatt et al., 2005), flowing along the North Weddell Ridge. A local reductive dissolution of particles coming from the slope sediments of the ridge may explain the high values of labile Fe(II) at this station.

## Labile Fe(II) concentrations in the South Atlantic

G. Sarhou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 4.3 Oxidation rates

The Fe(II) half-life values in the surface waters were similar to previous values estimated in natural surface seawater at near-ambient concentrations (6–28 min, Croot et al., 2008). Our values were higher in the deep than in the surface waters. To our knowledge, our study is the first one to measure Fe(II) oxidation rates in natural deep seawater at near-ambient concentrations. The deep values were never as high as the ones estimated by Hansard et al. (2009, up to 690 min) in the Pacific Ocean. However, their  $[O_2]$  at  $\sim 1000$  m were as low as  $13 \mu\text{M}$ , whereas  $[O_2]$  was never lower than  $100 \mu\text{M}$  along the BGH transect. The overall oxidation rate of Fe(II) is a function of oxidant concentrations (e.g. oxygen, hydrogen peroxide, superoxide, etc., Gonzalez-Davila et al., 2006),  $T$ , pH, as well as Fe(II) chemical speciation (mainly  $\text{Fe}^{2+}$ ,  $\text{Fe}(\text{OH})^+$ ,  $\text{Fe}(\text{OH})_2$ ,  $\text{FeCO}_3$ ,  $\text{Fe}(\text{CO}_3)_2^{2-}$ ,  $\text{Fe}(\text{CO}_3)(\text{OH})^-$ , Millero, 1989; King, 1998; Santana-Casiano et al., 2006; Trapp and Millero, 2007). To compare our data with theoretical ones, we used two published models of Fe(II) oxidation kinetics and in situ physical-chemical conditions (Model I: Santana-Casiano et al., 2005; Model II: Trapp and Millero, 2007; see supplementary material for detailed calculations). Theoretical values are given in Table 3. The two models differ in the equations used for the calculations of the oxidation rate constants of the individual species for oxidation by oxygen (see Supplement). Moreover, Model I considers the  $\text{Fe}(\text{CO}_3)(\text{OH})^-$  species, as well as the oxidation by the superoxide. None of the two models considers organic matter effects, and differences in oxidation rates among samples are only related to  $T$ , pH,  $S$ , and carbonate effects.

In the upper waters, for Model I, the Fe(II) half-life times ranged from 7.5 to 18.4 min, with a mean value of  $12.4 \pm 3.0$  min. For Model II, values ranged from 5.5 to 13.6 min, with a mean value of  $9.5 \pm 2.3$  min. The ranges of variations of the two theoretical data sets were similar, although a paired-t test showed that the two data sets were significantly different ( $P < 0.01$ ,  $n = 12$ ), with values from Model I always higher than values from Model II (up to 5 min). The measured  $t_{1/2}$  (2.9–11.3 min, mean value  $6.7 \pm 2.6$  min) showed systematically lower values than the theoretical ones of both models (by up to 10–15 min at station L2), except at station S5 where the measured value was slightly

---

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 higher than the Model II one. A much larger difference was observed in the deep waters between the two models. The Model I values ranged from 86 to 138 min (mean value  $112 \pm 19$  min), whereas the Model II ones were about 4 times lower, ranging from 19 to 32 min (mean value  $26 \pm 5$  min). This difference may come from the uncertainties in the model parameterization and/or on superoxide concentrations which could vary at depth more than oxygen concentration (see Supplement). However, what both models indicated was that in the deep waters, the half life times were almost constant (less than a factor of two) compared to the measured values which varied by a factor of 7 (10–72 min). Indeed, in the deep waters, the pH and  $T$  ranges of variation are relatively small, inducing a small range of variation of theoretical values. The largest range of variation of the measured values could be induced by a change in oxygen concentrations. Indeed, oxygen may have been consumed by micro-organisms in the sample during the 24 h storage or increase if an oxygen contamination occurred. Also, when the oxidation rates were measured at 4 °C, the deep samples and some of the surface samples were heated while most of the surface samples were cooled. These differences can affect the intermediate species and equilibrium processes. Moreover, after heating, changes in pH due to CO<sub>2</sub> dissolution-exchange can modify the Fe(II) speciation.

10  
15  
20  
25 Another explanation for the variability of the measured  $t_{1/2}$  and the discrepancy between measured and theoretical is organic complexation. Although dissolved Fe(III) is now well known to be strongly bound by organic chelators in seawater (Gledhill and van den Berg, 1994; Rue and Bruland, 1997; Gerringa et al., 2006, 2008; Thuróczy et al., 2010), organic complexation of Fe(II) has been suggested but never directly measured (Croot et al., 2007, 2008; Roy et al., 2008). In the Subarctic Pacific, Roy et al. (2008) observed a significant difference between the measured Fe(II) oxidation rates in natural surface water and the ones in UV-treated surface water, which strongly suggested that organic ligands influenced Fe(II) speciation in seawater. Like for Fe(III) species, the Fe(II) organic speciation may help maintaining Fe(II) in the dissolved phase. However, numerous studies on the effect of Fe organic complexation on the oxidation kinetics

---

**Labile Fe(II)  
concentrations in the  
South Atlantic**G. Sarthou et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



showed that the organic complexation can either increase or decrease the Fe(II) oxidation rates (Rijkenberg et al., 2006 and references herein). Variability of organic compounds within the water column could thus induce variability in the observed oxidation rates.

## 5 Conclusions

Concentrations of labile Fe(II) in the surface waters were systematically higher than the detection limit of our analytical method. The highest values were observed where sampling was done between 12:00 and 16:00, suggesting a biological production of Fe(II) in the SML linked to photosynthesis. South of the section, local minima coinciding with the Winter Waters confirm that direct biological reduction of Fe(III) may occur in the SML. This would explain why our nighttime surface samples have concentrations higher than the detection limit. At intermediate depths, sub-surface maxima were observed all along the section, although more pronounced in the STZ. A bloom at a senescent stage in the STZ, together with a good consistency between the maxima of labile Fe(II) and the increase in  $^{234}\text{Th}/^{238}\text{U}$  towards values over 1 suggested that Fe remineralization occurred at those depths and that Fe was mainly regenerated as Fe(II) species. In the deep waters, labile Fe(II) concentrations were higher than the detection limit at two stations: one located at the vicinity of the Agulhas ridge and another one in the north of the Weddell Gyre. Here we propose that this was likely due to hydrothermal and/or sediment inputs. Fe(II) oxidation rates were measured in the surface and deep waters. Our study is the first one, to our knowledge, to measure Fe(II) oxidation rates in natural deep seawater at near-ambient concentrations. In the deep waters,  $t_{1/2}$  values were on average 6 times higher than in the surface waters. The comparison of our measured  $t_{1/2}$  with theoretical ones using two different models suggested that organic complexation may strongly influence the oxidation rates, although more studies are needed to better constrain the organic speciation of Fe(II) and its influence on the half-lives of Fe(II). The global data set of Fe(II) also needs to be increased and this will be done in the framework of the GEOTRACES programme and the associated cruises.

### Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Supplementary material related to this article is available online at:  
[http://www.biogeosciences-discuss.net/8/4163/2011/  
bgnd-8-4163-2011-supplement.pdf](http://www.biogeosciences-discuss.net/8/4163/2011/bgnd-8-4163-2011-supplement.pdf).

*Acknowledgements.* We are grateful to Frank Dehairs, the cochief scientist with S. S., the captain and the crew of the R. V. *Marion Dufresne* for their excellent support and commitment onboard. This work was supported by the Institut National des Sciences de L'Univers of the Centre National de la Recherche Scientifique, the French Polar Institute (Institut Polaire Emile Victor), the French Research Institute for Exploitation of the Sea, and the National Agency for Research Funding (ANR-07-BLAN-0146). We thank Josephine Ras and Hervé Claustre for providing pigment data, Damien Cardinal for barium data, and Erika Key for the solar radiation data. Mikael Trapp is also thanked for his help with the Fe(II) oxidation model.

We also acknowledge the Go-Flo sampling team: J. Bown, M. Boyé, F. Lacan, A. Radic, and B. Wake.

## References

- Alldredge, A. L. and Cohen, Y.: Can microscale patches persist in the sea?, *Microelectrode study of marine snow, fecal pellets*, *Science*, 235, 689–601, 1987.
- Arhan, M., Mercier, H., and Park, Y.: On the deep water circulation of the eastern South Atlantic Ocean, *Deep Sea Res. Pt. I*, 50(7), 889–916, 2003.
- Arhan, M., Speich, S., Dencausse, G., Messenger, C., Fine, R., and Boye, M.: Anticyclonic and cyclonic eddies of subtropical origin in the subantarctic zone south of Africa, submitted to *J. Geophys. Res. Oceans*, 2011.
- Barbeau, K., Rue, E. L., Bruland, K. W., and Butler, A.: Photochemical cycling of iron in the surface ocean mediated by microbial iron(III)-binding ligands, *Nature*, 413, 409–413, 2001.
- Bennett, S. A., Achterberg, E. P., Connelly, D. P., Statham, P. J., Fones, G. R., and German, C. R.: The distribution and stabilisation of dissolved Fe in deep-sea hydrothermal plumes, *Earth Planet. Sci. Lett.*, 270(3–4), 157–167, 2008.
- Blain, S., Quéguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C., Brussaard, C., Carlotti, F., Christaki, U., Corbière, A., Durand, I., Ebersbach, F., Fuda,

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


J.-L., Garcia, N., Gerringa, L., Griffiths, B., Guigue, C., Guillerm, C., Jacquet, S., Jeandel, C., Laan, P., Lefèvre, D., Lomonaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.-H., Picheral, M., Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M., Thuiller, D., Timmermans, K., Trull, T., Uitz, J., van-Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L., and Wagener, T.: Impact of natural iron fertilisation on carbon sequestration in the Southern Ocean, *Nature*, 7139, 1070–1074, 2007.

Boebel, O., Lutjeharms, J., Schmid, C., Zenk, W., Rossby, T., and Barron, C.: The Cape Cauldron: a regime of turbulent inter-ocean exchange, *Deep-Sea Res. Pt. II*, 50, 57–86, 2003.

Bowie, A. R., Achterberg, E. P., Sedwick, P. N., Ussher, S., and Worsfold, P. J.: Real-time monitoring of picomolar concentrations of iron(II) in marine waters using automated flow injection chemiluminescence instrumentation, *Environ. Sci. Technol.*, 36, 4600–4607, 2002.

Boyd, P.: Ironing out algal issues in the southern ocean, *Science*, 304(5669), 396–397, 2004.

Boyd, P. W. and Ellwood, M. J.: The biogeochemical cycle of iron in the ocean, *Nat. Geosci.*, 3, 675–682, doi:10.1038/NCEO964, 2010.

Boyd, P. W., Watson, A. J., Law, C. S., Abraham, E. R., Trull, T., Murdoch, R., Bakker, D. C. E., Bowie, A., Buesseler, K. O., Chang, H., Charette, M., Croot, P., Downing, K., Frew, R., Gall, M., Hadfield, M., Hall, J., Harvey, M., Jameson, G., LaRoche, J., Liddicoat, M., Ling, R., Maldonado, M.T., McKay, R. M., Nodder, S., Pickmere, S., Pridmore, R., Rintoul, S., Safi, K., Sutton, P., Strzepek, R., Tanneberger, K., Turner, S., Waite, A., and Zeldis, J.: A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization, *Nature*, 407, 695–702, 2000.

Boye, M., Aldrich, A., van den Berg, C. M. G., de Jong, J. T. M., Veldhuis, M., and de Baar, H. J. W.: Horizontal gradient of the chemical speciation of iron in surface waters of the northeast Atlantic Ocean, *Mar. Chem.*, 80, 129–143, 2003.

Boye, M., Aldrich, A., van den Berg, C. M. G., de Jong, J. T. M., Nirmaier, H., Veldhuis, M., Timmermans, K. R., and de Baar, H. J. W.: The chemical speciation of iron in the north-east Atlantic Ocean, *Deep Sea Res. Pt. I*, 53, 667–683, 2006.

Boyle, E. and Jenkins, W., Hydrothermal Iron in the Deep Western South Pacific, *Geochim. Cosmochim. Ac.*, 72(12), Supplement 1, A107, 2008.

Bruland, K. W. and Rue, E. L.: Analytical Methods for the Determination of Concentrations and Speciation of Iron, edited by: Turner, S. and Hunter, K. A., *Biogeochemistry of Fe in Seawater*, SCOR/IUPAC, 255–290, 2001.

Bucciarelli, E., Sarthou, G., and Chever, F.: Hydrogen peroxide distributions along a transect

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


from the subtropical domain to the Weddell Sea Gyre in the Atlantic sector of the Southern Ocean, in prep., 2011.

Cai, P. H., Chen, W. F., Dai, M. H., Wan, Z. W., Wang, D. X., Li, Q., Tang, T. T., and Lv, D. W.: A high resolution study of particle export in the southern South China Sea based on  $^{234}\text{Th}$ - $^{238}\text{U}$  disequilibrium, *J. Geophys. Res. Ocean.*, 113(C4), C04019, doi:10.1029/2007JC004268, 2008.

Chever, F., Bucciarelli, E., Sarthou, G., Speich, S., Arhan, M., Penven, P., and Tagliabue, A.: Physical speciation of iron in the Atlantic sector of the Southern Ocean, along a transect from the subtropical domain to the Weddell Sea Gyre, *J. Geophys. Res. Ocean.*, 115, C10059, doi:10.1029/2009JC005880, 2010.

Chin, C. S., Coale, K. H., Elrod, V. A., and Johnson, K. S.: In situ observations of dissolved iron and manganese in hydrothermal vent plumes, Juan de Fuca Ridge, *J. Geophys. Res.*, 99(B3), 4969–4984, 1994.

Coale, K. H., Chin, C. S., Massoth, G. J., Johnson, K. S., and Baker, E. T.: In-situ chemical mapping of dissolved iron and manganese in hydrothermal plumes, *Nature*, 352, 325–328, 1991.

Coale, K. H., Johnson, K. S., Fitzwater, S. E., Gordon, R. M., Tanner, S., Chavez, F. P., Ferioli, L., Sakamoto, C., Rogers, P., Millero, F., Steinberg, P., Nightingale, P., Cooper, D., Cochlan, W. P., Landry, M. R., Constantinou, J., Rollwagen, G., Trasvina, A., and Kudela, R.: A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean, *Nature*, 383, 495–501, 1996.

Coale, K. H., Johnson, K. S., Chavez, F. P., Buesseler, K. O., Barber, R. T., Brzezinski, M. A., Cochlan, W. P., Millero, F. J., Falkowski, P. G., Bauer, J. E., Wanninkhof, R. H., Kudela, R. M., Altabet, M. A., Hales, B. E., Takahashi, T., Landry, M. R., Bidigare, R. R., Wang, X. J., Chase, Z., Strutton, P. G., Friederich, G. E., Gorbunov, M. Y., Lance, V. P., Hiltling, A. K., Hiscock, M. R., Demarest, M., Hiscock, W. T., Sullivan, K. F., Tanner, S. J., Gordon, R. M., Hunter, C. N., Elrod, V. A., Fitzwater, S. E., Jones, J. L., Tozzi, S., Koblizek, M., Roberts, A. E., Herndon, J., Brewster, J., Ladizinsky, N., Smith, G., Cooper, D., Timothy, D., Brown, S. L., Selph, K. E., Sheridan, C. C., Twining, B. S., and Johnson, Z. I.: Southern ocean iron enrichment experiment: Carbon cycling in high- and low-Si waters, *Science*, 304(5669), 408–414, 2004.

Croot, P. L. and Laan, P.: Continuous shipboard determination of Fe(II) in polar waters using flow injection analysis with chemiluminescence detection, *Anal. Chim. Acta*, 466(2), 261–



273, 2002.

Croot, P. L., Bowie, A. R., Frew, R. D., Maldonado, M. T., Hall, J. A., Safi, K. A., La Roche, J., Boyd, P. W., and Law, C. S.: Retention of dissolved iron and Fe-II in an iron induced Southern Ocean phytoplankton bloom, *Geophys. Res. Lett.*, 28(18), 3425–3428, 2001.

5 Croot, P. L., Laan, P., Nishioka, J., Strass, V., Cisewski, B., Boye, M., Timmermans, K. R., Bellerby, R. G., Goldson, L., Nightingale, P., and de Baar, H. J. W.: Spatial and temporal distribution of Fe(II) and H<sub>2</sub>O<sub>2</sub> during EisenEx, an open ocean mesocoscale iron enrichment, *Mar. Chem.*, 95, 65–88, 2005.

10 Croot, P. L., Frew, R. D., Sander, S., Hunter, K. A., Ellwood, M. J., Pickmere, S. E., Abraham, E. R., Law, C. S., Smith, M. J., and Boyd, P. W.: Physical mixing effects on iron biogeochemical cycling: Fe-Cycle experiment, *J. Geophys. Res.*, 112(C06015), doi:10.1029/2006JC003748, 2007.

15 Croot, P. L., Bluhm, K., Schlosser, C., Streu, P., Breitbarth, E., Frew, R., and Ardelan, M. V.: Regeneration of Fe(II) during EIFeX and SOFeX, *Geophys. Res. Lett.*, 35, L19606, doi:10.1029/2008GL035063, 2008.

de Baar, H. J. W. and de Jong, J. T. M.: Distributions, Sources and Sinks of Iron in Seawater, in: *Biogeochemistry of Fe in Seawater*, edited: Turner, D. R. and Hunter, K. A., SCOR-UPAC series, J. Wiley, Baltimore, 123–253, 2001.

20 Ducet, N., Traon, P. Y. L., and Reverdin, G.: Global high-resolution mapping of ocean circulation from TOPEX/Poseidon and ERS-1 and -2, *J. Geophys. Res.*, 105(C8), 19477–19498, 2000.

Elrod, V. A., Berelson, W. M., Coale, K. H., and Johnson, K. S.: The flux of iron from continental shelf sediments: A missing source for global budgets, *Geophys. Res. Lett.*, 31, L12307, doi:10.1029/2004GL020216, 2004.

25 Field, M. P. and Sherrell, R. M.: Dissolved and particulate Fe in a hydrothermal plume at 9°45' N, East Pacific Rise: Slow Fe (II) oxidation kinetics in Pacific plumes, *Geochim. Cosmochim. Ac.*, 64(4), 619–628, 2000.

Gerringa, L. J. A., Veldhuis, M. J. W., Timmermans, K. R., Sarthou, G., and de Baar, H. J. W.: Co-variance of dissolved Fe-binding ligands with biological observations in the Canary Basin., *Mar. Chem.*, 102, 276–290, 2006.

30 Gerringa, L. J. A., Blain, S., Laan, P., Sarthou, G., Veldhuis, M. J. W., Viollier, E., and Timmermans, K. R.: Dissolved organic ligands of Fe near the Kerguelen Archipelago in the Southern Ocean (Indian sector), *Deep Sea Res. Pt. II*, 55(5–7), 606–621, 2008.

Gervais, F., Riebesell, U., and Gorbunov, M. Y.: Changes in primary productivity and chlorophyll

**BGD**

8, 4163–4208, 2011

---

**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





---

**Labile Fe(II)  
concentrations in the  
South Atlantic**G. Sarthou et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

a in response to iron fertilization in the Southern Polar Frontal Zone, *Limnol. Oceanogr.*, 47(5), 1324–1335, 2002.

Gladyshev, S., Arhan, M., Sokov, A., and Speich, S.: A hydrographic section from South Africa to the southern limit of the Antarctic Circumpolar Current at the Greenwich meridian, *Deep Sea Res. Pt. I*, 55, 1284–1303, 2008.

Gledhill, M. and van den Berg, C. M. G.: Determination of complexation of iron(III) with natural organic complexing ligands in seawater using cathodic stripping voltammetry, *Mar. Chem.*, 47, 41–54, 1994.

Gledhill, M. and van den Berg, C. M. G.: Measurement of the redox speciation of iron in seawater by catalytic cathodic stripping voltammetry, *Mar. Chem.*, 50(1–4), 51–62, 1995.

Gobler, C. J., Donat, J. R., Consolvo, J. A., and Sanudo, W. S. A.: Physicochemical speciation of iron during coastal algal blooms, *Mar. Chem.*, 77(1), 71–89, 2002.

Godrant, A., Rose, A. L., Sarthou, G., and Waite, T. D.: A new method for the determination of extracellular production of superoxide from phytoplankton cells using the chemiluminescence probes MCLA and the Red-CLA, *Limnol. Oceanogr.-Meth.*, 7, 682–692, 2009.

Gonzalez-Dávila, M., Santana-Casiano, J. M., and Millero, F. J.: Oxidation of iron (II) nanomolar with  $H_2O_2$  in seawater, *Geochim. Cosmochim. Ac.*, 69, 83–93, 2005.

Gonzalez-Dávila, M., Santana-Casiano, J. M., and Millero, F. J.: Competition between  $O_2$  and  $H_2O_2$  in the oxidation of Fe(II) in natural waters, *J. Sol. Chem.*, 35, 95–111, 2006.

Gonzalez-Dávila, M., Santana-Casiano, J. M., Rueda, M. J., Llinás, O., and Gonzalez-Dávila, E. F.: Seasonal and interannual variability of seasurface carbon dioxide species at the European Station for Time Series in the Ocean at the Canary Islands (ESTOC) between 1996 and 2000, *Global Biogeochem. Cy.*, 17(3), 1076, doi:10.1029/2002GB001993, 2003.

Gordon, A. L., Weiss, R. F., Smethie, J. W. M., and Warner, M. J.: Thermocline and intermediate water communication between the South Atlantic and Indian Oceans, *J. Geophys. Res.*, 97(C5), 7223–7240, 1992.

Hansard, S. P., Landing, W. M., Measures, C. I., and Voelker, B. M.: Dissolved iron(II) in the Pacific Ocean: Measurements from the PO2 and P16NCLIVAR/CO2 repeat hydrography expeditions, *Deep Sea Res. Pt. I*, 56, 1117–1129, 2009.

Hansard, S. P., Vermilyea, A. W., and Voelker, B. M.: Measurements of superoxide radical concentration and decay kinetics in the Gulf of Alaska, *Deep-Sea Res. Pt. I*, 57, 1111–1119, 2010.

Hong, H. and Kester, D. R.: Redox state of iron in the offshore waters of Peru, *Limnol.*

---

**Labile Fe(II)  
concentrations in the  
South Atlantic**G. Sarthou et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Oceanogr., 31, 512–524, 1986.

Hopkison, B. M. and Barbeau, K. A.: Organic and Redox speciation of iron in the tropical North Pacific suboxic zone, *Mar. Chem.*, 106, 2–17, 2007.

Hutchins, D. A. and Bruland, K. W.: Grazer-mediated regeneration and assimilation of Fe, Zn, and Mn from planktonic prey, *Mar. Ecol. Prog. Ser.*, 110, 259–269, 1994.

Hutchins, D. A., DiTullio, G. R., and Bruland, K. W.: Iron and regenerated production: Evidence for biological iron recycling in two marine environments, *Limnol. Oceanogr.*, 38(6), 1242–1255, 1993.

Hutchins, D. A., Wang, W., and Fisher, N. S.: Copepod grazing and the biogeochemical fate of diatom iron, *Limnol. Oceanogr.*, 40, 989–994, 1995.

Johnson, K. S., Coale, K.H., Elrod, V. A., and Tindale, N. W.: Iron Photochemistry in seawater from the Equatorial Pacific, *Mar. Chem.*, 46, 319–334, 1994.

Johnson, K. S., Gordon, R. M., and Coale, K. H.: What controls dissolved iron concentrations in the world ocean?, *Mar. Chem.*, 57, 137–161, 1997.

Journet, E., Desboeufs, K. V., Sofikitis, A., Varrault, G., and Colin, J.-L.: In situ speciation of trace Fe(II) and Fe(III) in atmospheric waters by the FZ method coupled to GFAAS analysis, *Int. J. Environ. Anal. Chem.*, 87(9), 647–658, 2007.

Kieber, R.J., Williams, K., Willey, J. D., Skrabal, S., Avery, G., and R. B. J.: Iron speciation in coastal rainwater: concentration and deposition to seawater, *Mar. Chem.*, 73(2), 83–95, 2001.

King, D. W.: Role of carbonate speciation on the oxidation rate of Fe(II) in aquatic systems, *Environ. Sci. Technol.*, 32, 2997–3003, 1998.

King, D. W., Aldrich, R. A., and Charnecki, S. E.: Photochemical redox cycle of iron in NaCl solutions, *Mar. Chem.*, 44, 105–120, 1993.

King, D. W., Lounsbury, H. A., and Millero, F. J.: Rate and mechanism of Fe(II) oxidation at nanomolar total iron concentration, *Environ. Sci. Technol.*, 29(3), 818–824, 1995.

Klatt, O., Fahrbach, E., Hoppema, M., and Rohardt, G.: The transport of the Weddell Gyre across the Prime meridian, *Deep Sea Res. Pt. II*, 52, 513–528, doi:10.1016/j.dsr2.2004.12.015, 2005.

Klunder, M., Laan, P., Middag, R., Baar, H. J. W. D., and Ooijen, J. V.: Dissolved Iron the Southern Ocean (Atlantic Sector), *Deep Sea Res. Pt. II*, in press, doi:10.1016/j.dsr2.2010.10.042, 2010.

Kuma, K., Nakabayashi, S., Suzuki, Y., Kudo, I., and Matsunaga, K.: Photoreduction of Fe(III)

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- by dissolved organic substances and existence of Fe(II) in seawater during spring blooms, *Mar. Chem.*, 37, 15–27, 1992a.
- Kuma, K., Nakabayashi, S., Suzuki, Y., and Matsunaga, K.: Dissolution rate and solubility of colloidal hydrous ferric oxide in seawater, *Mar. Chem.*, 38, 133–143, 1992b.
- 5 Kuma, K., Nishioka, J., and Matsunaga, K.: Controls on iron(III) hydroxide solubility in seawater: the influence of pH and natural organic ligands, *Limnol. Oceanogr.*, 41(3), 396–407, 1996.
- Kustka, A. B., Shaked, Y., Milligan, A. J., King, D. W., and Morel, F. M. M.: Extracellular production of superoxide by marine diatoms: Contrasting effects on iron redox chemistry and bioavailability, *Limnol. Oceanogr.*, 50(4), 1172–1180, 2005.
- 10 Legendre, L., Demers, S., Garside, C., Haugen, E. M., Phinney, D. A., Shapiro, L. P., Therriault, J. C., and Yentsch, C. M.: Circadian photosynthetic activity of natural marine phytoplankton isolated in a tank, *J. Plankton Res.*, 10, 1–6, 1988.
- Lewis, E. and Wallace, D. W. R.: Program Developed for CO<sub>2</sub> System Calculations. ORNL/CDIAC-105, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 1998.
- 15 Liu, X. and Millero, F. J.: The solubility of iron in seawater, *Mar. Chem.*, 77(1), 43–54, 2002.
- Lohan, M. C. and Bruland, K. W.: Elevated Fe(II) and dissolved Fe in hypoxic shelf waters off Oregon and Washington: An enhanced source of iron to coastal upwelling regimes, *Environ. Sci. Technol.*, 42(17), 6462–6468, 2008.
- 20 Maiti, K., Benitez-Nelson, C. R., and Buesseler, K. O.: Insights into particle formation and remineralization using the short-lived radionuclide, Thorium-234, *Geophys. Res. Lett.*, 37, L15608, doi:10.1029/2010GL044063, 2010.
- Maldonado, M. T. and Price, N. M.: Reduction and transport of organically bound iron by *Thalassiosira oceanica* (Bacillariophyceae), *J. Phycol.*, 37(2), 298–309, 2001.
- 25 Maldonado, M. Y. and Price, N. M.: Utilization of iron bound to strong organic ligands by plankton communities in the subarctic Pacific Ocean, *Deep Sea Res. Pt. II*, 46(11–12), 2447–2473, 1999.
- Maldonado, M. T., Allen, A. E., Chong, J. S., Lin, K., Leus, D., Karpenko, N., and Harris, S. L.: Copper-dependent iron transport in coastal and oceanic diatoms, *Limnol. Oceanogr.*, 51(4), 1729–1743, 2006.
- 30 Meredith, M. P., Locarnini, R. A., Scoy, K. A. V., Watson, A. J., Heywood, K. J., and King, B. A.: On the sources of Weddell Gyre Antarctic Bottom Water, *J. Geophys. Res.*, 105(C1), 1093–1104, 2000.

---

**Labile Fe(II)  
concentrations in the  
South Atlantic**G. Sarthou et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Middag, R., Baar, H. J. W. d., Laan, P., Cai, P. H., and van Ooijen, J. C.: Dissolved manganese in the atlantic sector of the Southern Ocean, Deep Sea Res. Pt. II, in press, doi:10.1016/j.dsr2.2010.10.043, 2010.

5 Millero, F. J.: Effect of ionic interactions on the oxidation of iron(II) and copper(I) in natural waters, Mar. Chem., 28, 1–18, 1989.

Millero, F. J. and Sotolongo, S.: The oxidation of Fe(II) with H<sub>2</sub>O<sub>2</sub> in seawater, Geochim. Cosmochim. Ac., 53, 1867–1873, 1989.

Millero, F. J., Sotolongo, S., and Izaguirre, M.: The oxidation kinetics of Fe(II) in seawater, Geochim. Cosmochim. Ac., 51, 793–801, 1987.

10 Mintrop, L., Pérez, F. F., González Dávila, M., Körtzinger, A., and Santana-Casiano, J. M.: Alkalinity determination by potentiometry: intercalibration using three different methods, Cien. Mar., 26, 23–37, 2000.

Moffett, J., Goepfert, T. J., and Naqvi, W. A.: Reduced iron associated with secondary nitrite maxima in the Arabian Sea, Deep Sea Res. Pt. I, 54, 1341–1349, 2007.

15 Morel, F. M. M., Kustka, A. B., and Shaked, Y.: The role of unchelated Fe in the iron nutrition of phytoplankton, Limnol. Oceanogr., 53(1), 400–404, 2008.

O'Sullivan, D. W., Hanson, A. K., Miller, W. L., and Kester, D. R.: Measurement of Fe(II) in equatorial Pacific surface seawater, Limnol. Oceanogr., 36(8), 1727–1741, 1991.

20 Obata, H., Karatani, H., and Nakayama, E.: Automated determination of iron in seawater by chelating resin concentration and chemiluminescence, Anal. Chem., 65, 1524–1528, 1993.

Orsi, A. H., Nowlin, W. D., and Whitworth III, T.: On the circulation and stratification of the Weddell Gyre, Deep Sea Res. Pt. I, 40, 169–203, 1993.

Ozsoy, T. and Saydam, A. C.: Iron speciation in precipitation in the north-eastern Mediterranean and its relationship with Sahara dust, J. Atmos. Chem., 40, 41–76, 2001.

25 Planchon F., Cavagna A. J., Cardinal, D., and Dehairs, F.: Dynamics of biogenic particles in the Southern Atlantic Ocean as revealed by <sup>234</sup>Th proxy along the Bonus-Good-Hope transect from Cape Basin to Northern Weddell gyre, submitted to Biogeosciences, 2011.

Pollard, R., Sanders, R., Lucasa, M., and Statham, P.: The Crozet Natural Iron Bloom and Export Experiment (CROZEX), Deep Sea Res. Pt. II, 54(18–20), 1905–1914, 2007.

30 Rich, H. W. and Morel, F. M. M.: Availability of well-defined iron colloids to the marine diatom *Thalassiosira weissflogii*, Limnol. Oceanogr., 35(3), 652–662, 1990.

Rijkenberg, M. J. A., Gerringa, L. J. A., Carolus, V. E., Velzeboer, I., and de Baar, H. J. W.: Enhancement and inhibition of iron photoreduction by individual ligands in open ocean sea-

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



water, *Geochim. Cosmochim. Ac.*, 70, 2790–2805, 2006.

Rose, A. L. and Waite, T. D.: Chemiluminescence of luminol in the presence of iron(II) and oxygen: Oxidation mechanism and implications for its analytical use, *Anal. Chem.*, 73, 5909–5920, 2001.

Rose, A. L. and Waite, T. D.: Kinetic model for Fe(II) oxidation in seawater in the absence and presence of natural organic matter, *Environ. Sci. Technol.*, 36, 433–444, 2002.

Rose, A. L. and Waite, T. D.: Predicting iron speciation in coastal waters from the kinetics of sunlight-mediated iron redox cycling, *Aquat. Sci.*, 65, 375–383, 2003.

Rose, A. L., Salmon, T. P., Lukondeh, T., Neilan, B. A., and Waite, T. D.: Use of Superoxide as an Electron Shuttle for Iron Acquisition by the Marine Cyanobacterium *Lyngbya majuscula*, *Environ. Sci. Technol.*, 39(10), 3708–3715, doi:10.1021/es048766c, 2005.

Roy, E. G., Wells, M. L., and King, D. W.: Persistence of iron(II) in surface waters of the western subarctic Pacific, *Limnol. Oceanogr.*, 53, 89–98, 2008.

Rue, E. L. and Bruland, K. W.: The role of organic complexation on ambient iron chemistry in the equatorial Pacific Ocean and the response of a mesoscale iron addition experiment, *Limnol. Oceanogr.*, 42(5), 901–910, 1997.

Rutgers van der Loeff, M. M., Cai, P. H., Stimac, I., Bracher, A., Middag, R., Klunder, M. B., and Van Heuven, S.:  $^{234}\text{Th}$  in surface waters: distribution of particle export flux across the Antarctic Circumpolar Current and in the Weddell Sea during the GEOTRACES expedition ZERO and DRAKE, *Deep-Sea Res. Pt. II*, in press, doi:10.1016/j.dsr2.2011.02.004, 2011.

Salmon, T. P., Rose, A. L., Neilan, B. A., and Waite, T. D.: The FeL model of iron acquisition: Nondissociative reduction of ferric complexes in the marine environment, *Limnol. Oceanogr.*, 51(4), 1744–1754, 2006.

Santana-Casiano, J. M., Gonzalez-Davila, M., and Millero, F. J.: The oxidation of Fe(II) in NaCl- $\text{HCO}_3^-$  and seawater solutions in the presence of phthalate and salicylate ions: a kinetic model, *Mar. Chem.*, 85 (1–2), 27–40, 2004.

Santana-Casiano, J. M., Gonzalez-Davila, M., and Millero, F. J.: Oxidation of nanomolar levels of Fe(II) with oxygen in natural waters, *Environ. Sci. Technol.*, 39, 2073–2079, 2005.

Santana-Casiano, J. M., Gonzalez-Davila, M., and Millero, F. J.: The role of Fe(II) species on the oxidation of Fe(II) in natural waters in the presence of  $\text{O}_2$  and  $\text{H}_2\text{O}_2$ , *Mar. Chem.*, 99, 70–82, 2006.

Sarma, V. V. S. S., Kumar, M. D., and Saino, T.: Impact of sinking carbon flux on accumulation of deep-ocean carbon in the Northern Indian Ocean, *Biogeochem.*, 82, 89–100,

doi:10.1007/s10533-006-9055-1, 2007.

Sarthou, G., Jeandel, C., Brisset, L., Amouroux, D., and Donard, O. F. X.: Fe and H<sub>2</sub>O<sub>2</sub> distributions in the upper 250 meters of the water column in the Indian sector of the Southern Ocean, *Earth Planet. Sci. Lett.*, 147, 83–92, 1997.

5 Sarthou, G., Baker, A. R., Blain, S., Achterberg, E. P., Boye, M., Bowie, A. R., Croot, P. L., Laan, P., de Baar, H. J. W., Jickells, T. D., and Worsfold, P. J.: Atmospheric iron deposition and sea-surface dissolved iron concentrations in the eastern Atlantic Ocean, *Deep Sea Res. Pt. I*, 50(10–11), 1339–1352, 2003.

10 Sarthou, G., Vincent, D., Christaki, U., Obernosterer, I., Timmermans, K. R., and Brussaard, C. P. D.: The fate of biogenic iron during a phytoplankton bloom induced by natural fertilization: impact of copepod grazing, *Deep Sea Res. Pt. II*, 55, 734–751, 2008.

Shaked, Y., Kustka, A. B., Morel, F. M. M., and Erel, Y.: Simultaneous determination of iron reduction and uptake by phytoplankton, *Limnol. Oceanogr.-Meth.*, 2, 137–145, 2004.

15 Speich, S., Arhan, M., Gladyshev, S., Rupolo, V. and Perrot, X.: Upper layer structure and dynamics along BGH, submitted to *Ocean Sci.*, 2011.

Statham, P. J., German, C. R. M. and Connelly, D. P.: Iron(II) distribution and oxidation kinetics in hydrothermal plumes at the Kairei and Edmond vent sites, *Indian Ocean, Earth Planet. Sci. Lett.*, 236, 588–596, 2005.

20 Sunda, W. G.: Trace metal interactions with marine phytoplankton, *Biol. Oceanogr.*, 6, 411–442, 1989.

Tagliabue, A., Bowie, A., Chever, F., Baptiste, P.-J., Dutay, J.-C., Bucciarelli, E., Lannuzel, D., Remenyi, T., Sarthou, G., Aumont, O., Gehlen, M., and Bopp, L.: On the importance of hydrothermalism to the oceanic dissolved iron inventory, *Nat. Geosci.*, 3, 252–256, doi:10.1038/NGEO818, 2010.

25 Thuróczy, C.-E., Gerringa, L. J. A., Klunder, M. B., Middag, R., Laan, P., Timmermans, K. R., and de Baar, H. J. W.: Speciation of Fe in the Eastern North Atlantic Ocean, *Deep Sea Res. Pt. I*, 57(11), 1444–1453, 2010.

Trapp, J. M. and Millero, F. J.: The oxidation of iron(II) with oxygen in NaCl brines, *J. Sol. Chem.*, 36, 1479–1493, 2007.

30 Tsuda, A., Takeda, S., Saito, H., Nishioka, J., Nojiri, Y., Kudo, I., Kiyosawa, H., Shiimoto, A., Imai, K., Ono, T., Shimamoto, A., Tsumune, D., Yoshimura, T., Aono, T., Hinuma, A., Kinugasa, M., Suzuki, K., Sohrin, Y., Noiri, Y., Tani, H., Deguchi, Y., Tsurushima, N., Ogawa, H., Fukami, K., Kuma, K., and Saino, T.: A mesoscale iron enrichment in the western Subarctic

**BGD**

8, 4163–4208, 2011

---

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pacific induces a large centric diatom bloom, *Science*, 300, 958–961, 2003.
- Ussher, S. J., Yaqoob, M., Achterberg, E. P., Nabi, A., and Worsfold, P. J.: Effect of model ligands on iron redox speciation in natural waters using flow injection with luminol chemiluminescence detection, *Anal. Chem.*, 77(7), 1971–1978, 2005.
- 5 Ussher, S. J., Worsfold, P. J., Achterberg, E. P., Laes, A., Blain, S., Laan, P., and de Baar, H. J. W.: Distribution and redox speciation of dissolved iron on the European continental margin, *Limnol. Oceanogr.*, 52(6), 2530–2539, 2007.
- Voelker, B. M. and Sedlak, D. L.: Iron reduction by photoproduct superoxide in seawater, *Mar. Chem.*, 50(1–4), 93–102, 1995.
- 10 Waite, T. D. and Morel, F. M. M.: Coulometric study of the redox dynamics of iron in seawater, *Anal. Chem.*, 56, 787–792, 1984.
- Waite, T. D., Szymczak, R., Espey, Q. I., and Furnas, M. J.: Diel variations in iron speciation in northern Australian shelf waters, *Mar. Chem.*, 50(1–4), 79–92, 1995.
- Wells, M. L., Mayer, L. M., Donard, O. F. X., de Souza Sierra, M. M., and Ackelson, S. G.: The photolysis of colloidal iron in the oceans, *Nature*, 353, 248–250, 1991.
- 15 Whitworth III, T. and Nowlin, J. W. D.: Water masses and currents of the Southern Ocean at the Greenwich Meridian, *J. Geophys. Res.*, 96(C6), 6462–6476, 1987.
- Windom, H. L., Moore, W. S., Niencheski, L. F. H., and Jahnke, R. A.: Submarine groundwater discharge: a large, previously unrecognized source of dissolved iron to the South Atlantic Ocean, *Mar. Chem.*, 102, 252–266, 2006.
- 20 Yuan, J. and Shiller, A. M.: Determination of Subnanomolar Levels of Hydrogen Peroxide in Seawater by Reagent-Injection Chemiluminescence Detection, *Anal. Chem.*, 71, 1975–1980, 1999.

**BGD**

8, 4163–4208, 2011

---

**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Location of the stations sampled during the cruise in relation to the domains and fronts crossed. Mixed layer depth observed from the vertical profile of temperature for nearby CTD station, as well as day and time of sampling are indicated. Dates are in DD/MM/YYYY format (D = day, M = month and Y = year).

Oceanographic Domain	Fronts crossed	Station	Position	MLD (m)	Day of sampling	Time of sampling
Subtropical		L1	34.43° S, 14.40° E	50–60	17/02/2008	10:50
		S1	36.50° S, 13.10° E	40–50	20/02/2008	00:15
		L2	41.18° S, 09.92° E	25	25/02/2008	23:15
ACC	S-STF		42.2° S			
		S2	42.47° S, 08.93° E	50–80	27/02/2008	18:30
	SAF		44.2° S			
		L3	44.88° S, 06.88° E	60–80	01/03/2008	17:45
		L4	46.02° S, 05.87° E	80	03/03/2008	05:40
		S3	47.55° S, 04.37° E	80–100	05/03/2008	15:50
		L5	49.03° S, 02.84° E	100–110	07/03/2008	16:30
	PF		50.2° S			
	SACCF		51.5° S			
		S4	51.85° S, 00.00° E	120–150	10/03/2008	12:10
Eastern part of the Weddell Sea Gyre (EWSG)		L7	55.23° S, 00.03° E	80–110	13/03/2008	22:30
	Sbdy		55.5° S			
		S5	57.55° S, 00.03° W	100	16/03/2008	16:00



**Table 2.** Labile Fe(II) concentrations and percentage of labile Fe(II) over dissolved Fe (DFe, Chever et al., 2010). Uncertainties on the concentrations correspond to standard deviation of a same sample measured 3 times. EWSG = eastern part of the Weddell Sea Gyre. nd = not determined, when no DFe data were available.

Domain	Station	Position	Bottom depth (m)	Depth (m)	Fe(II) (nM)	STD (nM)	Fe(II)/DFe (%)	STD (%)			
Subtropical	L1	34.43° S, 14.40° E	4505	20	0.017	0.001	7.2	0.6			
				40	0.016	0.001	3.2	0.2			
				60	0.025	0.001	nd				
				80	0.063	0.003	30.1	1.8			
				200	0.076	0.004	10.2	0.5			
				700	0.033	0.002	2.8	0.2			
				800	0.049	0.002	7.7	0.5			
				1000	0.035	0.002	5.4	0.3			
				1200	0.017	0.001	2.7	0.2			
				2100	0.014	0.001	2.1	0.1			
				S1	36.50° S 13.10° E	4915	20	0.038	0.002	5.5	0.3
							30	0.039	0.002	5.5	0.4
							40	0.099	0.005	nd	
							70	0.107	0.005	nd	
							300	0.112	0.006	nd	
							500	0.107	0.005	nd	
							700	0.091	0.005	nd	
1000	0.039	0.002	3.9				0.3				
1200	0.029	0.001	3.1				0.2				
1400	0.015	0.001	2.0				0.1				
1600	0.009	0.000	1.3				0.3				
2000	0.009	0.000	nd								
2700	0.009	0.000	1.4				0.1				
3050	0.009	0.000	1.0	0.0							
3500	0.009	0.000	1.6	0.1							
3800	0.009	0.000	0.8	0.0							
L2	41.18° S 09.92° E	4525	4000	0.010	0.000	0.6	0.0				
			15	0.040	0.002	25.4	1.3				
			35	0.009	0.000	nd					
			45	0.009	0.000	1.5	0.2				
			95	0.094	0.005	32.3	3.1				
			300	0.102	0.005	nd					
			600	0.047	0.002	6.4	0.4				
			800	0.025	0.001	2.3	0.2				
			1200	0.022	0.001	2.7	0.2				
			1400	0.015	0.001	2.0	0.2				
			2100	0.009	0.000	1.0	0.1				

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 2.** Continued.

Domain	Station	Position	Bottom depth (m)	Depth (m)	Fe(II) (nM)	STD (nM)	Fe(II)/DFe (%)	STD (%)
ACC	S2	42.47° S 08.93° E	4070	15	0.024	0.001	13.3	1.6
				30	0.019	0.001	14.6	2.1
				35	0.021	0.001	21.5	2.8
				45	0.116	0.006	nd	
				196	0.125	0.007	70.4	5.8
				314	0.105	0.006	62.3	4.9
				461	0.102	0.006	nd	
				598	0.093	0.005	nd	
				809	0.088	0.005	20.2	1.3
				1029	0.078	0.004	18.0	1.3
				1250	0.071	0.004	15.5	1.1
				1441	0.051	0.003	7.9	0.5
				1764	0.032	0.002	5.0	0.3
				2156	0.033	0.002	3.2	0.2
				2548	0.027	0.001	3.4	0.2
				2891	0.020	0.001	1.4	0.1
				3234	0.025	0.001	3.2	0.3
				3626	0.020	0.001	2.8	0.2
				3940	0.023	0.001	3.7	0.2
				L3	44.88° S, 06.88° E	4315	30	0.043
100	0.019	0.002	9.3				1.7	
150	0.017	0.002	14.4				2.7	
270	0.023	0.002	11.6				1.4	
400	0.016	0.002	4.7				0.7	
600	0.013	0.001	3.1				0.4	
1200	0.009	0.001	0.9				0.1	
1400	0.009	0.001	1.1				0.1	
2100	0.011	0.001	1.8				0.2	
L4	46.02° S 05.87° E	4147	30				0.035	0.002
			60	0.030	0.002	nd		
			100	0.046	0.002	22.3	2.3	
			150	0.023	0.001	10.1	1.1	
			270	0.029	0.001	10.4	0.7	
			480	0.024	0.001	6.3	0.7	
			800	0.018	0.001	5.3	0.3	
			1600	0.018	0.001	2.5	0.2	
			2050	0.016	0.001	2.1	0.2	
			S3	47.55° S 04.37° E	4480	20	0.065	0.003
30	0.066	0.003				38.7	2.8	
40	0.032	0.002				nd		
70	0.021	0.001				11.5	1.4	
100	0.018	0.001				9.6	1.6	
200	0.048	0.002				33.8	3.9	
300	0.063	0.003				22.6	1.4	
450	0.055	0.003				19.1	1.0	
600	0.050	0.002				12.2	0.9	
800	0.041	0.002				7.3	0.5	
1070	0.037	0.002	5.6	0.4				
1500	0.028	0.001	4.6	0.3				
2020	0.018	0.001	1.6	0.1				

**Labile Fe(II) concentrations in the South Atlantic**

G. Sarthou et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 2.** Continued.

Domain	Station	Position	Bottom depth (m)	Depth (m)	Fe(II) (nM)	STD (nM)	Fe(II)/DFe (%)	STD (%)
				2500	0.015	0.001	1.9	0.2
				3000	0.010	0.001	1.7	0.1
				3500	0.009	0.000	1.2	0.1
				3980	0.010	0.001	1.3	0.1
	L5	49.03° S 02.84° E	4025	40	0.018	0.001	12.6	1.4
				80	0.060	0.001	nd	
				150	0.065	0.002	49.2	1.7
				170	0.087	0.002	nd	
				250	0.055	0.002	25.6	1.2
				350	0.048	0.003	13.6	1.4
				700	0.036	0.004	6.0	1.0
				1000	0.039	0.003	7.7	0.7
				1600	0.026	0.003	5.5	0.8
				2200	0.023	0.001	3.2	0.2
	L6	50.38° S 01.33° E	3576	30	0.039	0.002	15.8	1.8
				60	0.047	0.002	nd	
				100	0.021	0.001	9.6	1.5
				135	0.015	0.001	nd	
				180	0.018	0.001	8.4	0.8
				300	0.018	0.001	4.3	1.0
				600	0.014	0.001	1.6	0.1
				850	0.012	0.001	2.8	0.2
				1600	0.010	0.000	2.3	0.2
				2100	0.011	0.001	1.0	0.1
	S4	51.85° S 00.00° E	2632	30	0.116	0.006	63.5	4.6
				60	0.081	0.004	nd	
				130	0.060	0.003	44.4	4.8
				160	0.064	0.003	50.6	4.9
				180	0.054	0.003	29.4	1.8
				250	0.066	0.003	34.5	2.6
				300	0.050	0.003	24.9	2.7
				350	0.060	0.003	29.6	1.9
				400	0.047	0.002	21.9	1.6
				500	0.057	0.003	12.8	0.7
				700	0.035	0.002	8.3	0.5
				900	0.049	0.002	nd	
				1117	0.026	0.001	5.6	0.3
				1950	0.021	0.001	nd	
				2300	0.012	0.001	nd	
				2500	0.009	0.000	1.1	0.1
	L7	55.23° S 00.03° E	2770	30	0.016	0.001	14.6	1.0
				60	0.012	0.001	18.7	2.1
				100	0.017	0.001	26.0	3.7
				120	0.024	0.001	26.1	2.4
				200	0.037	0.002	17.4	1.1
				300	0.031	0.002	8.3	0.5
				650	0.020	0.001	4.5	0.2
				1000	0.012	0.001	2.5	0.2
				1500	0.019	0.001	4.0	0.3

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Table 2. Continued.

Domain	Station	Position	Bottom depth (m)	Depth (m)	Fe(II) (nM)	STD (nM)	Fe(II)/DFe (%)	STD (%)
EWSG	S5	57.55° S 00.03° W	3932	30	0.063	0.003	66.6	8.4
				60	0.042	0.002	44.4	2.5
				120	0.017	0.001	8.3	0.6
				140	0.018	0.001	24.4	3.4
				190	0.014	0.001	16.0	1.0
				250	0.025	0.001	22.5	1.8
				350	0.011	0.001	11.8	0.7
				550	0.014	0.001	12.9	1.0
				750	0.021	0.001	4.4	0.5
				800	0.024	0.001	7.1	0.5
				1250	0.009	0.000	2.2	0.1
				1700	0.009	0.000	3.5	0.2
				2150	0.035	0.002	12.2	0.7
				2600	0.036	0.002	10.0	0.9
3050	0.047	0.002	11.2	0.6				
3500	0.050	0.003	13.1	0.8				
3840	0.044	0.002	8.5	0.8				

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 3.** Theoretical values of Fe(II) half lives (in min) for surface and deep water samples at 4 °C using models of Fe(II) oxidation kinetics (Santana-Casiano et al., 2005: Model I; Trapp and Millero, 2007: Model II; see Supplement). The ionic strength has been calculated from salinity using the equation  $I = (19.9201 * S) / (1000 - 1.00488 * S)$  (Millero, 1995).

Stations	Latitude (° S)	<i>I</i> (M)	[O <sub>2</sub> ] (µM)	[H <sub>2</sub> O <sub>2</sub> ] (nM)	pH	[CO <sub>3</sub> <sup>2-</sup> ] (µM)	Theoretical values		Measured
							<i>t</i> 1/2 (min)	<i>t</i> 1/2 (min)	values <i>t</i> 1/2 (min)
Upper waters									
L1	34.43	0.74	233.7	27.0	8.09	216.1	7.5	5.5	2.9
S1	36.50	0.73	245.4	19.6	8.12	209.1	7.7	5.6	2.9
L2	41.18	0.71	251.3	2.4	7.99	110.3	18.4	13.6	3.8
S2	42.47	0.71	273.5	2.1	8.04	117.9	15.4	11.4	7.6
L3	44.88	0.70	293.2		8.07	128.0	12.6	9.7	6.4
L4	46.02	0.70	300.9	21.5	8.07	125.0	11.3	9.0	6.2
S3	47.55	0.70	311.3	15.1	8.08	119.8	12.0	9.4	9.0
L5	49.03	0.70	313.5	19.8	8.07	120.0	11.6	9.2	8.3
L6	50.38	0.70	326.2	22.6	8.08	112.1	12.2	9.6	5.9
S4	51.85	0.69	340.7	9.0	8.05	106.1	13.4	10.4	8.8
L7	55.23	0.70	349.2	31.1	8.06	98.8	13.1	10.3	7.0
S5	57.55	0.70	353.3	31.5	8.06	93.9	14.1	10.9	11.3
Deep waters									
S1	36.50	0.72	218.0		7.93	90.5	91.1	20.4	43.8
L2	41.18	0.71	187.7	1.1	7.85	74.6	127.0	30.8	63.4
S2	42.47	0.71	185.4		7.86	75.3	132.6	31.0	11.8
S2	42.47	0.72	227.8		7.88	91.3	85.6	19.2	14.6
L3	44.88	0.71	185.5		7.85	76.2	130.5	30.5	10.0
L4	46.02	0.72	187.0		7.87	85.5	114.8	26.1	39.2
S3	47.55	0.71	181.2		7.86	74.5	137.6	32.3	36.7
S3	47.55	0.72	218.7		7.84	81.7	101.3	23.3	37.6
L5	49.03	0.71	180.5	1.4	7.85	77.6	125.5	30.3	28.2
S4	51.85	0.71	183.9		7.88	79.5	126.7	29.2	33.6
S4	51.85	0.72	218.5		7.87	84.0	98.8	22.6	40.5
L7	55.23	0.72	201.9	6.3	7.92	84.4	89.9	23.4	30.9
S5	57.55	0.72	209.1		7.91	76.8	115.8	26.8	72.3
S5	57.55	0.72	252.4		7.85	79.7	89.8	20.8	47.2

**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 4.** Fe(II) concentration ranges in previous published studies.

Latitude/ Longitude	Environments/ Experiments	Depth range (m)	[Fe(II)] range	Reference
33.43° S–57.55° S/14.40° E– 00.03° W	Open Ocean South Atlantic/Southern Ocean	15–4000 m	9–125 pM	This study
30° N/135° E–118° W 14° S–56° N/152° W	Open Ocean Pacific Ocean and northern Philippine Sea	13–1010 m	<12–280 pM <12–76 pM	Hansard et al. (2009)
3.0° S–9.1° N/140° W	Open Ocean Equatorial Pacific	0–100 m	<120–530 pM	O'Sullivan et al. (1991)
178.72° E/46.24° S	Open Ocean, South West Pacific, FeCycle experiment.	2 m	Up to 46 pM (during night- time)	Croot et al. (2007)
23°18'S–24°48'S/8°39'E– 9°59'E	Open Ocean, South Atlantic	1–2 m	<12–45 pM	Bowie et al. (2002)
50°92'S–51°25'S/143°38'E– 143°03'E	Open Ocean, Southern Ocean, Subantarctic Front	1–2 m	<12–29 pM	Bowie et al. (2002)
42–51° N/23° E–2° W	Eastern North Atlantic, Euro- pean continental shelf and English Channel	2 m	<160 pM (oceanic waters) Up to 250 pM (shelf waters) 500–1800 pM (coastal wa- ters).	Boye et al. (2003)
37–42° N/23° W	Eastern North Atlantic	0–2000 m	<100–550 pM	Boye et al. (2006)
46° N–52.4° N/ 8° W–4.3° E	European continental margin. Open Ocean and shelf waters	3–4000 m	<12 → 200 pM	Ussher et al. (2007)
9.02° S–12.27° S/127.43° E– 144.19° E	Northern Australian shelf waters	2–3 m	Up to 3 nM	Waite et al. (1995)
31.53° N–h56.50° N/ 0.39° E–0.83° E	Northern North Sea	0–70 m	Up to 1.2 nM	Gledhill and van den Berg (1995)
9.5° S–10.9° S/78.1° W–79.1° W	Suboxic zone, near the coast of Peru	0–2300 m	Up to 40 nM	Hong and Kester (1986)
17° N–23.5° N/ 57° E–74° E	Suboxic zone, Arabian Sea	0–1000 m	Up to 600 pM	Moffett et al. (2007)
15° N–18° N/ 105° W–115° W	Oxic-suboxic zone, Eastern Tropi- cal North Pacific	0–300 m	Up to 150 pM	Hopkison and Barbeau (2007)
61° S/140° E	Southern Ocean SOIREE fertili- zation experiment	2–3 m	Up to 1 nM	Croot et al. (2001)
48° S/21° E	Southern Ocean EISENEX fertili- zation experiment	0–100 m	Up to 1 nM	Croot and Laan (2002); Croot et al. (2005)
56.2° S–6–6° S/172° W	Southern Ocean SOFEX fertili- zation experiment	2–3 m	Up to 300 pM in patch	Croot et al. (2008)
50° S/2° E	Southern Ocean EIFEX fertili- zation experiment	2–3 m	Up to 800 pM in patch	Croot et al. (2008)
46.7° N/165.8° E	Sub-Arctic Pacific Ocean SEEDS II fertilization experiment	0–80 m	UP to > 200 pM in patch	Roy et al. (2008)

**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 1.** Location of the stations sampled during the BONUS-GoodHope cruise along with the three main oceanographic provinces encountered. The three domains crossed were the subtropical domain (stations L1, S1 and L2), the ACC domain (stations S2, L3, L4, S3, L5, L6, S4 and L7) and the eastern part of the Weddell Sea Gyre (station S5). Five fronts were crossed: the southern-subtropical front (S-STF), the sub Antarctic front (SAF), the polar front (PF), the southern ACC front (SACCF) and the southern boundary (Sbdy).

**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion








(a)



(b)



(c)

**Fig. 2.** Vertical distribution of salinity (a), theta (b), and oxygen (c) measured along the transect from the north (left) to the south (right) of the section. Water masses are indicated: AAIW: Antarctic Intermediate Water. This water mass is coming from the Indian Ocean through the Agulhas Current (I-AAIW) north of  $\sim 37^\circ$  S and from the Atlantic sector (A-AAIW) south of  $37^\circ$  S (Gordon et al., 1992). NADW: North Atlantic Deep Water. The highest salinity values close to the African continental slope reflect advection by a southeastward deep boundary current (SE-NADW, Arhan et al., 2003). LCDW: Lower Circumpolar Deep Water and AABW: Antarctic Bottom Water.

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion






(a)




(b)

**Fig. 3.** Vertical section of **(a)** labile Fe(II) in nM and **(b)** %Fe(II)/DFe from the north (left) to the south (right) of the section. DFe data are from Chever et al. (2010).



(a)




(b)


**Fig. 4.** Natural logarithm transformation of the chemiluminescent integral vs. time for Fe(II) spikes (0.25–1 nM) in a surface sample **(a)** and a deep one **(b)**. See text for more details.

## Labile Fe(II) concentrations in the South Atlantic


G. Sarthou et al.



(a)



(b)



(c)

**Fig. 5.** Measured Fe(II) half-lives at 4 °C in minutes for **(a)** surface waters (20–300 m), **(b)** both surface and deep waters (500–2000 m), and **(c)** surface, 500–2000 m deep and 2300–3600 m deep waters. The bars represent the mean value calculated from the four Fe spikes and the error bars the standard deviation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 6.** Percentage of labile Fe(II) relative to dissolved Fe concentrations in the surface mixed layer for all the stations vs. the time at which sampling was done.

**Labile Fe(II)  
concentrations in the  
South Atlantic**

G. Sarthou et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 7.** Vertical profiles of  $^{234}\text{Th}/^{238}\text{U}$ , labile Fe(II) and  $\text{Ba}_{\text{xs}}$  at station S2.  $\text{Ba}_{\text{xs}}$  is the biogenic particulate Ba calculated by subtracting the lithogenic Ba from total particulate Ba.

## Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Labile Fe(II) concentrations in the South Atlantic

G. Sarthou et al.



**Fig. 8.** Vertical profiles in the upper 1500 m of labile Fe(II) and  $T$  at the 4 stations south of the PF where the AAWW were clearly seen.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

