

THE SUBTIDAL EPIBENTHIC COMMUNITIES OFF PUERTO DEL CARMEN (LANZAROTE, CANARY ISLANDS)

C. NIKE BIANCHI, RICARDO HAROUN, CARLA MORRI & PETER WIRTZ



BIANCHI, C. N., R. HAROUN, C. MORRI & P. WIRTZ 2000. The subtidal epibenthic communities off Puerto del Carmen (Lanzarote, Canary Islands). *Arquipélago. Life and Marine Sciences*. Supplement 2(Part A): 145-155. Ponta Delgada. ISSN 0873-4704.

SCUBA diving down to ca 50 m depth in February 1996 allowed the recording of 150 conspicuous species (or species groups) by visual census and photography. Five main communities were identified using cluster analysis on presence / absence data. Rock in shallow water (< 10 m) was covered by a species-rich community dominated by photophilic algae. A community characterised by several fish species inhabited sandy bottom at a depth 10 to 20 m deep. Between 20 and 50 m, a volcanic reef ran almost parallel to the coastline. A community characterised chiefly by echinoderms, encrusting invertebrates and few algae thrived on the reef head, whereas a community of sessile animals (sponges, cnidarians etc.) was recognisable on the reef wall and in caves. A further community was found on rocks at the base of the reef, and was constituted mainly by bushy cnidarians. Species composition and general physiognomy of these five epibenthic communities were similar to those already described from other sites of the Canaries and showed analogies with corresponding communities in the rest of the Atlantic-Mediterranean region.

C. Nike Bianchi, Marine Environment Research Centre, ENEA Santa Teresa, PO Box 316, IT-19100 La Spezia (Italy). (e-mail: bianchi@estof.santateresa.enea.it) - Ricardo Haroun, Dpto Biología, Fac. Ciencias del Mar, Univ. Las Palmas de Gran Canaria, ES-35017 Las Palmas (Spain). (e-mail: ricardo.haroun@biologia.ulpgc.es) - Carla Morri, Dipartimento per lo studio del Territorio e delle sue Risorse (Zoologia), Università di Genova, via Balbi 5, IT-16126 Genova (Italy). (e-mail: zoologia@igecuniv.csita.unige.it) - Peter Wirtz, Centro de Ciências Biológicas e Geológicas, Universidade da Madeira, Largo do Colégio, PT-9000 Funchal (Portugal).

INTRODUCTION

The lack of a comprehensive and structured classification system of the European marine habitats, such as that developed for the terrestrial habitats (COMMISSION OF THE EUROPEAN COMMUNITIES 1991), has been highlighted by CONNOR et al. (1995). Much reference work is available for the Mediterranean Sea (AUGIER 1982; BELLAN-SANTINI et al. 1994), and PÉRÈS (1982) extended to a worldwide scale his still largely employed "Nouveau manuel de bionomie benthique de la Mer Méditerranée" (PÉRÈS & PICARD 1964). Reference is also available for the northern European seas (GLÉMAREC 1973;

HISCOCK & MITCHELL 1980), whereas little exists for the Eastern Atlantic.

The Canary Islands are an important part of the Eastern Atlantic, because of their geographic position and the relationships within both the "Macaronesia" puzzle (BEYHL et al. 1995) and the Atlantic-Mediterranean region (TORTONESE 1960).

The marine flora and fauna of the Canaries are sufficiently known (BACALLADO ARÁNEGA 1984; GONZÁLES HENRÍQUEZ et al. 1986) and guides to invertebrates and fish are available (BRITO 1991; PÉREZ SÁNCHEZ & MORENO BATET 1991; WIRTZ 1994, 1995), but comparatively few studies have been undertaken on the classification of marine communities. Most of these studies deal with

algal vegetation (*e. g.*, LAWSON & NORTON 1971; HAROUN TABRAUE *et al.* 1984; BALLESTEROS 1993; MEDINA *et al.* 1995), but HERRERA *et al.* (1993) and ARÍSTEGUI *et al.* (1987) examined whole epibenthic communities.

The aim of this paper is to characterise the main epibenthic communities and their zonation according to bottom types off Puerto del Carmen, a small town on the SE coast of the Island of Lanzarote, the easternmost of the Canaries.

METHODS

We used visual census and underwater photography by SCUBA diving down to ca 50 m depth. Four sites were investigated (Fig. 1): 1) El Agujero, 2) Bajamar, 3) Punta Tiñosa; 4) El Muelle. The first three sites were explored by SCUBA diving along depth profiles (Fig. 2). Bajamar was visited four times, to explore the deep reef in detail (Fig. 3). El Muelle (Fig. 4) is a shallow water area near the dock, and dives were conducted by simply wandering around over the bottom. Depths were measured with a diving computer, and subsequently corrected to chart datum.

In each site, 2 to 5 different stations were identified physiognomically (BIANCHI *et al.* 1991), *i. e.*, a "station" was a relatively large area homogenous in aspect. In each of such stations, the "conspicuous" (HISCOCK 1987) species of flora and fauna were recorded and photographed with an underwater camera equipped with a wide-angle (15 mm) lens and an electronic strobe.

A matrix "stations \times species" was produced, using presence / absence data. Similarity between stations was calculated applying Sorensen coefficient (BOUDOURESQUE 1971) and stations




Fig. 1. Location of study area at Puerto del Carmen Lanzarote (Canary Islands), with diving sites indicated.

were subsequently clustered into a dendrogram through complete linkage.

RESULTS

A total of 152 species/species groups was observed and/or photographed in the 4 diving sites: 21 macroalgae, 11 sponges, 32 cnidarians, 10 molluscs, 6 polychaetes, 7 crustaceans, 6 lophophorates, 9 echinoderms, 4 ascidians, and 46 fishes (Table 1).

Species-groups were used in the case of organisms not easily recognised underwater or on the slides, but sharing the general aspect (HISCOCK 1987). The name "algal turf" was given to a mixture of different small-sized algal thalli, as defined by NETO (1992). A similar definition applies to hydroid mat.

Encrusting coralline algae, encrusting




Fig. 2. Schematic sea-bottom profiles at three dive sites: El Agujero, Bajamar, Tiñosa. Horizontal distances are taken from maps and are thus simply indicative. Letters indicate stations.

bryozoans and encrusting sponges were especially abundant in sea-urchin barrens where *Baltzella inops* and *Phorbas fictitius* were the most common sponges, and *Reptadeonella violacea* and *Schizoporella longirostris* the most common bryozoans. However, similar encrusting species occurred also in other habitats, and a consistent identification down to species level was not possible.

Many different species of sponges shared a massive growth-form, but species of *Acanthella*, *Aplysina*, *Axinella*, *Ircinia* and *Spongionella* were excluded from the group and consistently recognised to at least genus level. The group "large hydroids" comprised species of *Aglaophenia*, *Eudendrium* and *Halecium*, among others. Serpulidae and Vermetidae possibly included only one species each, whereas *Ircinia* sp.p. might well correspond to different massive sponges: CRUZ SIMÓ (1984) listed five *Ircinia* species for the Canaries, all occurring in habitats such as those studied here. A similar remark applies for the alga *Hypnea* sp.p., several species being known in the region (HAROUN & PRUD'HOMME VAN REINE 1993).

Cluster analysis applied to the

presence / absence data matrix identified 5 major group of stations (Fig. 5).

The best defined cluster was composed by five stations (Q Y U D K), all located on the sand slope which, starting from the base of the littoral cliff at about 10 m, reached 25 m depth at a distance of about 150-180 m from the shore (Fig. 2). It was a rather steep slope of bare volcanic sand nearly deprived of visible flora and fauna. Thus, this cluster resulted defined primarily by a number of fish species (Table 1), among which the conger eel *Heteroconger longissimus* was noticeable for its characteristic "gardens". Swarms of the opossum shrimp *Paramysis arenosa* were also common (WITTMAN & WIRTZ 1998).

A second well-defined cluster included six stations (E G F Z H V) located on rock in shallow water (Fig. 5). These stations corresponded to the base of the littoral rocky cliff, ending generally within 7 m depth and leaving place to a boulder field (Fig. 4). The rocky substrate generally exhibited high biological cover, with the dominance of photophilic algae, such as *Dictyota dichotoma*, *Pterocladia capillacea* and many others (Table 1). Small areas of bare rock were




Fig. 3. A pictorial view of the submerged volcanic reef off Bajamar, running almost parallel to the coastline of Puerto del Carmen. Letters indicate stations.

nevertheless observed chiefly where sea urchins (especially *Arbacia lixula*) were abundant. The sponge *Aplysina aerophoba* and the sea anemone *Anemonia viridis* were the most important sessile invertebrates: both species are known to harbour phototrophic endosymbionts (BIANCHI et al. 1998). Many different fish species were frequent (Table 1). The boundary between rocks and sand was characterised by crowding *Arachnanthus nocturnus*, a tube anemone locally reaching the density of 200 individuals·m².

The other three clusters were less sharply separated and embraced the stations located on the impressive submerged reef that ran almost parallel to the coastline at the outer end of the sand slope, between 25 and 50 m depth (Fig. 3). The volcanic rock forming the reef was highly fissured and uneven, thus originating many different habitats, among which submarine caves and overhangs (Fig. 2).

Three stations (P J C) located on the reef head and two (X O) located on the (upper) reef wall clustered together (Fig. 5). This cluster was characterised by few algae and several sessile or sedentary animals, among which different echinoderms occurred (Table 1). Rocks at the reef head exhibited low biological cover probably because of the intense grazing by the sea-urchin *Diadema antillarum*; other common organisms were the sea-star *Narcissia canariensis*, the algae

Lobophora variegata, *Lophocladia trichoclados* and *Cottoniella filamentosa*, and the ascidian *Pycnoclavella* sp., together with encrusting sponges and bryozoans.

Another cluster was formed by height stations, five of which (B R N I M) corresponded to caves or overhangs, two (S A) to the reef wall, and only one (T) to attract of reef head near a cave (Fig. 2). A diverse sessile fauna was characteristic of this cluster (sponges, cnidarians, polychaetes etc.), whereas fish were less important (Table 1). Hydroids were abundant on the reef wall. Stations I and M, corresponding to deep overhangs characterised by the tree coral *Dendrophyllia ramea*, formed a slightly distinct subcluster (Fig. 5).

Finally, the last cluster was made up by only two stations (L W), located on the deep rocks at the base of the reef. This cluster was defined especially by its exclusive sessile biota, with the black coral *Anthipathes wollastoni* forming extensive "forests". Different gorgonian species and the epizoid zoanthid *Gerardia savaglia* were also common (Table 1).

DISCUSSION

The five station clusters may easily be interpreted as different benthic habitats, each one having its peculiar biotic community (in the sense of HISCOCK & MITCHELL 1980).

These communities fit well within the general scheme outlined for the Canaries by BACALLADO ARÁNEGA (1984) and GONZÁLES HENRÍQUEZ et al. (1986). Analogies can be found with those described by HERRERA et al. (1993) at Playa del Cabrón, Gran Canaria, and by BALLESTEROS (1993) in some stations off Fuerteventura and Lanzarote.

The main difference between the latter study and the present one, is the relatively limited extension of photophilic algal assemblages at




Fig. 4. Schematic map of the sea-bottom in the dive site called "El Muelle". Letters indicate stations (left). Vertical profile and zonation at station Z (right): a) *Ulva rigida* and *Actinia equina*; b) *Corynactis viridis*; c) *Corallina elongata*; d) *Arbacia lixula* and encrusting corallines; e) *Parazoanthus* sp.; f) cave with encrusting and massive sponges; g) encrusting corallines; h) *Dictyota dichotoma* and other algae.




Fig. 5. Dendrogram resulting from cluster analysis (Sorensen coefficient, complete linkage). Letters indicate stations (see text for explanation).

Puerto del Carmen, but this may be explained by the fact that the shallow water area investigated was rather sciophilous, under steep cliffs.

Anyway, the general dominance by Dictyotales (here especially *Dictyota dichotoma*) is a common feature of such algal assemblages in the whole Atlantic-Mediterranean region (BIANCHI et al. 1998).

At Playa del Cabrón, HERRERA et al. (1993) limited their study to a depth of 30 m, whereas at Puerto del Carmen we extended investigation down to about 50 m. The community we observed in deep water is similar to that found by BALLESTEROS (1993) at 53 m depth off Punta Tiñosa and by ARISTEGUI et al. (1987) when dredging from 40 to 200 m depth off Tenerife. Diving allowed us to allocate *Dendrophyllia ramea* and the other bushy anthozoa (gorgonians and antipatharians) to different "sub-habitats": the former was found under overhangs, the latter on sub-horizontal rocks. Gorgonians characterise a similar depth zone on

hard bottoms also in the Mediterranean Sea, but antipatharians and *Dendrophyllia ramea* live deeper (PÉRES & PICARD 1964). A gorgonian-

Table 1

List of the species observed or photographed during the dives at Puerto del Carmen, February 1996, 0 to 54 m depth, and their presence in the five major habitats recognised through cluster analysis and ranked according to increasing depth. Notes: ¹ confused within "encrusting sponges"; ² on shells occupied by the hermit crab *Pagurus prideaux*; ³ confused within "large hydroids"; ⁴ inside sponges, in very high densities (WIRTZ 1996); ⁵ the rich crustacean fauna associated with this species has been described by WIRTZ (1997); ⁶ associated with the sea-star *Astropecten aranciaceus* and visible at night; ⁷ associated with the black coral *Antipathes wollastoni* (FRANSEN & WIRTZ 1998); ⁸ confused within "encrusting bryozoans".

	Shallow rock (< 10 m)	Sand slope (10-20 m)	Reef head (~ 25 m)	Caves and reef wall (30-40 m)	Deep rocks (~ 50 m)
ALGAE					
algal turf	+	.	+	+	+
<i>Asparagopsis taxiformis</i>	+	.	+	.	.
<i>Carpomitra costata</i>	+
<i>Caulerpa racemosa</i>	+
<i>Caulerpa webbiana</i>	+	.	+	+	+
<i>Cottoniella filamentosa</i>	.	+	+	.	+
<i>Dictyota bartayresii</i>	+
<i>Dictyota dichotoma</i>	+	.	+	+	.
encrusting corallines	+	.	+	+	+
<i>Falkenbergia rufolanosa</i>	+	+	+	.	+
<i>Halopteris filicina</i>	+	.	+	.	+
<i>Hypnea</i> sp. p.	+	+	+	.	.
<i>Lobophora variegata</i>	+	+	+	.	.
<i>Lophocladia trichoclados</i>	+	+	+	.	+
<i>Microdictyon tenuis</i>	+
<i>Padina pavonica</i>	+	+	+	.	.
<i>Palmophyllum crassum</i>	+	.	+	+	+
<i>Pterocladia capillacea</i>	+
<i>Stypocaulon scoparium</i>	+	.	+	.	.
<i>Stypopodium zonale</i>	+	+	+	.	.
<i>Ulva rigida</i>	+
PORIFERA					
<i>Acanthella acuta</i>	.	.	.	+	.
<i>Aplysina aerophoba</i>	+	.	+	.	.
<i>Axinella damicornis</i>	.	.	+	+	+
<i>Axinella polypoides</i>	.	.	.	+	.
<i>Baltzella inops</i> ¹	+	.	+	.	.
encrusting sponges	+	.	+	+	+
<i>Ircinia</i> sp. p.	+	.	.	+	.
massive sponges	+	.	+	+	+
<i>Phorbast fictitius</i> ¹	+	.	+	.	.
<i>Plakortis simplex</i>	.	.	+	+	.
<i>Spongionella pulchella</i>	+	.	+	+	.
CNIDARIA					
<i>Actinia equina</i>	+
<i>Adamsia palliata</i> ²	.	+	.	.	.
<i>Aglaophenia pluma</i> ³	.	.	+	.	.
<i>Aiptasia mutabilis</i>	+
<i>Alicia mirabilis</i>	+
<i>Anemonia melanaster</i>	+

(Table 1 continued)

<i>Anemonia viridis</i>	+
<i>Antipathes wollastoni</i>	.	.	.	+	+
<i>Arachnanthus nocturnus</i>	+	+	.	.	.
<i>Balanophyllia regia</i>	+	.	+	.	.
<i>Caryophyllia inornata</i>	.	.	+	+	.
<i>Corynactis viridis</i>	+
<i>Dendrophyllia ramea</i>	.	.	.	+	.
<i>Ellisella paraplexauroides</i>	+
<i>Eudendrium</i> sp. ³	.	.	.	+	.
<i>Eunicella verrucosa</i>	+
<i>Gerardia savaglia</i>	+
<i>Halecium</i> sp. ³	.	.	.	+	.
<i>Hoplangia durotrix</i>	.	.	.	+	.
hydroid mat	+	.	.	+	.
large hydroids	.	.	+	+	+
<i>Leptopsammia pruvoti</i>	.	.	+	+	+
<i>Lophogorgia ruberrima</i>	+
<i>Lophogorgia viminalis</i>	+
<i>Madracis asperula</i>	.	.	.	+	.
<i>Madracis pharensis</i>	.	.	.	+	.
<i>Nausithoe punctata</i> ⁴	.	.	.	+	.
<i>Pachycerianthus</i> sp.	.	.	+	+	.
<i>Paracyathus pulchellus</i>	.	.	+	.	.
<i>Parazoanthus</i> sp.	+	.	.	+	+
<i>Phyllangia mouchezii</i>	.	.	+	+	.
<i>Telmatactis cricoides</i> ⁵	+	.	+	.	.
MOLLUSCA					
<i>Chama gryphoides</i>	.	.	+	.	.
<i>Coryphella pedata</i>	.	.	+	.	.
<i>Discodoris atromaculata</i>	+
<i>Hypselodoris fontandraui</i>	+
<i>Hypselodoris picta</i>	.	.	.	+	.
<i>Neopycnodonte cochlear</i>	.	.	.	+	.
<i>Pinna rudis</i>	+
<i>Spurilla neapolitana</i>	+
<i>Umbraculum umbraculum</i>	+
Vermetidae gen. sp.	.	.	.	+	.
POLYCHAETA					
<i>Acholoe squamosa</i> ⁶	.	+	.	.	.
<i>Bispira viola</i>	.	.	.	+	.
<i>Hermodice carunculata</i>	.	.	.	+	+
<i>Protula tubularia</i>	.	.	+	+	.
Serpulidae gen. sp.	.	.	+	+	.
<i>Vermiliopsis</i> sp.	.	.	.	+	.
CRUSTACEA					
<i>Balanus trigonus</i>	.	.	+	+	.
<i>Eualus occultus</i>	.	+	.	.	.
<i>Pagurus prideaux</i>	.	+	.	.	.
<i>Paramysis arenosa</i>	.	+	.	.	.
<i>Percnon gibbesi</i>	+
<i>Periclimenes wirtzi</i> ⁷	+
<i>Stenorhynchus lanceolatus</i>	+	.	+	+	.

(Table 1 continued)

LOPHOPHORATA					
<i>Bugula plumosa</i>	.	.	.	+	.
encrusting bryozoans	+	.	+	+	+
<i>Phoronis hippocrepia</i>	.	.	.	+	.
<i>Reptadeonella violacea</i> ⁸					
<i>Schizoporella longirostris</i> ⁸					
<i>Smittina cervicornis</i>	.	.	.	+	.
ECHINODERMATA					
<i>Arbacia lixula</i>	+
<i>Astropecten aranciatus</i>	.	+	.	.	.
<i>Diadema antillarum</i>	+	.	+	+	.
<i>Hacelia attenuata</i>	+
<i>Holothuria</i> cf. <i>forskali</i>	+
<i>Holothuria</i> sp.	+	+	+	.	.
<i>Martasterias glacialis</i>	.	.	+	.	.
<i>Narcissa canariensis</i>	.	.	+	+	.
<i>Sphaerechinus granularis</i>	+	.	+	.	.
TUNICATA					
<i>Ascidia mentula</i>	+
<i>Clavelina lepadiformis</i>	.	.	.	+	.
<i>Didemnum</i> sp.	+	.	.	+	+
<i>Pycnoclavella</i> sp.	.	.	+	+	+
PISCES					
<i>Abudefduf luridus</i>	+	.	+	+	.
<i>Anthias anthias</i>	.	.	+	.	+
<i>Apogon imberbis</i>	+	.	+	+	.
<i>Bodianus scrofa</i>	.	.	.	+	.
<i>Boops boops</i>	+	+	.	.	.
<i>Bothus podas madeirensis</i>	.	+	.	.	.
<i>Canthigaster rostrata</i>	+	+	+	+	.
<i>Centrolabrus trutta</i>	+
<i>Chelon labrosus</i>	+	+	.	.	.
<i>Chromis limbata</i>	+	.	+	.	+
<i>Diplodus cervinus</i>	+
<i>Diplodus sargus cadenati</i>	+	.	+	+	.
<i>Diplodus vulgaris</i>	+
<i>Epinephelus marginatus</i>	+	.	+	+	.
<i>Gobius niger</i>	.	+	.	.	.
<i>Heteroconger longissimus</i>	.	+	.	.	.
<i>Lithognathus mormyrus</i>	+	+	.	.	.
<i>Liza aurata</i>	+	+	.	.	.
<i>Mullus surmuletus</i>	+	+	.	.	.
<i>Muraena augusti</i>	.	.	+	.	.
<i>Mycteroperca fusca</i>	+
<i>Oblada melanura</i>	+
<i>Ophioblennius atlanticus</i>	+
<i>Pagellus erythrinus</i>	.	+	.	.	.
<i>Parablennius parvicornis</i>	+
<i>Parablennius pilicornis</i>	+
<i>Pseudocaranx dentex</i> juv.	.	+	.	.	.
<i>Sardinella maderensis</i>	+	+	.	.	.
<i>Sarpa salpa</i>	+
<i>Scartella cristata</i>	+
<i>Scorpaena maderensis</i>	+	.	+	+	.

(Table 1 continued)

<i>Serranus atricauda</i>	.	.	.	+	.
<i>Sparisoma cretense</i>	+	.	+	+	.
<i>Sphoeroides marmoratus</i>	.	.	+	.	.
<i>Sphyræna viridensis</i>	.	+	+	+	.
<i>Spondyllosoma cantharus</i>	+
<i>Squatina squatina</i>	.	+	.	.	.
<i>Stephanolepis hispidus</i>	+
<i>Synodus saurus</i>	+	+	.	.	.
<i>Synodus synodus</i>	+
<i>Taeniura grabata</i>	+
<i>Thalassoma pavo</i>	+	.	+	+	+
<i>Trachinotus ovatus</i>	+
<i>Trachinus draco</i>	.	+	.	.	.
<i>Uranoscopus scaber</i>	.	+	.	.	.
<i>Xyrichtys novacula</i>	.	+	.	.	.

antipatharian community similar to that observed at Puerto del Carmen, although composed by different species, was described from the Archipelago of Cape Verde by MORRI & BIANCHI (1995).

The discussion above suggests that an economic and "low-tech" approach, mostly based on visual census and photography, may be sufficient for the description of main epibenthic communities. Recording 150 species in a limited area with such a low "sampling" effort stands for the high marine biodiversity of the Canaries.

Clearly, a major problem is that not everything can be recognised underwater or on the photographs to the species level. This was especially true for encrusting organisms, which might well be represented by different species in the different habitats we studied, and could in some cases have dampened habitat discrimination (for example, poor sorting of reef-head and reef-wall stations in cluster analysis).

Another limit is that small species were not easily seen underwater, but might be characteristic too. In some cases, these small species lived strictly associated with other, larger organisms (see notes in Table 1). Being prevalently linked to their host rather than to the physical environment, they are species-specific not really biotope-specific (BIANCHI et al. 1989; MORRI et al. 1991). Large organisms, which monopolise or dominate the substratum on which they live, are better "indicators" (KÖNNECKER 1977) of the

environmental conditions or factors to which they are adapted and should be preferred for the description and classification of epibenthic communities.

ACKNOWLEDGEMENTS

The "Centro de Buceo Atlantica" and "Safari Diving" provided logistic support to our dives at Puerto del Carmen. K. Wittman (Wien) participated in some of the dives. J. Núñez (Tenerife), M. Pansini (Genova) and J. C. Miquel (Monaco) provided references. J. C. den Hartog (Leiden), G. Jarms (Hamburg) and P. Knight-Jones (Swansea) kindly identified *Anemonia melanaster*, *Nausithoe punctata* and *Bispira viola*, respectively. This work was done within the frame of the Research Project "MACMED" (comparing marine benthic communities and zonation in MACaronesia and the MEDiterranean) of the Marine Environment Research Centre of La Spezia.

REFERENCES

- ARÍSTEGUI, J., A. BRITO, T. CRUZ, J. J. BACALLADO, J. BARQUÍN, J. NÚÑEZ & G. PÉREZ-DIONIS 1987. El poblamiento de los fondos de *Dendrophyllia ramea* (Anthozoa, Scleractinia) en las Islas Canarias. *Cuadernos marisqueros: Publicaciones técnicas* 11: 163-181.

- AUGIER, H. 1982. *Inventory and classification of marine benthic biocenoses of the Mediterranean*. Council of Europe, Strasbourg (Nature and environment series No. 25). 57 pp.
- BACALLADO ARÁNEGA, J. J. (Ed.) 1984. *Fauna marina y terrestre del Archipiélago Canario*. Edirca, Las Palmas de Gran Canaria. 356 pp.
- BALLESTEROS, E. 1993. Algunas observaciones sobre las comunidades de algas profundas en Lanzarote and Fuerteventura (Islas Canarias). *Vieraea* 22: 17-27.
- BELLAN-SANTINI, D. J. C. LACAZE & C. POIZAT (Eds) 1994. *Les biocénoses marines et littorales de Méditerranée: synthèse, menaces et perspectives*. Secrétariat de la faune et de la flore, Muséum National d'Histoire Naturelle, Paris. 246 pp.
- BEYHL, F. E., B. MIES & P. OHM 1995. Macaronesia: a biogeographical puzzle. *Boletim do Museu municipal do Funchal (História Natural)*, Supplement 4 A: 107-113.
- BIANCHI, C. N., D. BEDULLI, C. MORRI & A. OCCHIPINTI AMBROGI 1989. L'herbier de Posidonies: écosystème ou carrefour éco-éthologique? Pp. 257-272 in: BOUDOURESQUE, C. F., A. MEINESZ, E. FRESI & V. GRAVEZ (Eds). *2nd International Workshop on Posidonia Beds*. GIS Posidonie, Marseille. 321 pp.
- BIANCHI, C. N., S. COCITO, C. MORRI & S. SGORBINI 1991. Rilevamento bionomico subacqueo. Pp. 67-83 in: ABBIATI, M. (Ed.) *Lezioni del corso formativo per ricercatore scientifico subacqueo*. International School for Scientific Diving, Pisa. 240 pp.
- BIANCHI, C. N., C. MORRI, G. F. SARTONI & P. WIRTZ 1998. Sublittoral epibenthic communities around Funchal (Ilha da Madeira, NE Atlantic). *Boletim do Museu Municipal do Funchal (História Natural)* Suplemento 5: 51-80
- BRITO, A 1991. Catálogo de los peces de las islas Canarias. F. Lemus, La Laguna, 230 pp.
- BOUDOURESQUE, C. F. 1971. Méthodes d'étude qualitative et quantitative du benthos (en particulier du phytobenthos). *Tethys* 3(1): 79-104.
- COMMISSION OF THE EUROPEAN COMMUNITIES 1991. *CORINE biotopes*. Office for Official Publications of the European Communities, Luxembourg (1st edition). 300 pp.
- CONNOR, D. W., K. HISCOCK, R. L. FOSTER-SMITH & R. COVEY 1995. A classification system for benthic marine biotopes. Pp. 155-159 in: ELEFThERIOU, A., A. D. ANSELL & C. J. SMITH (Eds). *Biology and ecology of shallow coastal waters*. Olsen & Olsen, Fredensborg. 391 pp.
- CRUZ SIMÓ, T. & J. J. BACALLADO ARÁNEGA 1984. Introducción a los poblamientos de espongiarios de las Islas Canarias. *Actas do IV simpósio ibérico de estudos do benthos marinho*. Lisboa, 3: 141-150.
- FRANSEN, C. H. J. M. & P. WIRTZ 1998. Contribution to the knowledge of decapod crustaceans from Madeira and the Canary Islands. *Zoologische Mededelingen* 71 (19): 215-230.
- GLÉMAREC, M. 1973. The benthic communities of the European north Atlantic shelf. *Oceanography and marine Biology: an annual Review* 11: 263-289.
- GONZÁLES HENRÍQUEZ, M. N., J. D. RODRIGO PÉREZ & C. SUÁREZ RODRÍGUEZ 1986. *Flora y vegetación del Archipiélago Canario*. Edirca, Las Palmas de Gran Canaria. 335 pp.
- HAROUN, R. J. & W. PRUD'HOMME VAN REINE 1993. A biogeographical study of *Laurencia* and *Hypnea* species of the Macaronesian region. *Courier Forschungsinstitut Senckenberg* 159: 119-125.
- HAROUN TABRAUE, R. J., M. C. GIL-RODRÍGUEZ, J. AFONSO CARRILLO & W. WILDPRET DE LA TORRE 1984. Vegetación bentónica del Roque de los Organos (Gomera). *Anales de Biología* 2(sección especial, 2): 107-117.
- HERRERA, R., D. MONTERO & R. HAROUN 1993. Bionomía bentónica del litoral de la playa del Cabrón (Gran Canaria). *Publicaciones especiales del Instituto español de Oceanografía* 11: 291-298.
- HISCOCK, K. 1987. Subtidal rock and shallow sediments using diving. Pp. 198-237 in: BAKER, J. M. & W. J. WOLFF (Eds). *Biological surveys of estuaries and coasts*. Cambridge University Press, Cambridge. 449 pp.
- HISCOCK, K. & R. MITCHELL 1980. The description and classification of sublittoral epibenthic ecosystems. Pp. 323-370 in: PRICE, J. H., D. E. G. IRVINE & W. F. FARNHAM (Eds) *The shore environment* (Volume 2: Ecosystems). Academic Press, London. 945 pp.
- KÖNNECKER, G. 1977. Epibenthic assemblages as indicators of environmental conditions. Pp. 391-395 in: KEEGAN, B. F., P. O'CEIDIGH & P. J. S. BOADEN (Eds) *Biology of benthic organisms*. Pergamon Press, Oxford. 630 pp.
- LAWSON, G. W. & T. A. NORTON 1971. Some observations on littoral and sublittoral zonation at Tenerife (Canary Isles). *Botanica marina* 14: 116-120.
- MEDINA, M., R. J. HAROUN & W. WILDPRET 1995. Phytosociological study of the *Cystoseira abies-marina* (Gmelin) C. Agardh (Cystoseiraceae, Phaeophyceae) community in the Canarian Archipelago. *Boletim do Museu Municipal do Funchal (História Natural)* Suppl. 4 B: 433-440.

- MORRI, C. & C. N. BIANCHI 1995. Cnidarian zonation at Ilha do Sal (Arquipélago de Cabo Verde). *Beiträge zur Paläontologie* 20: 41-49.
- MORRI, C., G. BAVESTRELLO & C. N. BIANCHI 1991. Faunal and ecological notes on some benthic cnidarian species from the Tuscan Archipelago and eastern Ligurian Sea (western Mediterranean). *Bollettino dei Musei e degli Istituti biologici dell'Università di Genova* 54-55 (1988-1989): 27-47.
- NETO, A. I. 1992. Contribution to the taxonomy and ecology of the Azorean benthic marine algae. *Biological Journal of the Linnean Society* 46: 163-176.
- PÉRÈS, J. M. 1982. Major benthic assemblages. Pp. 372-508 in: KINNE, O. (Ed.) *Marine Ecology*, Volume V, *Ocean Management*. Part I. J. Wiley & Sons, Chichester. 642 pp.
- PÉRÈS, J. M. & J. PICARD 1964. Nouveau manuel de bionomie benthique de la Mer Méditerranée. *Recueil des Travaux de la Station marine d'Endoume* 31 (=47): 1-137.
- PÉREZ SANCHEZ, J. M. & MORENO BATET, E. 1991. Invertebrados marinos de Canarias. Ediciones del Cabildo Insular de Gran Canaria, Las Palmas, 335 pp.
- TORTONESE, E. 1960. The relations between the Mediterranean and Atlantic fauna. *Hidrobiologi* ser. B, 5 (1-2): 30-34.
- WIRTZ, P. 1994. *Unterwasserführer Madeira, Kanaren, Azoren: Fische / Underwater guide Madeira, Canary Islands, Azores: Fish*. Verlag Stephanie Naglschmid, Stuttgart. 159 pp.
- WIRTZ, P. 1995. *Unterwasserführer Madeira, Kanaren, Azoren: Niedere Tiere / Underwater guide Madeira, Canary Islands, Azores: Invertebrates*. Verlag Stephanie Naglschmid, Stuttgart. 247 pp.
- WIRTZ, P. 1996. Coronatenpolypen in Horschwämmen. *Das Aquarium* 329: 29-31.
- WIRTZ, P. 1997. Crustacean symbionts of the sea anemone *Telmatactis cricoides* at Madeira and Canary Islands. *Journal of Zoology* 242: 799-811.
- WITTMAN, K. J. & P. WIRTZ (in press). A first inventory of the mysid fauna (Crustacea: Mysidacea) in coastal waters of the Madeira and Canary Archipelagos. *Boletim do Museu municipal do Funchal (História Natural)*.

Accepted 1 February 1999.