

First evaluation of neustonic microplastics in the Macaronesian region, NE Atlantic

A. Herrera^{a,*}, E. Raymond^a, I. Martínez^a, S. Álvarez^b, J. Canning-Clode^{b,d,e}, I. Gestoso^{b,d}, C.K. Pham^c, N. Ríos^c, Y. Rodríguez^c, M. Gómez^a

^a*Marine Ecophysiology Group (EOMAR), IU- ECOAQUA. Universidad de Las Palmas de Gran Canaria, Canary Islands, Spain.*

^b*MARE-Marine and Environmental Sciences Centre, Agência Regional para o Desenvolvimento da Investigação Tecnologia e Inovação (ARDITI), Funchal, Madeira, Portugal.*

^c*IMAR/OKEANOS - Universidade dos Açores, Departamento de Oceanografia e Pescas, 9901-862 Horta, Portugal*

^d*Centre of IMAR of the University of the Azores, Department of Oceanography and Fisheries, Azores, Portugal.*

^e*Smithsonian Environmental Research Center, 647 Contees Wharf Road, Edgewater, MD 21037, U.S.A*

Abstract

Marine microplastic pollution is an issue of great concern nowadays since high concentrations have been detected in the ocean, mainly in the subtropical gyres that accumulate this type of debris. The long-term effects of this pollution on ecosystems and marine biota are still unknown. The aim of this study is to quantify and characterise microplastics and neustonic zooplankton in sub-surface waters of the Macaronesian region, an area that has been little studied to date. Our results show a great variability in the concentration of microplastics with values between 15,283 items/Km² in Los Gigantes (Tenerife, Canary Islands) and 1,007,872 items/Km² in Las Canteras (Gran

*Corresponding author. Tel.: +34 928 45 44 40; fax: +34 928 45 29 22
Email address: alicia.herrera@ulpgc.es (A. Herrera)

Canaria, Canary Islands). The main types of debris found were plastic fragments and fibres. The abundances of neustonic zooplankton were also very variable between the different sampling areas, being the main components copepods and eggs. Regarding the microplastics-zooplankton ratio, values were obtained between 0.002 and 0.22. In Las Canteras, the highest accumulation zone, was found twice as much microplastics as zooplankton for the 1-5mm fraction in dry weight. These values highlight the potential hazard of microplastics - and its associated chemical contaminants - for marine biota, especially for large filter feeders.

Keywords: marine debris, microdebris, plastic, zooplankton, manta net, marine litter, North Atlantic

1. Introduction

Large-scale plastic production has continued to grow from its beginnings in 1950 to the present day, reaching almost 350 million tonnes in 2017 (PlasticsEurope, 2018). The accumulation of this plastic waste and its entry into the ocean, estimated at 4.8 to 12.7 million metric tons per year (Jambeck et al., 2015), is one of the major environmental problems of the present time.

The United Nations Environmental Programme (UNEP) defines marine litter as ‘any persistent, manufactured or processed solid discarded material, disposed of or abandoned in the marine and coastal environment’ (UNEP, 2009), the great majority of these wastes are plastics, but glass, wood, tar, metal, natural fibres, etc. can also be found (Kroon et al., 2018). Nowadays there is no consensus on the size that defines microplastics and microdebris,

14 in 2009 NOAA proposes a definition in which microplastics are considered as
15 all plastic particles with $<5\text{mm}$ in diameter (Arthur et al., 2009), EU MSFD
16 WG-GES (MSDF Technical Subgroup on Marine Litter, 2013) proposes >20
17 and $<5000\mu\text{m}$, and recently Hartmann et al. (2019) propose the size between
18 >1 and $<1000\mu\text{m}$ to define microplastics. In the present study we use the
19 term microdebris to describe particles of anthropogenic origin with a size less
20 than 5mm .

21
22 Pollution by microplastics is an issue of growing concern in the scientific
23 community, environmental policy authorities and society due to the potential
24 risk it may have for ecosystems, marine biota, human health and the econ-
25 omy. The topic is being widely studied at a global level, on the other hand,
26 in the Macaronesian region although results are known in beaches (Baztan
27 et al., 2014; Herrera et al., 2018; Álvarez-Hernández et al., 2019; Gestoso
28 et al., 2019; Chambault et al., 2018; Ríos et al., 2018), and there are some
29 studies on marine biota (Rodríguez et al., 2012; Rodríguez and Pham, 2017;
30 Pham et al., 2014; Herrera et al., 2019b), microplastic contamination in sub-
31 surface waters has been little studied.

32
33 The Macaronesia region is conformed by a group of islands located in
34 the Eastern Atlantic, which form a biogeographic region. It includes more
35 than 40 islands grouped into five archipelagos: Azores, Madeira, Selvagens,
36 Canary Islands and Cape Verde. In total they occupy an area of more than
37 $14,600\text{ Km}^2$. The Macaronesian region has great biodiversity and many en-
38 demic species, 211 Sites of Community Importance (SCIs) and more than 65

39 Special Protection Areas (SPAs) have been declared (Sundseth, 2010). Due
40 to their oceanographic situation close to the North Atlantic subtropical gyre
41 (NASG), the islands are located at the flow of the Azores Current and the Ca-
42 nary Current, branches downstream of the Gulf Stream (Comas-Rodríguez,
43 2011). As a result, such oceanic islands are predicted to be particularly vul-
44 nerable to plastic pollution.

45
46 One approach of assess the potential risk of microplastics to marine or-
47 ganisms, particularly filter feeders, is to study the ratio between the amount
48 of neustonic microplastics and zooplankton Moore et al. (2001). This ratio
49 increases in areas of the ocean with low productivity where the number of or-
50 ganisms decreases and the amount of plastic accumulates, such as in oceanic
51 gyres.

52
53 For the above mentioned reasons, the main objectives of this study are
54 to determine the abundance and characterize the floating microplastics in
55 different zones of the Macaronesian region, and to study the microplastics-
56 zooplankton ratio.

57 58 **2. Materials and Methods**

59 *2.1. Samples collection*

60 A total of 45 neustonic samples were collected during daylight (9:00-14:00
61 hs.), 24 in the Canary Islands archipelago, 12 in Madeira and 9 in the Azores
62 in the Macaronesian region (Fig. 1). Samples were collected in opportunistic

63 samplings in different periods between 2015 and 2018. The collection dates
64 and locations of each sample are shown in table 1 of supplementary material.
65 In the Canary Islands and Madeira the neustonic samples were collected with
66 a manta net with a rectangular mouth opening of 25 x 60 cm, and a 200 μ m
67 mesh size. At each location 3 samples were taken, except in Taliarte where
68 only 2 samples were collected and in Los Gigantes where 4 samples were
69 collected. The manta net was trawled at a speed of ~3 knots, during periods
70 of 20 minutes. GPS coordinates were taken to measure the length of each
71 transect. The net trawls were towed at a horizontal angle of 45° with respect
72 to the ship's trails. Kukulka et al. (2012) demonstrated that in strong wind
73 conditions the neuston net tends to collect less plastic particles because it
74 is distributed vertically in the mixed layer due to wind-induced mixing. For
75 this reason the sampling was only carried out under optimal sea conditions,
76 with a Beaufort Sea Scale between 0 and 2.

77 In the Azores, three parallel transects were carried out at each site using
78 200 μ m mesh bongo nets 50 cm in diameter. Each tow lasted 2min20 sec-
79 onds at a constant speed of ~2 knots. The volume of water filtered in the
80 tows was calculated using a flowmeter only in Azores archipelago. The start
81 and end coordinates were also recorded to determine the length of each tran-
82 sect.

83 Samples were collected and preserved in 4% of formaldehyde for later anal-
84 ysis.

85 *2.2. Samples processing*

86 All debris range from 0.2-5mm were identified and counting by visual
87 inspection under a binocular stereomicroscope (Leica S9i) with integrated

CMOS camera, at 55x magnification. Microdebris were classified in different categories regarding its typology; irregular plastic fragments (Fragments), industrial pellets (Pellets), fibres (Fibres), films and sheets (Films), plastic microbeads (Microbeads), EPS and XPS (Foams), fishing lines (Lines) and others debris including glass, paint, aluminium foil and tar (Other) (Fig. 2). Since FTIR was not available to identify the type of particles in the category fibres are included synthetic, semi-synthetic and natural. During the entire process in the laboratory cotton lab coats were worn and all materials were carefully rinsed with bidistilled water to prevent contamination of the samples. A petri dish with clean 50 μ m mesh was placed near the stereomicroscope during the visual inspection as contamination control. No contamination was found in any of the controls.

Zooplankton neuston samples were separated in 200-500; 500-1000 and >1000 μ m fraction size. Then, an aliquot of 10 or 20ml, depending on plankton concentration, were scanned in a high resolution scan (Epson V800 Pro) and were counted and classified in large taxonomic groups using Zooprocess software V7.30 and ECOTAXA V2.0 (Picheral et al., 2017), as described in the protocol by Herrera et al. (2019a)

The microplastics and zooplankton (in number of units) collected were divided by the total area of filtered water and the concentration was expressed in items/Km². The concentration was expressed in items/m³ only for the Azores' samples.

113 2.3. Statistical analysis

114 The data were analyzed and plotted using R statistical program V3.5.3
115 (R Core Team, 2019). To confirm normality, microplastics and zooplank-
116 ton abundance data were analyzed by the Shapiro Wilk test and the ho-
117 moscedasticity of the residuals was assessed graphically. Microplastics and
118 zooplankton data were not normal and statistical differences between areas
119 were tested using Kruskal-Wallis test and Conover posthoc test.

120 3. Results

121 3.1. Microplastics and zooplankton abundance

122 The maximum values of microplastics (items/Km²) were 1,007,872 in the
123 Canary Islands, 467,259 in the Azores and 124,190 in Madeira (Table 2).
124 However, no significant differences were found between the abundances of mi-
125 croplastics (items/Km²) among the archipelagos (p-value=0.35). The mean
126 values found at each locality expressed per Km² are summarized in table 1.
127 If we consider the values obtained in each locality, the maximum abundance
128 found was in Las Canteras (1,007,872 items/Km²) and the minimum in Los
129 Gigantes (15,482 items/Km²) both in the Canary archipelago (Figs. 3a and
130 4b).

131 Regarding the differences between localities within each archipelago, differ-
132 ences were only observed in the Canary Islands archipelago, being the values
133 in Las Canteras significantly higher than in San Andres, Los Gigantes, and
134 Famara as shown in figure 3a.

135

136 The maximum zooplankton abundance found were 288.9×10^6 ind/Km² in
137 Porto Pim, Azores; 73.4×10^6 ind/Km² in Famara, Canary Islands; and 24×10^6
138 ind/Km² in Funchal, Madeira. Significant differences in zooplankton abun-
139 dance were observed between the Azores and Madeira (p-value= 7.4×10^{-7})
140 and between the Azores and the Canary Islands (p-value= 3.3×10^{-6}). The
141 mean abundances found in each locality expressed per Km² are summa-
142 rized in table 1. Within the Canary Island archipelago, in Famara, there
143 was significantly higher abundance of zooplankton than in Taliarte and San
144 Andres (p-values<0.05). However, within the archipelagos of Madeira and
145 Azores no significant differences were observed between the locations (p-
146 values>0.05)(Fig. 3b).

147

148 3.2. *Composition of debris and zooplankton*

149 In the samples collected in the Canary Islands and Azores archipelagos,
150 100% of the debris were microplastics and fibres (synthetic, semi-synthetic
151 or natural). In the Madeira archipelago, on the other hand, 16% were other
152 types of debris. Of the total microplastics collected in the Canary Islands
153 57.3% were fragments, 27.4% fibres, 9.9% lines and 5.3% films; in the Azores
154 archipelago 54% were fibres, 34.9% fragments, 6.3% lines and 4.8% films;
155 while in Madeira, from total debris 47.5% were fragments, 30% fibers, 4.6%
156 styrofoam, 0.5% films, and 16.8% were other debris such as glass, paint, alu-
157 minium foil and tar (Fig. 5).

158 Regarding particle size, in the Canary Islands 50.6% were between 0.2-1mm
159 and the rest between 1-5mm in size; in Madeira 39.4% of the particles had a
160 size between 1-5mm and the rest in the fraction between 0.2-1mm; while in

161 the Azores, only 17.5% of the particles were of the fraction of 1-5mm, being
162 82.5% of a size between 0.2-1mm (Fig. 5).

163 Neustonic zooplankton were classified into large taxonomic groups, in terms
164 of abundance copepods were the dominant group, and the other major group
165 were fish eggs. In the Canary Islands the percentage of each group was 85%
166 copepods, 12.5% eggs, 1% appendicularia, 0.5% salpidae and within the re-
167 maining 1% were found amphipods, annelids, chaetognats, decapods and
168 euphausiids. Also in the Azores, the most abundant group were copepods
169 with 44%, eggs 29.5%, cirripedia larvae 17.2% and ostracods 9.4% (Fig. 6b).
170 In contrast, the neustonic zooplankton collected in Madeira were 60% eggs ,
171 38.1% copepods, 1.2% appendicularia, and the remaining 1% were composed
172 by annelids, decapods, salpids and chaetognats (Fig. 6c).

173

174 3.3. *Ratio Microplastics/Zooplankton*

175 The average ratio of Microplastics/Zooplankton (Micro/Zoo) in each of
176 the archipelagos was 0.032 in the Canary Islands, 0.021 in Madeira and 0.002
177 in the Azores (Table 2). The mean values obtained in each locality are shown
178 in table 1. The highest Micro/Zoo ratios were found in the Canary Islands, in
179 the localities of Taliarte (0.22), Las Canteras (0.1) and Lambra (0.06) (Fig.
180 3c). In Madeira maximum values of 0.06 were found in Caniçal (Fig. 3c). In
181 the Azores archipelago the maximum values reached were 0.005 in Porto Pim
182 and Almoxarife (Fig. 3c). The Micro/Zoo ratio was significantly lower in
183 the Azores than in the Canary Islands and Madeira (p-values<0.05). Within
184 each archipelagos, significant differences were only observed in the Canary
185 Islands, with significantly higher MP/Zoo ratios in Taliarte and Las Canteras

186 than in Famara and Los Gigantes (p-values<0.05)(Fig. 3c).
187 The ratio of microplastics/zooplankton in dry weight was estimated only in
188 the samples from Las Canteras for the fraction >1mm, as they were the only
189 ones that contained enough microplastics to do that estimation. In the 3
190 samples collected within that fraction the Micro/Zoo dry weight ratio was
191 2.70, 2.67 and 0.50 respectively, being the average value 2.0 ± 1.3 .

192

193 **4. Discussion**

194 The mean values of MPs in items per Km² are in the range of those
195 found in other areas of the ocean (see review in table 2). The maximum
196 values found in Las Canteras are similar to those reported in areas of high
197 accumulations such as the North Pacific Central Gyre (Moore et al., 2001)
198 and the Mediterranean Sea (Collignon et al., 2012; Ruiz-Orejón et al., 2016);
199 but lower than those reported by Law et al. (2014) in the Eastern Pacific
200 accumulation zone, and Suaria et al. (2016) and Van Der Hal et al. (2017)
201 also for the Mediterranean.

202

203 High concentrations of microplastics were found in the three archipelagos,
204 especially in the localities of Las Canteras in the Canary Islands, and Porto
205 Pim in the Azores, both located in a semi-enclosed bays, acting as retention
206 zones. Other authors have also reported high abundances of microplastics in
207 bays of Tokyo and Brazil (Cheung et al., 2018; Figueiredo et al., 2018).

208

209 Las Canteras' sampling area is located within El Confital bay on the

210 northeast coast of Gran Canaria. According to the circulation model pro-
211 posed in the study carried out by Mcknight (2016) in El Confital bay, N and
212 NNW wind scenarios shows a recirculation pattern in the eastern-central of
213 the bay. In contrast, with the NE and NNE winds -the predominant winds
214 in the area- it shows a circulation pattern towards the west but intensified in
215 the northeast cape, where the flux is directed towards the bay in southwest
216 direction.

217 Mcknight (2016) analysed the relationship between near-shore surface cir-
218 culation and marine debris deposition on the beach, but there are no studies
219 in the region that relate surface circulation to floating debris. It is probable
220 that the recirculation pattern observed under N and NNW wind conditions
221 may affect the transport of floating debris, determining the accumulation in
222 the central areas, which would explain the high values found at Las Canteras.
223 Further studies are needed to understand the effect of coastal hydrodynamics
224 on the accumulation of neustonic microplastics.

225

226 As can be observed in table 1, there is a great variability in the concen-
227 trations found in the different sampling areas, even between nearby localities
228 such as Las Canteras and San Andres. Although significant differences were
229 observed between archipelagos, both in microplastics and zooplankton abun-
230 dance, it is probable that these differences are due to the fact that sampling
231 was opportunistic, at different times and with different methods. Other au-
232 thors have found that there are variability in the estimations of microplastics
233 according to the methodology used (Barrows et al., 2017; Eriksen et al., 2018;
234 Green et al., 2018) so we should be cautious when making this type of com-

235 parison. Therefore, it is necessary to carry out a specific study in the area
236 in order to determine the spatial variability.

237

238 Our data show for the first time that this region is an area highly pol-
239 luted by microplastics and other debris. The situation of the islands in
240 the flow of the descending branches of the Gulf Stream is probably mak-
241 ing them especially vulnerable to microplastic pollution, as demonstrated by
242 studies on beaches in the region (Baztan et al., 2014; Herrera et al., 2018;
243 Álvarez-Hernández et al., 2019; Gestoso et al., 2019) and marine organisms
244 (Rodríguez et al., 2012; Herrera et al., 2019b; Pham et al., 2017).

245

246 Regarding the categories, most of the microplastics were fragments and
247 fibres, these results agree with those reported worldwide (Aliabad et al.,
248 2019; Cózar et al., 2015; Di Mauro et al., 2017; Eriksen et al., 2013; Faure
249 et al., 2015; Figueiredo et al., 2018; Gewert et al., 2017; Suaria et al., 2016),
250 and with the types of microplastics found in the stomach of Atlantic chub
251 mackerel (*Scomber colias*) collected in Canary Islands waters (Herrera et al.,
252 2019b) and juvenile loggerhead turtles (Pham et al., 2017). However, the
253 percentages found in sub-surface waters off the beaches of Famara, Lambra
254 and Las Canteras do not correspond to those found in high tide line sedi-
255 ments. In Famara almost 44.3% of the microplastic samples collected from
256 beaches were composed of pellets, however in the sub-surface water samples
257 no pellets were found. Something similar occurs in Lambra that presented a
258 35.6% of tar in the sand samples, but this type of waste did not appear in the
259 samples collected with the manta net. These results suggest that the pattern

260 of accumulation of different types of microplastics at the tide line differs from
261 that at the sea surface. This could be due to the fact that the different types,
262 either by their shape or composition, are deposited in different ways in the
263 sand.

264

265 In the present study, samples showed a high percentage of microplastics
266 with respect to zooplankton in abundance, especially in some areas such as
267 Taliarte, Las Canteras and Lambra. Microplastics reached values of 22% of
268 the zooplankton samples in Taliarte, this could explain the high incidence
269 of microplastics in the planktivorous fish Atlantic chub mackerel (*Scomber*
270 *colias*) collected in the Canary Islands according to the study carry out by
271 Herrera et al. (2019b). This Micro/Zoo ratio in abundance is similar to the
272 ones found by Moore et al. (2001) in the North Pacific Central Gyre and Frias
273 et al. (2014) on the Portuguese coast, and much higher than that reported
274 by other authors (see table 2).

275

276 In addition, the dry weight ratio for the 1-5mm fraction in the Las Can-
277 teras area showed twice times much microplastics as zooplankton. Collignon
278 et al. (2012) found an average weight ratio of 0.5 and Moore et al. (2001)
279 found 6 times more plastic than zooplankton in the area near the accumu-
280 lation of the North Pacific Subtropical Gyre. Although the ratio is higher,
281 Moore et al. (2001) included the fraction greater than 5mm, whereas in the
282 present study the ratio of 2 was found taking into account only the fraction
283 of 1-5mm.

284

285 This high percentage of microplastics in the zooplankton samples could
286 have a great impact on the health of marine organisms, either because of the
287 physical danger of ingestion, the associated chemical contaminants or the
288 false sense of satiation that could affect the intake of nutrients, especially in
289 large filter feeders species. Fossi et al. (2017) demonstrated the overlap of
290 zones of microplastic accumulation with the feeding areas of fin whales in the
291 Pelagos Sanctuary in the Mediterranean, highlighting the high intake risk for
292 marine biota.

293

294 One of the main concerns of the scientific community is the effects that
295 microplastics can have on marine organisms and the food chain. Many stud-
296 ies have demonstrated the ingestion of microplastics in invertebrates, fish,
297 seabirds and cetaceans. However, the risk associated with this ingestion is
298 still unknown. On the other hand, microplastics have been shown to possess
299 various types of associated chemical contaminants (Endo et al., 2005; Hirai
300 et al., 2011; Ogata et al., 2009; Rios et al., 2007; Camacho et al., 2019) and
301 these could affect the health of organisms (Rochman et al., 2013; Derraik,
302 2002; Teuten et al., 2009).

303

304 Also, the high microplastics-zooplankton ratio found in this study demon-
305 strates the potential risk it may have for biota and marine ecosystems, espe-
306 cially if we consider that high levels of POP's and emerging chemical pollu-
307 tants have already been reported by Camacho et al. (2019) in microplastic
308 samples collected in the Canary Islands. The waters around Macaronesia are
309 important feeding grounds for some large filter feeders, such as the whale

310 shark (*Rhincodon typus*), the basking shark (*Cetorhinus maximus*), several
311 species of manta rays of the genus *Mobula* (*M. tarapacana*, *M. mobular*, *M.*
312 *birostris*); and filter whales of the genus *Balaenoptera* (*B. edeni*, *B. bryde*, *B.*
313 *physalus*, *B. borealis*, *B. musculus*) (Carrillo et al., 2010; Espino et al., 2014;
314 Sobral and Afonso, 2014; Das and Afonso, 2017; Prieto et al., 2014, 2017;
315 Silva et al., 2014). According to our results, these species among others have
316 a high potential risk of ingestion of microplastics and associated chemical
317 contaminants.

318 5. Conclusions

- 319 1- High levels of contamination by neustonic microplastics (0.2-5mm)
320 were found in various areas of Macaronesia, reaching values of more than 1
321 million particles per square kilometre.
- 322 2- The microplastics-zooplankton abundance ratio was very variable in the
323 different zones, reaching values of 0.22.
- 324 3- We found twice as much microplastics as zooplankton in dry weight for
325 the 1-5mm fraction in the area of greatest accumulation in Las Canteras.
- 326 4- It is necessary to carry out more studies of floating microplastics abun-
327 dance in order to understand the circulation and accumulation patterns in
328 the Macaronesian region.
- 329 5- In addition, studies on the abundance of neustonic microplastics and zoo-
330 plankton and their impact at different levels of the food web are needed to
331 assess possible risks to marine organisms.

332

333 Acknowledgements

334 This work was funded by projects PLASMAR (MAC/1.1a/030), with
335 the support of the European Union (EU) and co-financed by the Euro-
336 pean Regional Development Fund (ERDF) and the INTERREG V-A Spain-
337 Portugal MAC 2014-2020 (Madeira-Azores-Canarias) and MICROTROFIC
338 (ULPGC2015-04) awarded to A.H. Part of this work was performed in the
339 framework of the project LIXAZ (ACORES-01-0145-FEDER-00053), funded
340 through the operational program AÇORES 2020 and also was partially funded
341 by Fundação para a Ciência e Tecnologia, I.P. (FCT), Portugal, with national
342 funds (FCT/MCTES, orçamento de Estado, project reference PTDC/MAR-
343 PRO/1851/2014), and the European Regional Development Fund (ERDF)
344 through the COMPETE 2020 programme (POCI-01-0145-FEDER-016885)
345 through the project PLASTICGLOBAL Assessment of plastic-mediated chem-
346 icals transfer in food webs of deep, coastal and estuarine ecosystems under
347 global change scenarios. The project PLASTICGLOBAL is also funded by
348 the Lisboa 2020 programme (LISBOA-01-0145-FEDER-016885).

349 A.H. was supported by a postdoctoral fellowship granted by Universidad
350 de Las Palmas de Gran Canaria (ULPGC-2014). S.A. was supported by a
351 Research fellowship for Graduates granted by Agência Regional para o De-
352 senvolvimento da Investigação, Tecnologia e Inovação (ARDITI). I.G. was
353 financially supported by a post-doctoral grant in the framework of the 2015
354 ARDITI Grant Programme Madeira 14-20 (Project M1420-09-5369-FSE-
355 000001). J. C.-C. was supported by a starting grant in the framework of
356 the 2014 FCT Investigator Programme (IF/01606/2014/CP1230/CT0001).
357 This study had the support of Fundação para a Ciência e Tecnologia (FCT),

358 through the strategic project UID/MAR/04292/2019 granted to MARE.
359 We would like to thank Captain Jorge Cáceres for his support during the
360 sampling in La Graciosa Marine Reserve, and Carmen Meléndez from the
361 Farfalle Project for her collaboration in the collection of samples at Los Gi-
362 gantes.

363

364 6. References

365 Aliabad, M.K., Nassiri, M., Kor, K., 2019. Microplastics in
366 the surface seawaters of Chabahar Bay , Gulf of Oman (
367 Makran Coasts). Marine Pollution Bulletin 143, 125–133.
368 URL: <https://doi.org/10.1016/j.marpolbul.2019.04.037>,
369 doi:10.1016/j.marpolbul.2019.04.037.

370 Álvarez-Hernández, C., Cairós, C., López-Darias, J., Mazzetti, E.,
371 Hernández-Sánchez, C., González-Sálamo, J., Hernández-Borges,
372 J., 2019. Microplastic debris in beaches of Tenerife (Ca-
373 nary Islands , Spain). Marine Pollution Bulletin 146, 26–
374 32. URL: <https://doi.org/10.1016/j.marpolbul.2019.05.064>,
375 doi:10.1016/j.marpolbul.2019.05.064.

376 Arthur, C., Baker, J., Bamford, H., 2009. Proceedings of the International
377 Research Workshop on the Occurrence , Effects , and Fate of Microplastic
378 Marine Debris. Group , 530.

379 Aytan, U., Valente, A., Senturk, Y., Usta, R., Basak, F., Sahin, E.,
380 Evren, R., Agirbas, E., 2016. First evaluation of neustonic mi-

croplastics in Black Sea waters. *Marine Environmental Research* 119,
 22–30. URL: <http://dx.doi.org/10.1016/j.marenvres.2016.05.009>,
 doi:10.1016/j.marenvres.2016.05.009.

Barrows, A.P.W., Neumann, C.A., Berger, L., Shaw, S.D., 2017. Grab
 vs. neuston tow net: a microplastic sampling performance comparison
 and possible advances in the field. *Anal. Methods* 9, 1446–1453.
 doi:10.1039/C6AY02387H.

Baztan, J., Carrasco, A., Chouinard, O., Cleaud, M., Gabaldon, J.E.,
 Huck, T., Jaffrès, L., Jorgensen, B., Miguelez, A., Paillard, C.,
 Vanderlinden, J.P., 2014. Protected areas in the Atlantic facing
 the hazards of micro-plastic pollution: First diagnosis of three is-
 lands in the Canary Current. *Marine Pollution Bulletin* 80, 302–
 311. URL: <http://dx.doi.org/10.1016/j.marpolbul.2013.12.052>,
 doi:10.1016/j.marpolbul.2013.12.052.

Beer, S., Garm, A., Huwer, B., Dierking, J., Gissel, T., 2018. Sci-
 ence of the Total Environment No increase in marine microplas-
 tic concentration over the last three decades A case study from
 the Baltic Sea. *Science of the Total Environment* 621, 1272–
 1279. URL: <https://doi.org/10.1016/j.scitotenv.2017.10.101>,
 doi:10.1016/j.scitotenv.2017.10.101.

Camacho, M., Herrera, A., Gómez, M., Acosta-Dacal, A., Henríquez-
 Hernández, L.A., Martínez, I., Pérez-Luzardo, O., 2019. Or-
 ganic pollutants in marine plastic debris from Canary Is-
 lands beaches. *Science of the Total Environment* 662, 22–

31. URL: <https://doi.org/10.1016/j.scitotenv.2018.12.422>,
doi:10.1016/j.scitotenv.2018.12.422.
- Carrillo, M., Pérez-Vallazza, C., Álvarez-Vázquez, R., 2010. Cetacean diversity and distribution off Tenerife (Canary Islands). *Marine Biodiversity Records*, 3, 1–9. doi:10.1017/S1755267210000801.
- Chambault, P., Vandeperre, F., Machete, M., Carvalho, J., Kim, C., 2018. Distribution and composition of floating macro litter off the Azores archipelago and Madeira (NE Atlantic) using opportunistic surveys. *Marine Environmental Research* 141, 225–232. URL: <https://doi.org/10.1016/j.marenvres.2018.09.015>,
doi:10.1016/j.marenvres.2018.09.015.
- Cheung, L.T., Lui, C.Y., Fok, L., 2018. Microplastic contamination of wild and captive flathead grey mullet (*Mugil cephalus*). *International Journal of Environmental Research and Public Health* 15. doi:10.3390/ijerph15040597.
- Collignon, A., Hecq, J.H., Galgani, F., Collard, F., Goffart, A., 2014. Annual variation in neustonic micro- and meso-plastic particles and zooplankton in the Bay of Calvi (Mediterranean-Corsica). *Marine Pollution Bulletin* 79, 293–298. URL: <http://dx.doi.org/10.1016/j.marpolbul.2013.11.023>,
doi:10.1016/j.marpolbul.2013.11.023.
- Collignon, A., Hecq, J.H., Galgani, F., Voisin, P., Collard, F., Goffart, A., 2012. Neustonic microplastic and zooplankton in the North

428 Western Mediterranean Sea. Marine Pollution Bulletin 64, 861–
 429 864. URL: <http://dx.doi.org/10.1016/j.marpolbul.2012.01.011>,
 430 doi:10.1016/j.marpolbul.2012.01.011.

431 Comas-Rodríguez, I., 2011. The Azores Current System and the Canary
 432 Current from CTD and ADCP data. Ph.D. thesis. Universidad de Las
 433 Palmas de Gran Canaria. URL: <http://hdl.handle.net/10553/6250>,
 434 doi:978-84-694-9558-2.

435 Cózar, A., Sanz-Martín, M., Martí, E., González-gordillo, J.I., Ubeda, B.,
 436 Gálvez, J.Á., Irigoien, X., Duarte, C.M., 2015. Plastic Accumulation in
 437 the Mediterranean. PLoS ONE , 1–12doi:10.1371/journal.pone.0121762.

438 Das, D., Afonso, P., 2017. Review of the Diversity , Ecology , and Con-
 439 servation of Elasmobranchs in the Azores Region , Mid-North Atlantic.
 440 Frontiers in Marine Science 4, 1–19. doi:10.3389/fmars.2017.00354.

441 De Lucia, A.G., Caliani, I., Marra, S., Camedda, A., Coppa, S., Alcaro, L.,
 442 Campani, T., Giannetti, M., Coppola, D., Maria, A., Panti, C., Baini,
 443 M., Guerranti, C., Marsili, L., Massaro, G., Cristina, M., Matiddi, M.,
 444 2014. Amount and distribution of neustonic micro-plastic off the western
 445 Sardinian coast (Central-Western Mediterranean Sea). Marine Environ-
 446 mental Research 100, 10–16. doi:10.1016/j.marenvres.2014.03.017.

447 Derraik, J.G.B., 2002. The pollution of the marine environment by plas-
 448 tic debris. Marine Pollution Bulletin 44, 842–852. doi:10.1016/s0025-
 449 326x(02)00220-5.

- 450 Di Mauro, R., Kupchik, M.J., Ben, M.C., 2017. Abundant plankton-sized
451 microplastic particles in shelf waters of the northern Gulf of Mexico *.
452 Environmental Pollution 230, 798–809. doi:10.1016/j.envpol.2017.07.030.
- 453 Doyle, M.J., Watson, W., Bowlin, N.M., Sheavly, S.B., 2011.
454 Plastic particles in coastal pelagic ecosystems of the North-
455 east Pacific ocean. Marine Environmental Research 71, 41–52.
456 URL: <http://dx.doi.org/10.1016/j.marenvres.2010.10.001>,
457 doi:10.1016/j.marenvres.2010.10.001.
- 458 Endo, S., Takizawa, R., Okuda, K., Takada, H., Chiba, K., Kanehiro, H.,
459 Ogi, H., Yamashita, R., Date, T., 2005. Concentration of polychlorinated
460 biphenyls (PCBs) in beached resin pellets: Variability among individual
461 particles and regional differences. Marine Pollution Bulletin 50, 1103–1114.
462 doi:10.1016/j.marpolbul.2005.04.030.
- 463 Eriksen, M., Liboiron, M., Kiessling, T., Charron, L., Alling, A., Le-
464 breton, L., Richards, H., Roth, B., Ory, N.C., Hidalgo-Ruz, V.,
465 Meerhoff, E., Box, C., Cummins, A., Thiel, M., 2018. Microplas-
466 tic sampling with the AVANI trawl compared to two neuston trawls
467 in the Bay of Bengal and South Pacific *. Environmental Pollution
468 232, 430–439. URL: <https://doi.org/10.1016/j.envpol.2017.09.058>,
469 doi:10.1016/j.envpol.2017.09.058.
- 470 Eriksen, M., Maximenko, N., Thiel, M., Cummins, A., Lattin, G., Wilson,
471 S., Hafner, J., Zellers, A., Rifman, S., 2013. Plastic pollution in the South
472 Pacific subtropical gyre. Marine Pollution Bulletin 68, 71–76.

- 473 Espino, F., González, J., Boyra, A., Fernández, C., Tuya, F., Brito, A., 2014.
474 Diversity and biogeography of fishes in the Arinaga-Gando area, east coast
475 of Gran Canaria (Canary Islands). *Rev. Acad. Canar. Cienc.* XXVI, 9–25.
- 476 Faure, F., Demars, A.C., Wieser, A.O., Kunz, A.M., Felipe De Alen-
477 castro, L., 2015. Plastic pollution in Swiss surface waters: nature and
478 concentrations, interaction with pollutants. *Environmental Chemistry*
479 doi:10.1071/EN14218.
- 480 Figueiredo, G.M., Moraes, T., Vianna, P., Bay, G., 2018. Sus-
481 pended microplastics in a highly polluted bay : Abundance , size ,
482 and availability for mesozooplankton. *Marine Pollution Bulletin* 135,
483 256–265. URL: <https://doi.org/10.1016/j.marpolbul.2018.07.020>,
484 doi:10.1016/j.marpolbul.2018.07.020.
- 485 Fossi, M.C., Romero, T., Bains, M., Panti, C., Marsili, L., Campani, T.,
486 Canese, S., Galgani, F., Druon, J.N., Airolidi, S., Taddei, S., Fattorini,
487 M., Brandini, C., Lapucci, C., 2017. Plastic Debris Occurrence , Conver-
488 gence Areas and Fin Whales Feeding Ground in the Mediterranean Marine
489 Protected Area Pelagos Sanctuary : A Modeling Approach. *Frontiers in*
490 *Marine Science* 4, 1–15. doi:10.3389/fmars.2017.00167.
- 491 Frère, L., Paul-Pont, I., Rinnert, E., Petton, S., Jaffré, J., Bihannic,
492 I., Soudant, P., Lambert, C., Huvet, A., 2017. Influence of envi-
493 ronmental and anthropogenic factors on the composition, concentra-
494 tion and spatial distribution of microplastics: A case study of the Bay
495 of Brest (Brittany, France). *Environmental Pollution* 225, 211–222.
496 doi:10.1016/j.envpol.2017.03.023.

- 497 Frias, J.P.G.L., Otero, V., Sobral, P., 2014. Evidence of microplastics in
498 samples of zooplankton from Portuguese coastal waters. *Marine Environ-
499 mental Research* 95, 89–95. doi:10.1016/j.marenvres.2014.01.001.
- 500 Gago, J., Henry, M., Galgani, F., 2015. First observation on
501 neustonic plastics in waters off NW Spain (spring 2013 and
502 2014). *Marine environmental research* 111, 27–33. URL:
503 <http://www.sciencedirect.com/science/article/pii/S0141113615300167>,
504 doi:10.1016/j.marenvres.2015.07.009.
- 505 Gestoso, I., Cacabelos, E., Ramalhosa, P., Canning-clode, J., 2019.
506 Plasticrusts : A new potential threat in the Anthropocene ' s
507 rocky shores. *Science of the Total Environment* 687, 413–415.
508 doi:10.1016/j.scitotenv.2019.06.123.
- 509 Gewert, B., Ogonowski, M., Barth, A., Macleod, M., 2017. Abundance
510 and composition of near surface microplastics and plastic debris in the
511 Stockholm Archipelago , Baltic Sea. *Marine Pollution Bulletin* 120, 292–
512 302. URL: <http://dx.doi.org/10.1016/j.marpolbul.2017.04.062>,
513 doi:10.1016/j.marpolbul.2017.04.062.
- 514 Gilfillan, L.R., Doyle, M., Ohman, M.D., Watson, W., 2009. Occurrence of
515 plastic micro-debris in the southern California Current system. Technical
516 Report Vol 50 CalCOFI Rep. Scripps Institution of Oceanography.
- 517 Green, D.S., Kregting, L., Boots, B., Blockley, D.J., Brickle, P., Crow-
518 ley, Q., 2018. A comparison of sampling methods for seawater mi-
519 croplastics and a fi rst report of the microplastic litter in coastal wa-

ters of Ascension and Falkland Islands. *Marine Pollution Bulletin* 137,
695–701. URL: <https://doi.org/10.1016/j.marpolbul.2018.11.004>,
doi:10.1016/j.marpolbul.2018.11.004.

Hartmann, N.B., Hu, T., Thompson, R.C., Hassello, M., Verschoor, A., Dau-
gaard, A.E., Rist, S., Karlsson, T., Brennholt, N., Cole, M., Herrling,
M.P., Hess, M.C., Ivleva, N.P., Lusher, A.L., Wagner, M., 2019. Are We
Speaking the Same Language ? Recommendations for a Definition and
Categorization Framework for Plastic Debris. *Environ. Sci. Technol* 53,
1039–1047. doi:10.1021/acs.est.8b05297.

Herrera, A., Asensio, M., Martínez, I., Santana, A., Packard, T., Gómez,
M., 2018. Microplastic and tar pollution on three Canary Islands
beaches: An annual study. *Marine Pollution Bulletin* 129, 494–502.
doi:10.1016/j.marpolbul.2017.10.020.

Herrera, A., Martínez, I., Rapp, J., Raymond, E., Álvarez, S., Gestoso, I.,
Canning-Clode, J., Gómez, M., 2019a. Sampling and Processing Microplas-
tics from Surface Waters. Technical Report. PLASMAR Project. URL:
<http://hdl.handle.net/10553/56209>.

Herrera, A., Stindlová, A., Martínez, I., Rapp, J., Romero-Kutzner, V.,
Samper, M.D., Montoto, T., Aguiar-González, B., Packard, T., Gómez,
M., 2019b. Microplastic ingestion by Atlantic chub mackerel (*Scomber*
colias) in the Canary Islands coast. *Marine Pollution Bulletin* 139,
127–135. URL: <https://doi.org/10.1016/j.marpolbul.2018.12.022>,
doi:10.1016/j.marpolbul.2018.12.022.

543 Hirai, H., Takada, H., Ogata, Y., Yamashita, R., Mizukawa, K., Saha,
 544 M., Kwan, C., Moore, C., Gray, H., Laursen, D., Zettler, E.R., Far-
 545 rington, J.W., Reddy, C.M., Peacock, E.E., Ward, M.W., 2011. Or-
 546 ganic micropollutants in marine plastics debris from the open ocean
 547 and remote and urban beaches. *Marine Pollution Bulletin* 62, 1683–
 548 1692. URL: <http://dx.doi.org/10.1016/j.marpolbul.2011.06.004>,
 549 doi:10.1016/j.marpolbul.2011.06.004.

550 Isobe, A., Uchida, K., Tokai, T., Iwasaki, S., 2015. East Asian seas : A
 551 hot spot of pelagic microplastics. *Marine Pollution Bulletin* 101, 618–623.
 552 doi:10.1016/j.marpolbul.2015.10.042.

553 Isobe, A., Uchiyama-Matsumoto, K., Uchida, K., Tokai, T., 2017.
 554 Microplastics in the Southern Ocean. *MPB* 114, 623–626.
 555 URL: <http://dx.doi.org/10.1016/j.marpolbul.2016.09.037>,
 556 doi:10.1016/j.marpolbul.2016.09.037.

557 Ivar do Sul, J.A., Costa, M.F., 2014. Microplastics in the pelagic environment
 558 around oceanic islands of the Western Tropical Atlantic Ocean. *Water, Air,*
 559 *and Soil Pollution* doi:10.1007/s11270-014-2004-z.

560 Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M.,
 561 Andrady, A., Narayan, R., Law, K.L., 2015. Plastic waste in-
 562 puts from land into the ocean. *Science* 347, 768–771. URL:
 563 <http://www.sciencemag.org/cgi/doi/10.1126/science.1260352>,
 564 doi:10.1126/science.1260352.

565 Kang, J.h., Youn, O., Lee, K.w., Kyoung, Y., Joon, W.,

2015. Marine neustonic microplastics around the southeast-
ern coast of Korea. Marine Pollution Bulletin 96, 304–312.
URL: <http://dx.doi.org/10.1016/j.marpolbul.2015.04.054>,
doi:10.1016/j.marpolbul.2015.04.054.
- Kanhai, L.D.K., Officer, R., Lyashevskaya, O., Thompson, R.C., O'Connor,
I., 2017. Microplastic abundance, distribution and composition along a
latitudinal gradient in the Atlantic Ocean. Marine Pollution Bulletin 115,
307–314. doi:10.1016/j.marpolbul.2016.12.025.
- Kroon, F., Motti, C., Talbot, S., Sobral, P., Puotinen, M., 2018. A work-
flow for improving estimates of microplastic contamination in marine wa-
ters : A case study from North-Western Australia A work fl ow for im-
proving estimates of microplastic contamination in marine waters : A
case study from North-Western Australia *. Environmental Pollution
238, 26–38. URL: <https://doi.org/10.1016/j.envpol.2018.03.010>,
doi:10.1016/j.envpol.2018.03.010.
- Kukulka, T., Proskurowski, G., Meyer, D.W., Law, K.L., 2012. The effect
of wind mixing on the vertical distribution of buoyant plastic debris. Geo-
physical Research Letters 39, 1–6. doi:10.1029/2012GL051116.
- Lattin, G.L., Moore, C.J., Zellers, A.F., Moore, S.L., Weisberg, S.B., 2004.
A comparison of neustonic plastic and zooplankton at different depths
near the southern California shore. Marine Pollution Bulletin 49, 291–294.
doi:10.1016/j.marpolbul.2004.01.020.
- Law, K.L., More, S.E., Goodwin, D.S., Zettler, E.R., 2014. Distribution of

589 Surface Plastic Debris in the Eastern Pacific Ocean from an 11-Year Data
590 Set. *Environmental Science & Technology* doi:10.1021/es4053076.

591 Lima, A., Costa, M.F., Barletta, M., 2014. Distribu-
592 tion patterns of microplastics within the plankton of a
593 tropical estuary. *Environmental Research* 132, 146–155.
594 URL: <http://dx.doi.org/10.1016/j.envres.2014.03.031>,
595 doi:10.1016/j.envres.2014.03.031.

596 Lusher, A.L., Burke, A., Connor, I.O., Officer, R., 2014. Microplastic pollu-
597 tion in the Northeast Atlantic Ocean : Validated and opportunistic sam-
598 pling. *Marine Pollution Bulletin* .

599 Lusher, A.L., Tirelli, V., Connor, I.O., Officer, R., 2015. Microplastics in
600 Arctic polar waters : the first reported values of particles in surface and
601 sub-surface samples. *Scientific Reports* , 1–9doi:10.1038/srep14947.

602 Maes, T., Meulen, M.D.V.D., Devriese, L.I., Leslie, H.A., Huvet, A., Frère,
603 L., Robbens, J., Vethaak, A.D., 2017. Microplastics Baseline Surveys at
604 the Water Surface and in Sediments of the North-East Atlantic. *Frontiers*
605 *in Marine Science* 4, 1–13. doi:10.3389/fmars.2017.00135.

606 Mcknight, L., 2016. Nearshore circulation in the Confital bay Implications
607 on marine debris transport and deposition at Las Canteras Beach. Master
608 thesis. Universidad de Las Palmas de Gran Canaria.

609 Moore, C.J., Moore, S.L., Leecaster, M.K., Weisberg, S.B., 2001. A com-
610 parison of plastic and plankton in the North Pacific Central Gyre. *Marine*
611 *Pollution Bulletin* 42, 1297–1300. doi:10.1016/S0025-326X(01)00114-X.

- 612 Moore, C.J., Moore, S.L., Weisberg, S.B., Lattin, G.L., Zellers, a.F., 2002.
 613 A comparison of neustonic plastic and zooplankton abundance in south-
 614 ern California’s coastal waters. *Marine Pollution Bulletin* 44, 1035–1038.
 615 doi:10.1016/S0025-326X(02)00150-9.
- 616 MSDF Technical Subgroup on Marine Litter, 2013. Guidance on Mon-
 617 itoring of Marine Litter in European Seas. Technical Report. Euro-
 618 pean Commission. URL: [papers3://publication/doi/10.2788/99475](https://publications.europa.eu/en/publication-detail/-/publication/10.2788/99475),
 619 doi:10.2788/99475.
- 620 Ogata, Y., Takada, H., Mizukawa, K., Hirai, H., Iwasa, S., Endo, S.,
 621 Mato, Y., Saha, M., Okuda, K., Nakashima, A., Murakami, M., Zurcher,
 622 N., Booyatumanondo, R., Zakaria, M.P., Dung, L.Q., Gordon, M.,
 623 Miguez, C., Suzuki, S., Moore, C., Karapanagioti, H.K., Weerts, S.,
 624 McClurg, T., Burres, E., Smith, W., Van Velkenburg, M., Lang, J.S.,
 625 Lang, R.C., Laursen, D., Danner, B., Stewardson, N., Thompson,
 626 R.C., 2009. International Pellet Watch: global monitoring of persistent
 627 organic pollutants (POPs) in coastal waters. 1. Initial phase data on
 628 PCBs, DDTs, and HCHs. *Marine pollution bulletin* 58, 1437–46. URL:
 629 <http://www.sciencedirect.com/science/article/pii/S0025326X09002690>,
 630 doi:10.1016/j.marpolbul.2009.06.014.
- 631 Pedrotti, M.L., Petit, S., Elineau, A., Bruzard, S., 2016. Changes in the
 632 Floating Plastic Pollution of the Mediterranean Sea in Relation to the
 633 Distance to Land. *PLoS ONE* 11, 1–14. doi:10.1371/journal.pone.0161581.
- 634 Pham, C.K., Ramirez-llodra, E., Alt, C.H.S., Amaro, T., Bergmann, M.,
 635 Canals, M., Company, J.B., Davies, J., Duineveld, G., Howell, K.L.,

636 Huvenne, V.A.I., Isidro, E., Jones, D.O.B., Lastras, G., Gomes-pereira,
 637 N., Purser, A., Stewart, H., Morato, T., Tubau, X., Rooij, D.V.,
 638 Tyler, P.A., 2014. Marine Litter Distribution and Density in Euro-
 639 pean Seas , from the Shelves to Deep Basins. PLoS ONE 9, 1–13.
 640 doi:10.1371/journal.pone.0095839.

641 Pham, C.K., Rodríguez, Y., Dauphin, A., Carriço, R., Frias,
 642 J.P.G.L., Vandeperre, F., Otero, V., Santos, M.R., Martins, H.R.,
 643 Bolten, A.B., Bjorndal, K.A., 2017. Plastic ingestion in oceanic-
 644 stage loggerhead sea turtles (*Caretta caretta*) off the North At-
 645 lantic subtropical gyre. Marine Pollution Bulletin 121, 222–
 646 229. URL: <http://dx.doi.org/10.1016/j.marpolbul.2017.06.008>,
 647 doi:10.1016/j.marpolbul.2017.06.008.

648 Picheral, M., Colin, S., Irisson, J.O., 2017. EcoTaxa, a tool for the taxonomic
 649 classification of images.

650 PlasticsEurope, 2018. Plastics the Facts 2018. Technical Re-
 651 port. Plastics Europe-Association of Plastics Manufacturers. URL:
 652 www.plasticseurope.org.

653 Prieto, R., Silva, M.A., Waring, G.T., Gonçalves, J.M.A., 2014.
 654 Sei whale movements and behaviour in the North Atlantic inferred
 655 from satellite telemetry. Endangered Species Research 26, 103–113.
 656 doi:10.3354/esr00630.

657 Prieto, R., Tobeña, M., Silva, M.A., 2017. Habitat preferences of
 658 baleen whales in a mid-latitude habitat. Deep-Sea Research Part II

141, 155–167. URL: <http://dx.doi.org/10.1016/j.dsr2.2016.07.015>,
doi:10.1016/j.dsr2.2016.07.015.

R Core Team, 2019. R: A Language and Environment for Statistical Computing. URL: <https://www.r-project.org/>.

Rios, L.M., Moore, C., Jones, P.R., 2007. Persistent organic pollutants carried by synthetic polymers in the ocean environment. *Marine Pollution Bulletin* 54, 1230–1237. doi:10.1016/j.marpolbul.2007.03.022.

Ríos, N., Frias, J.P.G.L., Rodríguez, Y., Carriço, R., Garcia, M., 2018. Spatio-temporal variability of beached macro-litter on remote islands of the North Atlantic. *Marine Pollution Bulletin* 133, 304–311.

Rochman, C.M., Hoh, E., Kurobe, T., Teh, S.J., 2013. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Scientific Reports* 3, 3263. URL: <http://www.nature.com/articles/srep03263>, doi:10.1038/srep03263.

Rodríguez, A., Rodríguez, B., Nazaret Carrasco, M., 2012. High prevalence of parental delivery of plastic debris in Cory’s shearwaters (*Calonectris diomedea*). *Marine Pollution Bulletin* 64, 2219–2223. doi:10.1016/j.marpolbul.2012.06.011.

Rodríguez, Y., Pham, C.K., 2017. Marine litter on the sea floor of the Faial-Pico Passage , Azores Archipelago. *Marine Pollution Bulletin* 116, 448–453. URL: <http://dx.doi.org/10.1016/j.marpolbul.2017.01.018>, doi:10.1016/j.marpolbul.2017.01.018.

- 681 Ruiz-Orejón, L.F., Sard, R., Ramis-pujol, J., 2016. Floating plastic debris
682 in the Central and Western Mediterranean Sea. *Marine Environmental*
683 *Research* 120, 136–144. doi:10.1016/j.marenvres.2016.08.001.
- 684 Silva, M.A., Prieto, R., Cascão, I., Seabra, M.I., Machete, M., Baumgartner,
685 M.F., Santos, R.S., Silva, M.A., Prieto, R., Cascão, I., Seabra, M.I., Ma-
686 chete, M., 2014. Spatial and temporal distribution of cetaceans in the
687 mid-Atlantic waters around the Azores. *Marine Biology Research* 10,
688 123–137. URL: <http://dx.doi.org/10.1080/17451000.2013.793814>,
689 doi:10.1080/17451000.2013.793814.
- 690 Sobral, A.F., Afonso, P., 2014. Occurrence of mobulids in the Azores , central
691 North Atlantic. *Journal of the Marine Biological Association of the United*
692 *Kingdom* 94, 1671–1675. doi:10.1017/S0025315414000964.
- 693 Song, Y.K., Hong, S.H., Jang, M., Kang, J.H., Kwon, O.Y., Han, G.M.,
694 Shim, W.J., 2014. Large accumulation of micro-sized synthetic polymer
695 particles in the sea surface microlayer. *Environmental Science and Tech-*
696 *nology* 48, 9014–9021. doi:10.1021/es501757s.
- 697 Suaria, G., Avio, C.G., Mineo, A., Lattin, G.L., Magaldi, M.G., Belmonte,
698 G., Moore, C.J., Regoli, F., Aliani, S., 2016. The Mediterranean Plastic
699 Soup : synthetic polymers in Mediterranean surface waters. *Scientific*
700 *Reports* , 1–10doi:10.1038/srep37551.
- 701 Sundseth, K., 2010. Natura 2000 en la región macaronésica. Technical Re-
702 port. Unión Europea. Luxemburgo. doi:10.2779/70730.

703 Teuten, E.L., Saquing, J.M., Knappe, D.R.U., Barlaz, M.a., Jonsson, S.,
704 Björn, A., Rowland, S.J., Thompson, R.C., Galloway, T.S., Yamashita, R.,
705 Ochi, D., Watanuki, Y., Moore, C., Viet, P.H., Tana, T.S., Prudente, M.,
706 Boonyatumanond, R., Zakaria, M.P., Akkhavong, K., Ogata, Y., Hirai, H.,
707 Iwasa, S., Mizukawa, K., Hagino, Y., Imamura, A., Saha, M., Takada, H.,
708 2009. Transport and release of chemicals from plastics to the environment
709 and to wildlife. *Philosophical transactions of the Royal Society of London.*
710 *Series B, Biological sciences* 364, 2027–2045. doi:10.1098/rstb.2008.0284.

711 UNEP, 2009. Marine Litter : A Global Challenge Marine Litter : A Global
712 Challenge. Technical Report. United Nations Environment Programme.

713 Van Der Hal, N., Ariel, A., Angel, D.L., 2017. Exception-
714 ally high abundances of microplastics in the oligotrophic Israeli
715 Mediterranean coastal waters. *Marine Pollution Bulletin* 116, 151–
716 155. URL: <http://dx.doi.org/10.1016/j.marpolbul.2016.12.052>,
717 doi:10.1016/j.marpolbul.2016.12.052.

718 **7. Figures and Tables**



Figure 1: Study area. The numbers inside the circles show the number of samples collected at each site.




Figure 2: Types of debris found. a) Irregular fragments (Fragments), scale bar=5mm. b) Industrial pellets (Pellets), scale bar=5mm. c) Fibres, scale bar=2mm d) Microbeads, scale bar=500 μ m. e) EPS and XPS (Foams), scale bar=1mm. f) Films, scale bar=5mm. g,h) Fishing lines (Lines), scale bar=5mm. i) Paint (Other), scale bar=2mm.




Figure 3: (a) Abundance of microplastics (0.2-5mm) in items by Km² at each location. The central thick line of each box designates the median, the box height shows the interquartile range, and the whiskers indicate the lowest and the highest values. Significant differences between locations within each archipelago are shown ** ($p<0.05$), * ($p<0.01$).




Figure 3: (b) Neustonic zooplankton in individuals by Km² at each location. Y axis was log2 transformed in order to improve data visualization. Significant differences between locations within each archipelago are shown ** (p<0.05), * (p<0.01).




Figure 3: (c) Ratio Microplastics/Zooplankton abundance. Significant differences between locations within each archipelago are shown ** (p<0.05), * (p<0.01).




Figure 4: (a) Abundance of microplastics in items/Km² in coastal waters of Lanzarote and La Graciosa Islands, Canary Islands archipelago




Figure 4: (b) Abundance of microplastics in items/Km² in coastal waters of Gran Canaria Island, Canary Islands archipelago




Figure 4: (c) Abundance of microplastics in items/ Km^2 in coastal waters of Tenerife Island, Canary Islands archipelago




Figure 4: (d) Abundance of microplastics in items/ Km^2 in coastal waters of Madeira Island, Madeira archipelago.







Figure 4: (e) Abundance of microplastics in items/Km² in coastal waters of Faial Island, Azores archipelago.



(a)




(b)



(c)

40

Figure 5: Percentage of type and size of debris found at each archipelago. (a) Canary Islands archipelago. (b) Madeira archipelago. (c) Azores archipelago. Category “Other” include glass, paint, aluminum foil and tar



(a)



(b)



(c)

Figure 6: Percentage of taxonomic groups from total neustonic zooplankton collected at each archipelago. (a) Canary Islands archipelago. (b) Madeira archipelago. (c) Azores archipelago.

Table 1: Mean abundance of microplastics and zooplankton, and ratio of items of microplastics/number of zooplankton at each sampling location.

Location	Archipelago	Micro (items/Km ²)	Zoo (ind/Km ²)	Micro/Zoo
		mean±SD	mean±SD(x10 ⁶)	items ratio
Lambra	Canary Islands	153,304±95,348	5.7±2.2	0.032
Arrecife	Canary Islands	157,102±96,840	5.1±1.9	0.030
Famara	Canary Islands	68,020±75,654	38.0±31.2	0.002
Taliarte	Canary Islands	154,570±9,217	1.4±1.1	0.147
Las Canteras	Canary Islands	894,069±98,951	15.7±13.1	0.08
Gando	Canary Islands	125,949±61,630	22.6±18.4	0.008
San Andres	Canary Islands	21,326±6,281	3.1±0.9	0.007
Los Gigantes	Canary Islands	27,593 ±8,895	14.4±4.1	0.002
Canical	Madeira	87,538±12,223	5.3±4.0	0.028
Funchal	Madeira	40,054 ±4,711	9.5±12.5	0.013
Desertas	Madeira	66,568±19,379	7.3±4.8	0.021
Canico	Madeira	84,343±39,828	5.1±3.2	0.024
Praia do Norte	Azores	77,223±40,279	140.7±75.2	0.0007
Almoxarife	Azores	143,858±143,033	95.0±32.4	0.002
Porto Pim	Azores	300,352±164,345	177.9±98.7	0.002

Table 2: Bibliographic search in Web of Science by terms *debris and *neustonic from 1900 to 2019 showed 48 results. From these results we selected the articles that report data of neustonic microplastics or debris in the oceans.

Sampling area	Net mesh size μm	Mean MP items/ Km^2	Max MP items/ Km^2	Mean MP items/ m^3	Max MP items/ m^3	MP/zoo ratio items	Reference
Chabahar Bay, Gulf of Oman	neuston 333			0.49	1.14		Aliabad et al. (2019)
Black Sea	WP2 200			0.0012			Aytan et al. (2016)
Bornholm Basin, Baltic Sea	Bongo 150			0.21	0.28		Beer et al. (2018)
Pearl River estuary, Hong Kong waters	manta 333	334,780	1,675,982	3.97	29697		Cheung et al. (2018)
North Western Mediterranean Sea	manta 333	116,000	892,000				Collignon et al. (2012)
Bay of Calvi, Mediterranean Sea	manta 333	62,000	688,000			0.0006	Collignon et al. (2014)
Sardinian Sea, Western Mediterranean	manta 500			0.15	0.35		De Lucia et al. (2014)
Northern Gulf of Mexico	neuston 335			13	21.6	0.0058	Di Mauro et al. (2017)
Northeast Bering Sea, Pacific Ocean	Sameoto 505			0.017-0.072	0.072		Doyle et al. (2011)
Southern California, Pacific Ocean	Manta 505			0.004-0.19	0.19		Doyle et al. (2011)
South Pacific subtropical gyre	Manta 333	26,988	396,342				Eriksen et al. (2013)
Western Mediterranean Sea	Manta 333	129,682	420,000				Faure et al. (2015)
Guanabara Bay, Southeastern Brazil	neuston 64	900,000	1,900,000	4.8	11		Figueiredo et al. (2018)
Pelagos Sanctuary, Western Mediterranean Sea	mantan 333	82,000	264,000				Fossi et al. (2017)
Bay of Brest, France	manta 335			0.24	1.43		Frère et al. (2017)
Portuguese coastal waters, Aveiro	neuston 280-335			0.002		0.04	Frias et al. (2014)
Portuguese coastal waters, Lisboa	neuston 280-335			0.033		0.12	Frias et al. (2014)
Portuguese coastal waters, Costa Vicentina	neuston 280-335			0.036		0.14	Frias et al. (2014)
Portuguese coastal waters, Algarve	neuston 280-335			0.014		0.005	Frias et al. (2014)
Spanish northwest coast	manta 335	34,000					Gago et al. (2015)
Spanish northwest coast	manta 335	176,000					Gago et al. (2015)

Continued on next page

Sampling area	Net	Mean MP		Max MP		Mean MP		Max MP		MP/zoo ratio items	Reference
		mesh size μm	items/ Km^2	items/ Km^2	items/ Km^2	items/ m^3	items/ m^3	items/ m^3	items/ m^3		
Stockholm Archipelago, Baltic Sea	manta 335		110,000	618,000		1.37		7.73			Gewert et al. (2017)
Southern California Current, Pacific Ocean	manta 505					0.011					Gilfillan et al. (2009)
Southern California Current, Pacific Ocean	manta 505					0.033					Gilfillan et al. (2009)
Southern California Current, Pacific Ocean	manta 505					0.016					Gilfillan et al. (2009)
East Asian Seas, Japan	neuston 350			172,000		3.74		491			Isobe et al. (2015)
Southern Ocean, Antarctica	manta 350		100,000			0.031		0.099			Isobe et al. (2017)
Western Tropical Atlantic, Abrolhos	manta 300					0.04					Ivar do Sul and Costa (2014)
Western Tropical Atlantic, Fernando de Noronha	manta 300					0.015					Ivar do Sul and Costa (2014)
Western Tropical Atlantic, Trinidad	manta 300					0.025		0.13			Ivar do Sul and Costa (2014)
Western Tropical Atlantic Ocean	manta 300					0.03					Ivar do Sul and Costa (2014)
Atlantic Ocean	pump 250					1.15		8.5			Kanghai et al. (2017)
South East Sea of Korea	manta 330					1.92-5.51					Kang et al. (2015)
South East Sea of Korea	manta 330					2.30-38.77					Kang et al. (2015)
North Western Australia, Indian Ocean	manta 355					0.01-0.41					Kroon et al. (2018)
North Western Australia, Indian Ocean	plankton					0.00-0.09					Kroon et al. (2018)
Santa Monica Bay, California	manta 333					3.92					Latini et al. (2004)
Eastern Pacific Ocean (accumulation zone)	neuston 333		156,800	12,340,000							Law et al. (2014)
Eastern Pacific Ocean (outside)	neuston 333		1,864								Law et al. (2014)
Goiana Estuary, Northeast coast of Brazil	plankton 300										Lima et al. (2014)
Northeast Atlantic Ocean	pump 250					0.26		0.0019			Lusher et al. (2014)
Arctic waters, Norway	manta 333		28,000			2.46		22.5			Lusher et al. (2015)
Arctic waters, Norway	pump 250					0.34		1.31			Lusher et al. (2015)
North-East Atlantic	manta 333		36,623	375,854		2.68		11.5			Lusher et al. (2015)
North Pacific Central Gyre	manta 330		334,271	969,777		0.15		1.5			Maes et al. (2017)
						2.23		0.1819			Moore et al. (2001)

Continued on next page

Sampling area	Net	Mean MP items/Km ²	Max MP items/Km ²	Mean MP items/m ³	Max MP items/m ³	MP/zoo ratio items	Reference
Southern California coastal waters	manta 333			7.25			Moore et al. (2002)
Mediterranean Sea, near coast	manta 333	158,000	578,000			0.03	Pedrotti et al. (2016)
Mediterranean Sea, > 10Km from coast	manta 333	370,000				0.006	Pedrotti et al. (2016)
Central and Western Mediterranean Sea	manta 333	147,500	1,164,403				Ruiz-Orejón et al. (2016)
Southern coast Korea	manta 333			43			Song et al. (2014)
Mediterranean Sea	neuston 200	400,000	4,520,000	1	11.3		Suaria et al. (2016)
Israeli Mediterranean coast	manta 333	1,518,384	64,812,600	7.68	324		Van Der Hal et al. (2017)
Mediterranean Sea	manta 200	24,3853					Cózar et al. (2015)
North Atlantic Ocean, Azores	manta 200	173,811	467,260	0.44	1.19	0.002	present work
North Atlantic Ocean, Madeira	manta 200	69,626	124,190			0.021	present work
North Atlantic Ocean, Canary Islands	manta 200	194,951	1,007,872			0.032	present work

Supplementary material

Table 3: Sampling dates (mm/dd/yy), locations, distances to the coast in meters and type of net used.

Location	Island	Archipelago	Date	Longitud	Latitud	Distance	Net
Canical	Madeira	Madeira	8/11/17	-16.7084	32.7143	2503	Manta (200 μ m)
Canical	Madeira	Madeira	8/11/17	-16.7317	32.7132	2995	Manta (200 μ m)
Canical	Madeira	Madeira	8/11/17	-16.7496	32.6984	2332	Manta (200 μ m)
Funchal	Madeira	Madeira	8/11/17	-16.8393	32.6305	915	Manta (200 μ m)
Funchal	Madeira	Madeira	8/11/17	-16.8624	32.6329	1017	Manta (200 μ m)
Funchal	Madeira	Madeira	8/11/17	-16.8736	32.6156	1073	Manta (200 μ m)
Desertas	Madeira	Madeira	8/12/17	-16.6819	32.6751	5563	Manta (200 μ m)
Desertas	Madeira	Madeira	8/12/17	-16.7076	32.6666	7497	Manta (200 μ m)
Desertas	Madeira	Madeira	8/12/17	-16.7310	32.6573	6610	Manta (200 μ m)
Canico	Madeira	Madeira	8/12/17	-16.7577	32.6780	2470	Manta (200 μ m)
Canico	Madeira	Madeira	8/12/17	-16.7710	32.6623	2251	Manta (200 μ m)
Canico	Madeira	Madeira	8/12/17	-16.7884	32.6482	2723	Manta (200 μ m)
Lambra	La Graciosa	Canary Islands	11/6/15	-13.4883	29.2962	1680	Manta (200 μ m)
Lambra	La Graciosa	Canary Islands	11/6/15	-13.5303	29.3297	2245	Manta (200 μ m)
Lambra	La Graciosa	Canary Islands	11/6/15	-13.4489	29.2623	2287	Manta (200 μ m)
Arrecife	Lanzarote	Canary Islands	12/3/15	-13.5263	28.9606	834	Manta (200 μ m)
Arrecife	Lanzarote	Canary Islands	12/3/15	-13.5411	28.9553	659	Manta (200 μ m)
Arrecife	Lanzarote	Canary Islands	12/3/15	-13.5201	28.9549	1774	Manta (200 μ m)
Famara	Lanzarote	Canary Islands	3/4/16	-13.5434	29.1479	1682	Manta (200 μ m)
Famara	Lanzarote	Canary Islands	3/4/16	-13.5734	29.1372	978	Manta (200 μ m)
Famara	Lanzarote	Canary Islands	3/4/16	-13.5852	29.1533	2103	Manta (200 μ m)
Taliarte	Gran Canaria	Canary Islands	6/8/18	-15.3666	28.0140	856	Manta (200 μ m)
Taliarte	Gran Canaria	Canary Islands	6/8/18	-15.3622	27.9897	520	Manta (200 μ m)
Las Canteras	Gran Canaria	Canary Islands	6/26/18	-15.4689	28.1327	395	Manta (200 μ m)
Las Canteras	Gran Canaria	Canary Islands	6/26/18	-15.4533	28.1355	745	Manta (200 μ m)
Las Canteras	Gran Canaria	Canary Islands	6/26/18	-15.4402	28.1469	730	Manta (200 μ m)
Gando	Gran Canaria	Canary Islands	9/18/18	-15.3733	27.9511	713	Manta (200 μ m)
Gando	Gran Canaria	Canary Islands	9/18/18	-15.3736	27.9657	510	Manta (200 μ m)
Gando	Gran Canaria	Canary Islands	9/18/18	-15.3688	27.9797	648	Manta (200 μ m)
San Andres	Gran Canaria	Canary Islands	10/3/18	-15.5415	28.1583	915	Manta (200 μ m)
San Andres	Gran Canaria	Canary Islands	10/3/18	-15.5235	28.1593	710	Manta (200 μ m)
San Andres	Gran Canaria	Canary Islands	10/3/18	-15.5047	28.1607	1446	Manta (200 μ m)

Continued on next page

Location	Island	Archipelago	Date	Longitud	Latitud	Distance	Net
Los Gigantes	Tenerife	Canary Islands	10/4/18	-16.8506	28.2377	1107	Manta (200 μ m)
Los Gigantes	Tenerife	Canary Islands	10/4/18	-16.8511	28.2224	1171	Manta (200 μ m)
Los Gigantes	Tenerife	Canary Islands	10/4/18	-16.8463	28.2065	1184	Manta (200 μ m)
Los Gigantes	Tenerife	Canary Islands	10/4/18	-16.8465	28.2479	1178	Manta (200 μ m)
Praia do Norte	Faial	Azores	7/10/18	-28.7570	38.6132	172	Bongo (200 μ m)
Praia do Norte	Faial	Azores	7/10/18	-28.7571	38.6131	308	Bongo (200 μ m)
Praia do Norte	Faial	Azores	7/10/18	-28.7559	38.6139	303	Bongo (200 μ m)
Almoxarife	Faial	Azores	7/10/18	-28.6076	38.5559	116	Bongo (200 μ m)
Almoxarife	Faial	Azores	7/10/18	-28.6079	38.5547	139	Bongo (200 μ m)
Almoxarife	Faial	Azores	7/10/18	-28.6078	38.5546	133	Bongo (200 μ m)
Porto Pim	Faial	Azores	7/10/18	-28.6290	38.5241	78	Bongo (200 μ m)
Porto Pim	Faial	Azores	7/10/18	-28.6286	38.5231	101	Bongo (200 μ m)
Porto Pim	Faial	Azores	7/10/18	-28.6289	38.5240	83	Bongo (200 μ m)